

# Welcome & Roadmap

Manqi Ruan


# Physics Benchmarks

Benchmarks	Main observables	Key performances	Status
llH, H->X	Higgs recoil spectrum	Lepton Id efficiency, Tracker intrinsic momentum resolution	Well understood
H+X, H->di photon	Event reconstruction efficiency, Higgs invariant mass peak width	Tracker Material, Intrinsic ECAL energy Resolution	
ZH->4 jets,	Br(H->bb, cc, gg)	Jet clustering, PFA: Jet Energy Resolution, Jet Flavor Tagging	Studied at CEPC conceptual Detector (CEPC_v1)
vvH, H->di tau	Efficiency of Tau reconstruction with different tau decay mode	PFA separation, Impact parameter resolution	
qqH, H->invisible	Higgs recoil spectrum	PFA: Jet Energy Resolution	
vvH, H->WW->lvqq	Event Reconstruction Efficiency di-jet mass distribution	PFA, Simultaneous reconstruction of Lepton, Jets and Missing Energy	Studied at different Calorimeter Granularity
H+X, H->di muon	Event reconstruction efficiency, Higgs invariant mass peak width	Lepton Id efficiency, Tracker intrinsic momentum resolution	Studied at CEPC conceptual Detector
vvH, H->2 jets	Br(H->bb, cc, gg)	Jet Energy Resolution & Flavor Tagging	
WW->lvqq	W mass	Jet Energy resolution & Systematic controls	Full simulation analysis not accomplished yet

Each analysis will be repeated at different geometry, with full Higgs signal sample  
(and potentially WW sample)

# Goal

- Toward
  - The Physics Benchmark
    - Physics + EW
  - The Detector Key Performance & Key Reconstruction Technology
  - The Key Detector Geometry Design & Parameter
  - Focus on Critical questions: potential No-Go problems
  - The Theoretical Errors
  - The CDR & TDR



Back up

# Tasks ahead

- Detector Geometry Design

- ILD-like:

- Key parameter selections
    - Key performance
    - Key questions (Potentially No-Go)


- Significant Beyond ILD: any good idea?

- Systematic errors...

- Software Chain: A continuous effort of 3-5 years.

- Physics Potential




- Higgs                      Mostly done. Benchmarks need be iterated at new geometry
  - EW, Z                      ??
  - Flavor,                      ??
  - ... theoretical errors                      Progressing



# Fix Basic Geometry

- Benchmark detector geometry
  - ECAL Layout Done
  - HCAL Layout ~ 2 months
  - Tracker Size
    - for tracking performance Done
    - PFA Performance ~ 2.5 months
  - B-Field Strength ~ 1 months
  - Integrated with a reliable MDI ???
- ~ half a year is needed to fix the basic geometry of the detector, with a full support from the analysis team.
- Very unfortunate, many key players will leave us soon

# Detector performance



Acceptance	$ \cos(\theta)  < 0.995$ (from the inner radius of the outmost tracking disk)
Tracking Efficiency	For isolated charged particle with energy $> 1\text{ GeV}$ : $\sim 100\%$
Photon Reconstruction Efficiency	For isolated photon with energy $> 0.5\text{ GeV}$ : $\sim 100\%$
Tracker resolution	$\delta(1/P_T) = 2 \cdot 10^{-5} (\text{GeV}^{-1})$
ECAL intrinsic resolution	$\delta E/E = 16\%/\sqrt{E/\text{GeV}} \oplus 0.5\%$
HCAL intrinsic resolution	$\delta E/E = 60\%/\sqrt{E/\text{GeV}} \oplus 1\%$
Jet energy resolution	$\delta E/E = 4\%$
<b>Typical Distance for shower separation</b>	$< 3\text{ cm}$
<b>Lepton identification</b>	For charged particle with Energy $> 2\text{ GeV}$ : Lepton identification efficiency $> 99.5\%$ , $P(\text{hadron} \rightarrow \text{muon}) \sim P(\text{hadron} \rightarrow \text{electron})$ : 1%
b-tagging	At Z pole samples & $\text{eff}(b \rightarrow b) = 80\%$ , $P(\text{uds} \rightarrow b) < 1\%$ , $P(c \rightarrow b) \sim 10\%$
c-tagging	At Z pole samples & $\text{eff}(c \rightarrow c) = 60\%$ , $P(\text{uds} \rightarrow c) = 7\%$ , $P(b \rightarrow c) = 12\%$

Performance at full reconstruction

*Benchmark separation distance  $< 3\text{ cm}$   
(Testing on 10 GeV Pion + 5 GeV Photon Sample)*