

Implication of Higgs Factory Precision Measurements on New Physics Models



Shufang Su • U. of Arizona

IHEP

Nov 2, 2017

S. Su

Collaboration with J. Gu, H. Li, Z. Liu, W. Su.


Outline

- ➊ Higgs precision measurements
- ➋ Global fit framework
- ➌ Perturbative models
 - SM with a real singlet extension
 - 2HDM
- ➍ Strong dynamics
 - Minimal composite Higgs Model (MCHM)
 - General EFT patterns of strongly interacting models
- ➎ Conclusion


Higgs Precision Measurements

ATLAS

Individual analysis




LHC: 7+8 TeV



Higgs Precision Measurements

ATLAS Simulation Preliminary

$\sqrt{s} = 14 \text{ TeV}$: $\int L dt = 300 \text{ fb}^{-1}$; $\int L dt = 3000 \text{ fb}^{-1}$



ATL-PHYS-PUB-2014-016

$\Delta\mu/\mu$

LHC: 14 TeV, 300 fb⁻¹, 3000 fb⁻¹

$\Delta\mu/\mu$	300 fb^{-1}		3000 fb^{-1}		
	All unc.	No theory unc.	All unc.	No theory unc.	
$H \rightarrow \gamma\gamma$ (comb.)	0.13	0.09	0.09	0.04	
	(0j)	0.19	0.12	0.16	0.05
	(1j)	0.27	0.14	0.23	0.05
	(VBF-like)	0.47	0.43	0.22	0.15
	(WH-like)	0.48	0.48	0.19	0.17
	(ZH-like)	0.85	0.85	0.28	0.27
	(ttH-like)	0.38	0.36	0.17	0.12
$H \rightarrow ZZ$ (comb.)	0.11	0.07	0.09	0.04	
	(VH-like)	0.35	0.34	0.13	0.12
	(ttH-like)	0.49	0.48	0.20	0.16
	(VBF-like)	0.36	0.33	0.21	0.16
	(ggF-like)	0.12	0.07	0.11	0.04
	(0j)	0.18	0.09	0.16	0.05
	(1j)	0.30	0.18	0.26	0.10
$H \rightarrow WW$ (comb.)	0.13	0.08	0.11	0.05	
	(0j)	0.18	0.09	0.16	0.05
	(1j)	0.30	0.18	0.26	0.10
	(VBF-like)	0.21	0.20	0.15	0.09
	(ggF-like)	0.12	0.07	0.11	0.04
	(0j)	0.18	0.09	0.16	0.05
	(1j)	0.30	0.18	0.26	0.10
$H \rightarrow Z\gamma$ (incl.)	0.46	0.44	0.30	0.27	
	$H \rightarrow b\bar{b}$ (comb.)				
	(WH-like)	0.26	0.26	0.14	0.12
	(ZH-like)	0.57	0.56	0.37	0.36
	(ttH-like)	0.29	0.29	0.14	0.13
	(0j)	0.21	0.18	0.19	0.15
	(1j)	0.47	0.45	0.48	0.44
$H \rightarrow \tau\tau$ (VBF-like)	0.21	0.18	0.19	0.15	
	$H \rightarrow \mu\mu$ (comb.)				
	(incl.)	0.39	0.38	0.16	0.12
	(ttH-like)	0.47	0.45	0.48	0.44
	(0j)	0.21	0.18	0.19	0.15
	(1j)	0.74	0.72	0.27	0.23
	(VBF-like)	0.21	0.18	0.19	0.15

Higgs Precision Measurements

CEPC / FCC / ILC


collider	CEPC	FCC-ee	ILC					
\sqrt{s}	240 GeV	240 GeV	250 GeV	350 GeV		500 GeV		
$\int \mathcal{L} dt$	5 ab $^{-1}$	10 ab $^{-1}$	2 ab $^{-1}$	200 fb $^{-1}$		4 ab $^{-1}$		
production	Zh	Zh	Zh	Zh	$\nu\bar{\nu}h$	Zh	$\nu\bar{\nu}h$	$t\bar{t}h$
$\Delta\sigma/\sigma$	0.51%	0.4%	0.71%	2.1%	-	1.06	-	-
decay	$\Delta(\sigma \cdot BR)/(\sigma \cdot BR)$							
$h \rightarrow b\bar{b}$	0.28%	0.2%	0.42%	1.67%	1.67%	0.64%	0.25%	9.9%
$h \rightarrow c\bar{c}$	2.2%	1.2%	2.9%	12.7%	16.7%	4.5%	2.2%	-
$h \rightarrow gg$	1.6%	1.4%	2.5%	9.4%	11.0%	3.9%	1.5%	-
$h \rightarrow WW^*$	1.5%	0.9%	1.1%	8.7%	6.4%	3.3%	0.85%	-
$h \rightarrow \tau^+\tau^-$	1.2%	0.7%	2.3%	4.5%	24.4%	1.9%	3.2%	-
$h \rightarrow ZZ^*$	4.3%	3.1%	6.7%	28.3%	21.8%	8.8%	2.9%	-
$h \rightarrow \gamma\gamma$	9.0%	3.0%	12.0%	43.7%	50.1%	12.0%	6.7%	-
$h \rightarrow \mu^+\mu^-$	17%	13%	25.5%	97.6%	179.8%	31.1%	25.5%	-
$(\nu\bar{\nu})h \rightarrow b\bar{b}$	2.8%	2.2%	3.7%	-	-	-	-	-

Kappa framework and EFT Framework

Two model-independent approaches


kappa framework

$$\kappa_f = \frac{g(hff)}{g(hff; \text{SM})}, \quad \kappa_V = \frac{g(hVV)}{g(hff; \text{SM})}$$




EFT framework

$$\delta c_Z, \quad c_{ZZ}, \quad c_{Z\square}, \quad c_{\gamma\gamma}, \quad c_{Z\gamma}, \quad c_{gg}, \quad \delta y_u, \quad \delta y_d, \quad \delta y_e, \quad \lambda_Z$$



1704.02333

New Physics Implication




Kappa Framework and EFT Framework


limitations of model-independent approaches

- large level of degeneracy
parameter space for specific model much smaller
- correlation matrix often not provided
over conservative estimation when not include correlation
- assumptions and simplifications
may not be valid for a particular model


New Physics Implication



New Physics Implication



New Physics Implication



$$\chi^2 = \sum_i \frac{(\mu_i^{\text{BSM}} - \mu_i^{\text{obs}})^2}{\sigma_{\mu_i}^2} \quad \mu_i^{\text{BSM}} = \frac{(\sigma \times \text{Br})_{\text{BSM}}}{(\sigma \times \text{Br})_{\text{SM}}}$$

Perturbative Models

- SM with a real singlet extension
- 2HDM (Type I, II, L, F)

SM + Real Scalar Singlet

- SM + real scalar singlet

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \frac{1}{2}(\partial_\mu S)^2 - \frac{1}{2}m_S^2 S^2 - \Lambda_{SH} S(H^\dagger H) - \frac{1}{2}\lambda_{SH} S^2(H^\dagger H) - \frac{1}{3!}\Lambda_S S^3 - \frac{1}{4!}\lambda_S S^4$$


- after EWSB, 2 physical Higgses: CP-even Higgses: h_{SM} , singlet S
- Z_2 breaking: mixing between h_{SM} and S

$$h_{125} = \cos \theta \ h_{\text{SM}} + \sin \theta \ S$$

$$\kappa_i = g_i^{\text{SM+singlet}} / g_i^{SM} = \cos \theta$$


SM + Real Scalar Singlet

○ fit to $\sin \theta$



SM + Real Scalar Singlet

○ fit to $\sin \theta$




SM + Real Scalar Singlet

- fit to c_6 and c_H

$$\Delta\mathcal{L} = \frac{c_H}{\Lambda^2}\mathcal{O}_H + \frac{c_6}{\Lambda^2}\mathcal{O}_6$$

$$\mathcal{O}_H \equiv \frac{1}{2}(\partial_\mu|H^\dagger H|)^2 \quad \mathcal{O}_6 \equiv |H^\dagger H|^3$$

$$1 - \cos \theta \simeq \theta^2/2 \simeq \\ 1/2 \times c_H v^2/\Lambda^2$$



Perturbative Models

- SM with a real singlet extension
- 2HDM (Type I, II, L, F)

2HDM in one slide

- Two Higgs Doublet Model (CP-conserving)

$$\Phi_i = \begin{pmatrix} \phi_i^+ \\ (v_i + \phi_i^0 + iG_i)/\sqrt{2} \end{pmatrix}$$

$$v_u^2 + v_d^2 = v^2 = (246\text{GeV})^2$$
$$\tan \beta = v_u/v_d$$

$$\begin{pmatrix} H^0 \\ h^0 \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \phi_1^0 \\ \phi_2^0 \end{pmatrix}, \quad A^0 = -G_1 \sin \beta + G_2 \cos \beta$$
$$H^\pm = -\phi_1^\pm \sin \beta + \phi_2^\pm \cos \beta$$

after EWSB, 5 physical Higgses

CP-even Higgses: h^0, H^0 , CP-odd Higgs: A^0 , Charged Higgses: H^\pm

- h^0/H^0 VV coupling

$$g_{H^0VV} = \frac{m_V^2}{v} \cos(\beta - \alpha), \quad g_{h^0VV} = \frac{m_V^2}{v} \sin(\beta - \alpha).$$


alignment limit: $\cos(\beta-\alpha)=0$, h^0 is the SM Higgs with SM couplings.

2HDM parameters

	Φ_1	Φ_2
Type I	u,d,l	
Type II	u	d,l
lepton-specific	u,d	l
flipped	u,l	d



Model	κ_V	κ_u	κ_d	κ_ℓ
2HDM-I	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$\cos \alpha / \sin \beta$	$\cos \alpha / \sin \beta$
2HDM-II	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$-\sin \alpha / \cos \beta$	$-\sin \alpha / \cos \beta$
2HDM-L	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$\cos \alpha / \sin \beta$	$-\sin \alpha / \cos \beta$
2HDM-F	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$-\sin \alpha / \cos \beta$	$\cos \alpha / \sin \beta$

- parameters (CP-conserving, flavor limit, Z_2 symmetry)





Tree-level 2HDM fit

2HDM, LHC/CEPC fit




Tree-level 2HDM fit

2HDM, LHC/CEPC fit




Tree-level 2HDM fit

- κ -fit vs $\Delta\mu/\mu$ fit,
- CEPC/FCC/ILC,
- luminosity dependence



2HDM Model Distinction

Model	κ_V	κ_u	κ_d	κ_ℓ
2HDM-I	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$\cos \alpha / \sin \beta$	$\cos \alpha / \sin \beta$
2HDM-II	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$-\sin \alpha / \cos \beta$	$-\sin \alpha / \cos \beta$
2HDM-L	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$\cos \alpha / \sin \beta$	$-\sin \alpha / \cos \beta$
2HDM-F	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$-\sin \alpha / \cos \beta$	$\cos \alpha / \sin \beta$




2HDM: Loop in the Alignment Limit

• Type I


$$\kappa_{\text{loop}}^{\text{2HDM}} \equiv \frac{g_{\text{tree}}^{\text{2HDM}} + g_{\text{loop}}^{\text{2HDM}}}{g_{\text{tree}}^{\text{SM}} + g_{\text{loop}}^{\text{SM}}}$$

$$\kappa_{1-\text{loop}}^{\text{2HDM}}|_{\text{alignment}} = 1 + \Delta \kappa_{1-\text{loop}}^{\text{2HDM}}$$





2HDM: Loop in the Alignment Limit

• Type II




2HDM: Loop in the Alignment Limit

◎ Type II, varying luminosity



2HDM: Loop in the Alignment Limit


◎ Type II, κ -fit vs $\Delta\mu/\mu$ fit, CEPC/FCC/ILC



2HDM: Loop in the Alignment Limit


◎ Type II, varying m_{12}

$$\lambda v^2 \equiv m_\phi^2 - m_{12}^2 / (\sin \beta \cos \beta)$$



2HDM: Loop in the Alignment Limit

◎ 2HDM: distinguish different type



Strong Dynamics

- Minimum composite Higgs Model (**MCHM**)
- General EFT patterns of strong interacting models with a light Higgs

Composite Higgs in one slide

- Higgs is the PNGB of the spontaneous breaking of $G \Rightarrow H$
- EWSB is induced by vacuum misalignment, parametrized by $\xi = v^2/f^2$
- mass of SM fermion generated by mixing with composite states
- light top partners can be searched at the LHC
- minimal composite Higgs Model (MCHM): $SO(5)/SO(4)$

- hVV

$$\kappa_V \equiv \frac{g_{hVV}^{\text{CH}}}{g_{hVV}^{\text{SM}}} = \sqrt{1 - \xi}$$

- hff : depends on the fermion representation

$$F_1 \equiv \frac{1 - 2\xi}{\sqrt{1 - \xi}}, \quad F_2 \equiv \sqrt{1 - \xi}$$



MCHM

○ Fermion representation

MCHM: $\xi = v^2/f^2 < 10^{-3}$, $f > 4$ TeV

MCHM Reps.	5, 10 14-1-10 14-10-10 10-14-10	10-5-10	5-5-10	5-10-10 5-1-10	14-14-10	14-5-10	5-14-10
κ_t, κ_g	F_1	F_2	F_1	F_2	F_3	F_4	F_5
κ_b	F_1	F_1	F_2	F_2	F_1	F_1	F_1
CEPC							
$\xi \times 10^3$	2.56	2.36	4.19	3.87	2.78 – 2.56	2.71 – 2.36	2.36 – 2.04
f [TeV]	4.86	5.06	3.80	3.95	4.67 – 4.86	4.72 – 5.07	5.07 – 5.45
ILC							
$\xi \times 10^3$	2.19	2.02	3.44	3.20	2.31 – 2.19	2.06 – 2.01	1.87 – 1.72
f [TeV]	5.26	5.48	4.19	4.35	5.12 – 5.26	5.42 – 5.48	5.69 – 5.93
FCC-ee							
$\xi \times 10^3$	1.80	1.66	3.06	2.74	1.85 – 1.80	1.70 – 1.66	1.66 – 1.41
f [TeV]	5.79	6.04	4.45	4.70	5.72 – 5.80	5.97 – 6.05	6.05 – 6.56

MCHM



Strong Dynamics in EFT Language


① EFT operators

$$\mathcal{L}_6 = \frac{1}{m_*^2} \sum_i c_i \mathcal{O}_i$$

$\mathcal{O}_H = \frac{1}{2}(\partial_\mu H^2)^2$	$\mathcal{O}_{GG} = g_s^2 H ^2 G_{\mu\nu}^A G^{A,\mu\nu}$
$\mathcal{O}_W = \frac{ig}{2}(H^\dagger \sigma^a \overleftrightarrow{D}^\mu H) D^\nu W_{\mu\nu}^a$	$\mathcal{O}_{Y_u} = Y_u H ^2 \bar{Q}_L \tilde{H} u_R$
$\mathcal{O}_B = \frac{ig'}{2}(H^\dagger \overleftrightarrow{D}^\mu H) \partial^\nu B_{\mu\nu}$	$\mathcal{O}_{Y_d} = Y_d H ^2 \bar{Q}_L H d_R$
$\mathcal{O}_{HW} = ig(D^\mu H)^\dagger \sigma^a (D^\nu H) W_{\mu\nu}^a$	$\mathcal{O}_{Y_e} = Y_e H ^2 \bar{L}_L H e_R$
$\mathcal{O}_{HB} = ig'(D^\mu H)^\dagger (D^\nu H) B_{\mu\nu}$	$\mathcal{O}_{3W} = \frac{1}{3!} g \epsilon_{abc} W_\mu^{a\nu} W_{\nu\rho}^b W^{c\rho\mu}$
$\mathcal{O}_{BB} = g'^2 H ^2 B_{\mu\nu} B^{\mu\nu}$	


	\mathcal{O}_H	\mathcal{O}_W	\mathcal{O}_B	\mathcal{O}_{HW}	\mathcal{O}_{HB}	\mathcal{O}_{BB}	\mathcal{O}_{GG}	\mathcal{O}_{y_u}	\mathcal{O}_{y_d}	\mathcal{O}_{y_e}	\mathcal{O}_{3W}
ALH	g_*^2	1	1	1	1	1	1	g_*^2	g_*^2	g_*^2	$\frac{g^2}{g_*^2}$
GSILH	g_*^2	1	1	1	1	$\frac{y_t^2}{16\pi^2}$	$\frac{y_t^2}{16\pi^2}$	g_*^2	g_*^2	g_*^2	$\frac{g^2}{g_*^2}$
SILH	g_*^2	1	1	$\frac{g_*^2}{16\pi^2}$	$\frac{g_*^2}{16\pi^2}$	$\frac{y_t^2}{16\pi^2}$	$\frac{y_t^2}{16\pi^2}$	g_*^2	g_*^2	g_*^2	$\frac{g^2}{16\pi^2}$

Strong Dynamics in EFT Language




Strong Dynamics in EFT Language

individual fit




global fit



Conclusion

- future Higgs factories measure Higgs properties to a high precision
- Kappa-scheme/EFT scheme/model specific fit
- perturbative model




Conclusion


- strong dynamics models

- MCHM: $\xi = v^2/f^2 < 10^{-3}$, $f > 4$ TeV
- ALH/GSILH/SILH

individual fit



global fit



Conclusion



LHC



Lepton Collider



100 TeV pp

An exciting journey ahead of us!