

Charm physics at BESIII

报告人：马海龙 (中科院高能所)

实验物理中心讲座 EPD seminar

November 29th 2017

Introduction

Leptonic and hadronic decays of charmed hadrons (D^0 , D^+ , D_s^+ and Λ_c^+) provide ideal test-beds to explore weak and strong effects

➤ D leptonic decays

$f_{D(s)^+}$, $f_{K(\pi)^+}(0)$: better calibrate LQCD

$|V_{cs(d)}|$: better test on CKM matrix unitarity

Search for lepton flavor violation


$$U = \begin{bmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{bmatrix}$$

➤ D hadronic decays

$D^0\bar{D}^0$ mixing parameters and CPV

Strong phase difference in D^0 decays:

Constrain γ/ϕ_3 measurement in B decays



➤ Absolute BF's of Λ_c^+ decays


No absolute BF measurements of Λ_c^+ using near threshold data before BESIII

Contents


- Charm samples at BESIII
- (Semi-)leptonic $D_{(s)}$ decays
- Hadronic $D_{(s)}$ decays
- Λ_c^+ decays
- Summary

Recent $D^{0(+)}$, D_s^+ and Λ_c^+ samples

Taking from Longke Li's talk at joint workshop of BESIII/Belle/LHCb at Nankai

Experiment	Machine	C.M	Lumin.	N(D)	efficiency	advantage/disadvantage
 	CESR (e^+e^-)	3.77 GeV	0.8 fb $^{-1}$	2.9×10^6 $2.3 \times 10^6 (D^\pm)$	~10-30%	☺ extremely clean environment ☺ pure D-beam, almost no bkg ☺ quantum coherence ☹ no CM boost, no T-dep analyses
		4.17 GeV	0.6 fb $^{-1}$	0.6×10^6		
	BEPC-II (e^+e^-)	3.77 GeV	2.92 fb $^{-1}$	10.5×10^6 8.4×10^6 $D^{0(+)}$		
		4.18 GeV	3 fb $^{-1}$	3×10^6 D_s^+ Λ_c^+		
		4.6 GeV	0.567 fb $^{-1}$	★	★★★	
 	KEKB (e^+e^-)	10.58 GeV	1 ab $^{-1}$	1.3×10^9	~5-10%	☺ clear event environment ☺ high trigger efficiency ☺ high-efficiency detection of neutrals ☺ many high-statistics control samples ☺ time-dependent analysis ☹ smaller cross-section than pp colliders
	PEP-II (e^+e^-)	10.58 GeV	0.5 ab $^{-1}$	6.5×10^8 ★★		
	Tevatron ($p\bar{p}$)	1.96 TeV	9.6 fb $^{-1}$	1.3×10^{11}		☺ large production cross-section ☺ large boost: excellent time resolution
	LHC (pp)	7 TeV	1.0 fb $^{-1}$	5.0×10^{12} ★★★★	<0.5% ★	☹ dedicated trigger required ☹ hard to do neutrals and neutrinos
		8 TeV	2.0 fb $^{-1}$			


BEPCII: high luminosity double-ring collider



BESIII detector

Magnet: 1 T Super conducting

Nucl. Instr. Meth. A614, 345 (2010)



MDC: small cell & Gas:
 He/C₃H₈ (60/40), 43 layers
 $\sigma_{xy} = 130 \mu\text{m}$
 $\sigma_p/p = 0.5\% @ 1\text{GeV}$
 $dE/dx = 6\%$

TOF:
 $\sigma_T = 100 \text{ ps}$ Barrel
 110 ps Endcap

Muon ID: 9 layers RPC
 8 layers for endcap



EMC: CsI crystal, 28 cm
 $\Delta E/E = 2.5\% @ 1 \text{ GeV}$
 $\sigma_z = 0.6 \text{ cm}/\sqrt{E}$

Data Acquisition:
 Event rate = 4 kHz
 Total data volume ~ 50 MB/s


**2016年, ETOF升级改造
 后, 分辨明显改善**

$D^{0(+)}$, D_s^+ and Λ_c^+ samples (pb^{-1}) at BESIII

➤ $D^{0(+)}$ samples at $\psi(3770)$



➤ $D_s^+/\bar{D}_s^+/\Lambda_c^+$ samples



$$N_{ST}^i = 2 \times N_{D\bar{D}} \times B_{ST}^i \times \epsilon_{ST}^i$$

$$N_{DT}^i = 2 \times N_{D\bar{D}} \times B_{ST}^i \times B_{sig} \times \epsilon_{ST \text{ vs. sig}}^i$$

$$B_{sig} = \frac{N_{DT}^{tot}}{N_{ST}^{tot} \times \bar{\epsilon}_{sig}}$$



$$\bar{\epsilon}_{sig} = \sum_{i=1}^N (N_{ST}^i \times \epsilon_{ST \text{ vs. sig}}^i / \epsilon_{ST}^i) / \sum_{i=1}^N N_{ST}^i$$

(Semi)leptonic $D_{(s)}$ decays

- $l^+\nu$
- $Pl^+\nu$
- $VL^+\nu$
- $Sl^+\nu$
- Rare SL decays

(Semi-)leptonic D decays

粲介子(半)轻子衰变是测定CKM矩阵元 $|V_{cs(d)}|$ 的理想桥梁



$$\Gamma(D_{(s)}^+ \rightarrow \ell^+ \nu_\ell) = \frac{G_F^2 f_{D_{(s)}^+}^2}{8\pi} |V_{cd(s)}|^2 m_\ell^2 m_{D_{(s)}^+} \left(1 - \frac{m_\ell^2}{m_{D_{(s)}^+}^2}\right)^2$$

$$\frac{d\Gamma}{dq^2} = X \frac{G_F^2 |V_{cd(s)}|^2}{24\pi^3} p^3 |f_+(q^2)|^2$$

- 改进的 $f_{D(s)+}$, $f_+^{D \rightarrow K(\pi)}(q^2)$ 能够在更高精度上检验格点QCD的计算
- 格点QCD计算精度的改善为精密测量 $|V_{cs(d)}|$ 创造了条件

Searches for $D_{(s)}^+ \rightarrow l^+ \nu$ at BES I/II

22.3 pb⁻¹ at 4.03 GeV




22.3 pb⁻¹ at 4.03 GeV

33 pb⁻¹ around $\psi(3770)$

PRL74(1995)4599

PLB429(1998)188

PLB610(2005)183



$$f_{D_s^+} = (430^{+150+40}_{-130-40}) \text{ MeV}$$

$$f_{D^+} = (300^{+180+80}_{-150-40}) \text{ MeV}$$

$$f_{D^+} = (371^{+129}_{-119} \pm 25) \text{ MeV}$$

BESI

BESI


BESII

Newest results on $B[D^+ \rightarrow \mu^+ \nu]$, $f_{D^+}|V_{cd}|$

818 pb⁻¹ at $\psi(3770)$
(2004–2008)

2.93 fb⁻¹ data@ 3.773 GeV


PRD78(2008)052003



$$B_{D^+ \rightarrow \mu^+ \nu} = (3.82 \pm 0.32 \pm 0.09) \times 10^{-4}$$

$$f_{D^+} = 205.8 \pm 7.5 \pm 2.5 \text{ MeV}$$


PRD89(2014)051104R



$$B_{D^+ \rightarrow \mu^+ \nu} = (3.71 \pm 0.19 \pm 0.06) \times 10^{-4}$$

$$f_{D^+} = (203.2 \pm 5.3 \pm 1.8) \text{ MeV}$$

$$|V_{cd}| = 0.2210 \pm 0.0058 \pm 0.0047$$




20 fb⁻¹ 数据能够将 f_{D^+}
统计误差降至1%

Evidence of $D^+ \rightarrow \tau^+ \nu$ (4σ)

11

Fitting to DATA



$$B[D^+ \rightarrow \tau^+ \nu] = (1.20 \pm 0.24_{\text{stat.}}) \times 10^{-3}$$

$$R \equiv \frac{\Gamma(D^+ \rightarrow \tau^+ \nu)}{\Gamma(D^+ \rightarrow \mu^+ \nu)} = \frac{m_{\tau^+}^2 \left(1 - \frac{m_{\tau^+}^2}{M_{D^+}^2}\right)^2}{m_{\mu^+}^2 \left(1 - \frac{m_{\mu^+}^2}{M_{D^+}^2}\right)^2}$$

SM prediction: 2.66

BESIII: 3.21±0.64


12

Previous measurements of $D_s^+ \rightarrow l^+ \nu$


In the past 30 years, $D_s^+ \rightarrow l^+ \nu$ has been studied by WA75, CLEOII, E653, BESII, L3, OPAL, ALPHA, **CLEO-c, BELLE, Babar**

■ $D_s^+ D_s^-$, 600 pb⁻¹
@ 4.17 GeV [697 $l^+ \nu$]

PRD79(2009)052001




$f_{D_s^+} = 263.3 \pm 8.2 \pm 1.9$ MeV



Signal for $D_s^+ \rightarrow \tau^+ \nu$

$f_{D_s^+} = 252.2 \pm 11.1 \pm 5.2$ MeV

PRD80(2009)112004




$f_{D_s^+} = 257.8 \pm 13.3 \pm 5.2$ MeV

■ Belle, 913 fb⁻¹ at
10.58 GeV [2698 $l^+ \nu$]

$$e^+ e^- \rightarrow DKXD_s^{*-}$$

JHEP1309(2013)129




$f_{D_s^+} = 255.5 \pm 4.2 \pm 5.1$ MeV

■ Babar, 521 fb⁻¹ at
10.58 GeV [1023 $l^+ \nu$]

$$e^+ e^- \rightarrow DKXD_s^{*-}$$

PRD82(2010)091103




$f_{D_s^+} = 258.6 \pm 6.4 \pm 7.5$ MeV

Results on $B[D_s^+ \rightarrow l^+ \nu]$, $f_{D_{S^+}}|V_{cs}|$ at BESIII


0.48 fb⁻¹ data@4.01 GeV

3.19 fb⁻¹ data@4.178 GeV

PRD94(2016)072004




Use μ counter to suppress background



$f_{D_{S^+}} = (241.0 \pm 16.3 \pm 6.6) \text{ MeV}$

$f_{D_S}|V_{cs}| = 242.5 \pm 3.5 \pm 3.7 \text{ MeV}$




BESIII $f_{D_{S^+}}$ 精度达2%，
联合 $\tau^+ \nu$ 研究，能够降
至1.5%水平

Previous measurements of $f_+^{D \rightarrow K(\pi)_+}(0) |V_{cs(d)}|$


In the past 30 years, studies of $D \rightarrow K(\pi) l^+ \nu$ were made by MARKIII, E691, CLEO, CLEOII, BESII, FOCUS, **BELLE**, **Babar** and **CLEO-c**

■ BELLE, 282 fb⁻¹ at 10.58 GeV




PRL97(2006)061804

■ 2004-2009, CLEO-c, 818 pb⁻¹ at ψ''




PRD80(2009)032005

■ Babar, 75 fb⁻¹ at 10.58 GeV



■ Babar, 347.2 fb⁻¹ at 10.58 GeV




Before 2010, the LQCD calculated $f_+^{D \rightarrow K(\pi)}(0)$ precision is at 10% level, thus limiting $|V_{cs(d)}|$ measurement

Much improved LQCD calculations

Taking from Aida X. El-Khadra's talk at Beauty2014


errors (in %) comparison: **FLAG-2 averages** vs. **new results**




review by C. Bouchard @ Lattice 2014

Impact of $f^{D \rightarrow K(\pi)^+}(q^2)$ on LQCD



BESIII, PRD92(2015)072012





$D^0 \rightarrow K^- e^+ \nu$





$D^0 \rightarrow \pi^- e^+ \nu$



Comparisons of $f_{D \rightarrow K(\pi)_+}(0)$ with LQCD





BESIII, PRD96(2017)012002



First and second errors are statistical and systematic

Improvements on $|V_{cs(d)}|$

- 方法 1 $f_{D(s)}|V_{cd(s)}| \longrightarrow |V_{cd(s)}|$
- 方法 2 $f_{D \rightarrow K(\pi)_+}(0)|V_{cs(d)}| \longrightarrow |V_{cs(d)}|$





Taken from PDG, and the SL method suffers about 2.4% error from LQCD



方法2受 $f_{+}^{D \rightarrow K(\pi)}(0)$ 格点计算精度限制 [2.4(4.4)%]

Weights of $|V_{cs(d)}|$

BESIII $|V_{cd}|$ 权重>50%




$D_s^+ \rightarrow \tau^+ \nu$ 研究完成后, BESIII $D_s^+ \rightarrow l^+ \nu$ 对 $|V_{cs}|$ 贡献的权重有望达到50%左右



LFU test in CS decay $D^{0(+)} \rightarrow \pi l^+ \nu$ at BESIII


Evidence of LFV at 4σ in

$$R(D^{(*)}) = \frac{B(B \rightarrow D^{(*)} \tau \nu)}{B(B \rightarrow D^{(*)} l \nu)}$$



Evidence of LFV at 2.6σ in FCNC decays $B^+ \rightarrow K^+ \mu^+ \mu^- / K^+ e^+ e^-$


$$R_K = \frac{\Gamma(\bar{B} \rightarrow \bar{K} \mu^+ \mu^-)}{\Gamma(\bar{B} \rightarrow \bar{K} e^+ e^-)} = 0.745^{+0.090}_{-0.074} \pm 0.036$$



$$R_{LU}^{0(+)} = \frac{B(D^{0(+)} \rightarrow \pi^{-(0)} \mu^+ \nu)}{B(D^{0(+)} \rightarrow \pi^{-(0)} e^+ \nu)} \sim 0.97$$

B^{PDG16}: $R_{LU}^0 = 0.82 \pm 0.08$ ($\sim 2.0\sigma$)

$$B(D^0 \rightarrow \pi^- \mu^+ \nu) = (0.237 \pm 0.024)\%$$



改进测定 $B[D^0 \rightarrow \pi^- \mu^+ \nu] = (0.267 \pm 0.007 \pm 0.007)\%$

首次测定 $B[D^+ \rightarrow \pi^0 \mu^+ \nu] = (0.342 \pm 0.011 \pm 0.010)\%$

$$R_{LU}^0 = 0.918 \pm 0.036$$


$$R_{LU}^+ = 0.921 \pm 0.045$$

Amplitude analysis of $D^+ \rightarrow V e^+ \nu$


PRD94(2016)032001

PRD92(2015)071101(RC)



$D^+ \rightarrow K^- \pi^+ e^+ \nu$



Model independent form factors of $D^+ \rightarrow \bar{K}^{*0} e^+ \nu$



$D^+ \rightarrow \omega e^+ \nu$



$$r_V = V(0)/A_1(0) = 1.24 \pm 0.09 \pm 0.06$$


$$r_2 = A_2(0)/A_1(0) = 1.06 \pm 0.15 \pm 0.05$$

Observation of $D \rightarrow a_0(980)^-(0) e^+ \nu$

- Explore the nontrivial internal structure of light hadron mesons, traditional $q\bar{q}$ states, tetra quark system.
- With chiral unitarity approach in the coupled channels, BF is predicted to be order of $5(6) \times 10^{-5}$ for $D^{0(+)}$ decays
- Improve understanding of classification of light scalar mesons

$$R \equiv \frac{B(D^+ \rightarrow f_0 l^+ \nu) + B(D^+ \rightarrow \sigma l^+ \nu)}{B(D^+ \rightarrow a_0 l^+ \nu)}$$

$R=1(3)$ if traditional $q\bar{q}$ (tetra quark) system



$$\begin{aligned} & \odot B(D^0 \rightarrow a_0(980)^- e^+ \nu_e) \times B(a_0(980)^- \rightarrow \eta \pi^-) \\ & = (1.12^{+0.31}_{-0.28}(\text{stat}) \pm 0.10(\text{syst})) \times 10^{-4} \end{aligned}$$

5.9σ



$$\begin{aligned} & \odot B(D^+ \rightarrow a_0(980)^0 e^+ \nu_e) \times B(a_0(980)^0 \rightarrow \eta \pi^0) \\ & = (1.47^{+0.73}_{-0.59}(\text{stat}) \pm 0.14(\text{syst})) \times 10^{-4} \end{aligned}$$

3.0σ

$$\begin{aligned} & \odot B(D^+ \rightarrow a_0(980)^0 e^+ \nu_e) \times B(a_0(980)^0 \rightarrow \eta \pi^0) \\ & < 2.7 \times 10^{-4} \quad @ 90\% \text{ C.L.} \end{aligned}$$


Search for rare D decays

In SM, the BF's of charm rare decay are expected to be less than 10^{-6}



PRD 91(2015)112015

Consistent with Babar result



Blinding the selection, optimize cuts and study backgrounds based on MC and sideband data.


LNV decay

$\mathcal{B}(D^+ \rightarrow) \backslash [\times 10^{-6}]$	$K^+ e^+ e^-$	$K^- e^+ e^+$	$\pi^+ e^+ e^-$	$\pi^- e^+ e^+$
CLEO	3.0	3.6	3.9	1.1
Babar	1.0	0.9	1.1	1.9
PDG	1.0	0.9	1.1	1.1
This work	1.2	0.6	0.3	1.2


ST method

First searches for $D^+ \rightarrow \gamma e^+ \nu$ and $D^0 e^+ \nu$

Various theory models
predict BF's in 10^{-6} – 10^{-4}





Tree level amplitudes



$B[D^+ \rightarrow \gamma e^+ \nu]_{|E_\gamma > 10 \text{ MeV}} < 3.0 \times 10^{-4}$
@90% C.L.

Applying the SU(3) symmetry for the light quarks, this rare decay branching fraction can be predicted by theoretical calculation and its theoretical value is 2.78×10^{-13} [EPJC, 59:841-845(2009)].



$B[D^+ \rightarrow D^0 e^+ \nu] < 1 \times 10^{-4}$ @90% C.L.

D hadronic decays

- $D^0\bar{D}^0$ mixing parameters
- Strong phase difference
- SU(3) symmetry and break effect

$D^0\bar{D}^0$ mixing and CPV

- Open-flavor neutral meson transforms to its anti-meson and vice versa:

$$K^0 \Leftrightarrow \bar{K}^0, B_d^0 \Leftrightarrow \bar{B}_d^0, B_s^0 \Leftrightarrow \bar{B}_s^0, D^0 \Leftrightarrow \bar{D}^0$$

- Flavor eigenstate ($|D^0\rangle, |\bar{D}^0\rangle$) $\neq$ mass eigenstate $|D_{1,2}\rangle$ with $M_{1,2}$ and $\Gamma_{1,2}$)


$$|D_{1,2}\rangle \equiv p|D^0\rangle \pm q|\bar{D}^0\rangle \quad (\text{CPT: } p^2 + q^2 = 1)$$

- Mixing parameters definition:

$$\boxed{\mathbf{x} \equiv \frac{M_1 - M_2}{\Gamma}, \quad \mathbf{y} \equiv \frac{\Gamma_1 - \Gamma_2}{2\Gamma}, \quad \Gamma \equiv \frac{\Gamma_1 + \Gamma_2}{2}}$$

- under phase convention
 $CP|D^0\rangle = |\bar{D}^0\rangle, CP|\bar{D}^0\rangle = |D^0\rangle,$
- with CP conservation ($q = p = 1/\sqrt{2}$):
 $|D_{1,2}\rangle = |D_{+,-}\rangle$ (CP eigenstates)

- Unique: only the up-type meson for mixing
- Standard Model predicts: $\sim \mathcal{O}(1\%)$



(1) short distance ($< 0.1\%$ by CKM and GIM)



(2) long distance ($\sim 1\%$)

- Precise measurement of x, y : effectively limit New Physics(NP) modes;
- search for NP, eg: $|x| \gg |y|$

$D^0\bar{D}^0$ mixing and CPV

Decay Mode	Observables	Relationship
$D^0 \rightarrow K^+ K^- / \pi^+ \pi^-$	y_{CP} A_Γ	$2y_{CP} = (q/p + p/q) y \cos \phi -$ $(q/p - p/q) x \sin \phi$ $2A_\Gamma = (q/p - p/q) y \cos \phi -$ $(q/p + p/q) x \sin \phi$
$D^0 \rightarrow K_S^0 \pi^+ \pi^-$	x y $ q/p $ ϕ	
$D^0 \rightarrow K^+ \ell^- \nu$	R_M	$R_M = (x^2 + y^2)/2$
$D^0 \rightarrow K^+ \pi^- \pi^0$ (Dalitz plot analysis)	x'' y''	$x'' = x \cos \delta_{K\pi\pi} + y \sin \delta_{K\pi\pi}$ $y'' = y \cos \delta_{K\pi\pi} - x \sin \delta_{K\pi\pi}$
“Double-tagged” branching fractions measured in $\psi(3770) \rightarrow DD$ decays	R_M y R_D $\sqrt{R_D} \cos \delta$	$R_M = (x^2 + y^2)/2$
$D^0 \rightarrow K^+ \pi^-$	x'^2, y' x'^{2+}, x'^{2-} y'^+, y'^-	$x' = x \cos \delta + y \sin \delta$ $y' = y \cos \delta - x \sin \delta$ $A_M \equiv (q/p ^4 - 1)/(q/p ^4 + 1)$ $x'^{\pm} = [(1 \pm A_M)/(1 \mp A_M)]^{1/4} \times$ $(x' \cos \phi \pm y' \sin \phi)$ $y'^{\pm} = [(1 \pm A_M)/(1 \mp A_M)]^{1/4} \times$ $(y' \cos \phi \mp x' \sin \phi)$
$D^0 \rightarrow K^+ \pi^- / K^- \pi^+$ (time-integrated)	$\frac{\Gamma(D^0 \rightarrow K^+ \pi^-) + \Gamma(\bar{D}^0 \rightarrow K^- \pi^+)}{\Gamma(D^0 \rightarrow K^- \pi^+) + \Gamma(\bar{D}^0 \rightarrow K^+ \pi^-)}$ $\frac{\Gamma(D^0 \rightarrow K^+ \pi^-) - \Gamma(\bar{D}^0 \rightarrow K^- \pi^+)}{\Gamma(D^0 \rightarrow K^- \pi^+) + \Gamma(\bar{D}^0 \rightarrow K^+ \pi^-)}$	R_D A_D
$D^0 \rightarrow K^+ K^- / \pi^+ \pi^-$ (time-integrated)	$\frac{\Gamma(D^0 \rightarrow K^+ K^-) - \Gamma(\bar{D}^0 \rightarrow K^+ K^-)}{\Gamma(D^0 \rightarrow K^+ K^-) + \Gamma(\bar{D}^0 \rightarrow K^+ K^-)}$ $\frac{\Gamma(D^0 \rightarrow \pi^+ \pi^-) - \Gamma(\bar{D}^0 \rightarrow \pi^+ \pi^-)}{\Gamma(D^0 \rightarrow \pi^+ \pi^-) + \Gamma(\bar{D}^0 \rightarrow \pi^+ \pi^-)}$	$A_K + \frac{\langle t \rangle}{\tau_D} \mathcal{A}_{CP}^{\text{indirect}} \quad (\mathcal{A}_{CP}^{\text{indirect}} \approx -A_\Gamma)$ $A_\pi + \frac{\langle t \rangle}{\tau_D} \mathcal{A}_{CP}^{\text{indirect}} \quad (\mathcal{A}_{CP}^{\text{indirect}} \approx -A_\Gamma)$

混合参数:x,y

间接CPV参数:|q/p|,φ

直接CPV参数: A_D, A_K, A_π

R_D


强相差参数: $\delta_{K\pi}, \delta_{K\pi\pi 0}$

Evidence of $D^0\bar{D}^0$ mixing

■ Babar, 384 fb⁻¹@10.58 GeV

PRL98(2007)211802

$$\frac{T_{\text{WS}}(t)}{e^{-\Gamma t}} \propto R_D + \sqrt{R_D} y' \Gamma t + \frac{x'^2 + y'^2}{4} (\Gamma t)^2$$




$$\begin{aligned} R_D &= (0.303 \pm 0.016(\text{stat}) \pm 0.010(\text{syst}))\% \\ x'^2 &= (-0.22 \pm 0.30(\text{stat}) \pm 0.21(\text{syst})) \times 10^{-3} \\ y' &= (9.7 \pm 4.4(\text{stat}) \pm 3.1(\text{syst})) \times 10^{-3} \\ (x'^2, y') &\text{ with correlation } -0.95 \end{aligned}$$

■ BELLE, 540 fb⁻¹@10.58 GeV

PRL98(2007)211803

$$y_{CP} = \frac{\tau(K^- \pi^+)}{\tau(K^+ K^-)} - 1$$

$$A_\Gamma = \frac{\tau(\bar{D}^0 \rightarrow K^- K^+) - \tau(D^0 \rightarrow K^+ K^-)}{\tau(\bar{D}^0 \rightarrow K^- K^+) + \tau(D^0 \rightarrow K^+ K^-)}$$



$$\begin{aligned} A_\Gamma &= [0.01 \pm 0.30(\text{stat}) \pm 0.15(\text{syst})]\% \\ y_{CP} &= [1.31 \pm 0.32(\text{stat}) \pm 0.25(\text{syst})]\% \end{aligned}$$


Observation of $D^0\bar{D}^0$ mixing

■ LHCb, 3 fb⁻¹ p $\bar{p}$ at 7/8 TeV

PRL110(2013)101802 (1fb⁻¹)

PRL111(2013)251801

$$R(t) \approx R_D + \sqrt{R_D} y' \frac{t}{\tau} + \frac{x'^2 + y'^2}{4} \left(\frac{t}{\tau}\right)^2$$



$$x'^2 = (5.5 \pm 4.9) \times 10^{-5},$$

$$y' = (4.8 \pm 1.0) \times 10^{-3}$$


$$R_D = (3.568 \pm 0.066) \times 10^{-3}$$

■ CDFII, 9.6 fb⁻¹ p $\bar{p}$ at 1.96 TeV

PRL100(2008)121802(1.5 fb⁻¹)

PRL111(2013)231802

$$R(t/\tau) = R_D + \sqrt{R_D} y' (t/\tau) + \frac{x'^2 + y'^2}{4} (t/\tau)^2.$$



$$x'^2 = (0.08 \pm 0.18) \times 10^{-3}$$


$$y' = (4.3 \pm 4.3) \times 10^{-3},$$

$$R_D = (3.51 \pm 0.35) \times 10^{-3}$$

■ Belle, 976 fb⁻¹ at 10.58 GeV

PRL112(2014)111801

$$R(\tilde{t}/\tau) = \frac{\Gamma_{\text{WS}}(\tilde{t}/\tau)}{\Gamma_{\text{RS}}(\tilde{t}/\tau)} \approx R_D + \sqrt{R_D} y' \frac{\tilde{t}}{\tau} + \frac{x'^2 + y'^2}{4} \left(\frac{\tilde{t}}{\tau}\right)^2$$



$$x'^2 = (0.09 \pm 0.22) \times 10^{-3}$$

$$y' = (4.6 \pm 3.4) \times 10^{-3}$$

$$R_D = (3.53 \pm 0.13) \times 10^{-3}$$

$D^0\bar{D}^0$ mixing parameter y_{CP}

■ BESIII, 3 fb⁻¹ at 3.773 GeV

PLB744(2015)339

For D decay to CP eigenstates:

$$R_{CP\pm} \propto |A_{CP\pm}|^2 (1 \mp y_{CP})$$

$$y_{CP} = \frac{1}{2} [y \cos \phi (|\frac{q}{p}| + |\frac{p}{q}|) - x \sin \phi (|\frac{q}{p}| - |\frac{p}{q}|)]$$


For CP tagged semileptonic D decays:

$$R_{l,CP\pm} \propto |A_l|^2 |A_{CP\pm}|^2$$



$$y_{CP} \approx \frac{1}{4} \left(\frac{R_{l,CP+} R_{CP-}}{R_{l,CP-} R_{CP+}} - \frac{R_{l,CP-} R_{CP+}}{R_{l,CP+} R_{CP-}} \right)$$

Type	Modes
CP^+	$K^+ K^-, \pi^+ \pi^-, K_S \pi^0 \pi^0$
CP^-	$K_S^0 \pi^0, K_S^0 \omega, K_S^0 \eta$
$l^\pm$	$K e \nu, K \mu \nu$

$$y_{CP} = (-2.0 \pm 1.3 \pm 0.7) \%$$



Strong phase difference $\delta_{K\pi}$



$$\mathcal{A}_{CP \rightarrow K\pi} = \frac{\mathcal{B}_{D_2 \rightarrow K^- \pi^+} - \mathcal{B}_{D_1 \rightarrow K^- \pi^+}}{\mathcal{B}_{D_2 \rightarrow K^- \pi^+} + \mathcal{B}_{D_1 \rightarrow K^- \pi^+}}.$$

$$2r \cos \delta_{K\pi} + y = (1 + R_{WS}) \cdot \mathcal{A}_{CP \rightarrow K\pi},$$

$$|D_1\rangle \equiv \frac{|D^0\rangle + |\overline{D}^0\rangle}{\sqrt{2}} \quad |D_2\rangle \equiv \frac{|D^0\rangle - |\overline{D}^0\rangle}{\sqrt{2}}.$$

Type	Mode
Flavored	$K^- \pi^+, K^+ \pi^-$
CP+	$K^+ K^-, \pi^+ \pi^-, K_S^0 \pi^0 \pi^0, \pi^0 \pi^0, \rho^0 \pi^0$
CP-	$K_S^0 \pi^0, K_S^0 \eta, K_S^0 \omega$

$\delta_{K\pi}$ is related to mixing parameters x and y from x' and y'





目前最精确结果

$$A^{K\pi}_{CP} = (12.7 \pm 1.3 \pm 0.7) \times 10^{-2}$$

$$\cos \delta_{K\pi} = 1.02 \pm 0.11 \pm 0.06 \pm 0.01$$

PLB734(2014)227

$D^0\bar{D}^0$ mixing and CPV



$D^0\bar{D}^0$ mixing is observed, no direct CPV is found

Strong phase: bridge to constrain γ/ϕ_3

- Quantum correlated $D^0\bar{D}^0$ decays in $\psi(3770)$:
 - CP asymmetry in mixing and decays
 - Interference $\rightarrow$ strong phase parameters $\rightarrow$ Constrain γ/ϕ_3 , which is important for CKM UT

Direct measurement


$$\alpha/\phi_2 = (85.4^{+4.0}_{-3.9})^\circ$$

$$\beta/\phi_1 = (21.38^{+0.79}_{-0.77})^\circ$$

$$\gamma/\phi_3 = (68^{+8.0}_{-8.5})^\circ$$



γ is the worst measured angle,
mostly due to systematic error

Significant deviation from UT will
imply NP beyond SM



Strong phase difference $\delta_{K_S\pi^+\pi^-}$

Babar 2008 Optimal Bins



Mirrored binning over $x=y$ makes it so $c_i=c_i$ and $s_i=s_i$

□ Model prediction


● BESIII

▼ CLEO-c


BESIII preliminary

CLEO, PRD82, 112006


c_i



s_i




$c_i s_i$



- T_i : Bin yield measured in flavor decays
- r_B : color suppression factor ~ 0.1
- δ_B : strong phase of B decay
- c_i, s_i : weighted average of $\cos(\Delta\delta_D)$ and $\sin(\Delta\delta_D)$ respectively
where $\Delta\delta_D$ is the difference between phase of D^0 and D^0

Measured at B-Factories

Through $D^0 \rightarrow K_S \pi^+ \pi^-$ analysis



Constrain γ/ϕ_3 measurement


taken from Liming Zhang's talk at FPCPV2016




γ combination at LHCb

Determine γ from CPV measurements

LHCb-PAPER-2016-032



GLW: $D \rightarrow K^+ K^- \pi^+ \pi^-$
 $K_S^0 \pi^0$
 ADS: $D \rightarrow \pi^+ K^-$
 quasi-ADS: $D \rightarrow \pi^+ K^- \pi^+ \pi^-$
 $\pi^+ K^- \pi^0$
 GGSZ: $D \rightarrow K_S^0 \pi^+ \pi^-$
 $K_S^0 K^+ K^-$
 quasi-GLW: $D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$
 $K^+ K^- \pi^0$
 $\pi^+ \pi^- \pi^0$
 GLS: $D \rightarrow K_S^0 K^+ \pi^+$
 $K_S^0 \pi^+ K^+$



$$\gamma = (72.2_{-7.3}^{+6.8})^\circ \text{ syst. included}$$

$$\text{BaBar: } \gamma = (70 \pm 18)^\circ$$

$$\text{Belle: } \gamma = (73_{-15}^{+13})^\circ$$

Prospects

Sample	$\sigma_{\text{stat}}(\gamma)^\circ$
Run 1	8
Run 2	4
Upgrade	~ 1
Future upgrade	< 0.5

- Current one syst. $\sim 2^\circ$ from CLEO strong phase measurements
- 15-20 fb^{-1} $\psi(3770)$ data from BESIII are desired to avoid syst. limitation for upgrade scenario

More 15 fb^{-1} $\psi(3770)$ data@BESIII will avoid syst. limitation for γ/ϕ_3 measurement

SU(3) symmetries and breaking effect

- Ratio of branching fractions of D to KK and pi pi

$$R = \frac{Br(D^0 \rightarrow K^+ K^-)}{Br(D^0 \rightarrow \pi^+ \pi^-)} \approx 2.8$$

- R=1 in the SU(3) flavour symmetry limit
- Branching fraction of $\mathcal{B}(D^0 \rightarrow K^0 \bar{K}^0) = (0.320 \pm 0.038) \times 10^{-3}$ vanishes in the SU(3) limit
- DDbar mixing parameters
$$x, y \sim \sin^2 \theta_C \times [SU(3) \text{ breaking}]^2$$
- Non-zero mixing parameters indicate large SU(3) breaking effect


Theoretical calculations

Modes	Amplitudes	Br(FSI)	Br(diagrammatic)	Br(pole)	Br(FAT)	Br(FAT[mix])	Br(exp)
$D^0 \rightarrow \pi^+ \rho^-$	$T_V, (E_P)$	6.5	3.92 ± 0.46	3.5 ± 0.6	4.74	4.66	4.96 ± 0.24
$D^0 \rightarrow \pi^0 \rho^0$	$C_P, C_V, (E_P, E_V)$	1.7	2.96 ± 0.98	1.4 ± 0.6	3.55	3.83	3.72 ± 0.22
$D^0 \rightarrow \pi^0 \omega$	$C_P, C_V, (E_P, E_V)$	0.08	0.10 ± 0.18	0.08 ± 0.02	0.85	0.18	< 0.26
$D^0 \rightarrow \pi^0 \phi$	C_P	1.1	1.22 ± 0.08	1.0 ± 0.3	1.11	1.11	1.31 ± 0.10
$D^0 \rightarrow \pi^- \rho^+$	$T_P, (E_V)$	8.2	8.34 ± 1.69	10.2 ± 1.5	10.2	10.0	9.8 ± 0.4
$D^0 \rightarrow K^+ K^{*-}$	$T_V, (E_P)$	2.8	1.99 ± 0.24	1.6 ± 0.3	1.72	1.73	1.56 ± 0.12
$D^0 \rightarrow K^0 \bar{K}^{*0}$	E_P, E_V	0.99	0.29 ± 0.22	0.16 ± 0.05	1.1	1.1	< 1
$D^0 \rightarrow \bar{K}^0 K^{*0}$	E_P, E_V	0.99	0.29 ± 0.22	0.16 ± 0.05	1.1	1.1	< 0.56
$D^0 \rightarrow K^- K^{*+}$	$T_P, (E_V)$	4.5	4.25 ± 0.86	4.7 ± 0.8	4.37	4.37	4.38 ± 0.21
$D^0 \rightarrow \eta \rho^0$	$C_P, C_V, (E_P, E_V)$	0.24	1.11 ± 0.86	0.05 ± 0.01	0.54	0.45	
$D^0 \rightarrow \eta \omega$	$C_P, C_V, (E_P, E_V)$	1.9	3.08 ± 1.42	1.2 ± 0.3	2.4	2.0	
$D^0 \rightarrow \eta \phi$	$C_P, (E_P, E_V)$	0.57	0.31 ± 0.10	0.23 ± 0.06	0.19	0.18	0.14 ± 0.05
$D^0 \rightarrow \eta' \rho^0$	$C_P, C_V, (E_P, E_V)$	0.10	0.14 ± 0.02	0.08 ± 0.02	0.21	0.27	
$D^0 \rightarrow \eta' \omega$	$C_P, C_V, (E_P, E_V)$	0.001	0.07 ± 0.02	0.0001 ± 0.0001	0.04	0.02	
$D^+ \rightarrow \pi^+ \rho^0$	$T_V, C_P, (A_P, A_V)$	1.7		0.8 ± 0.7	0.42	0.58	0.81 ± 0.15
$D^+ \rightarrow \pi^+ \omega$	$T_V, C_P, (A_P, A_V)$	0.35		0.3 ± 0.3	0.95	0.80	< 0.34
$D^+ \rightarrow \pi^+ \phi$	C_P	5.9	6.21 ± 0.43	5.1 ± 1.4	5.65	5.65	$5.42^{+0.22}_{-0.24}$
$D^+ \rightarrow \pi^0 \rho^+$	$T_P, C_V, (A_P, A_V)$	3.7		3.5 ± 1.6	2.7	2.5	
$D^+ \rightarrow K^+ \bar{K}^{*0}$	$T_V, (A_V)$	2.5		4.1 ± 1.0	3.61	3.60	$3.675^{+0.14}_{-0.21}$
$D^+ \rightarrow \bar{K}^0 K^{*+}$	$T_P, (A_P)$	1.70		12.4 ± 2.4	11	11	32 ± 14
$D^+ \rightarrow \eta \rho^+$	$T_P, C_V, (A_P, A_V)$	0.002		0.4 ± 0.4	0.7	2.2	< 15
$D^+ \rightarrow \eta' \rho^+$	$T_P, C_V, (A_P, A_V)$	1.3		0.8 ± 0.1	0.7	0.8	
$D_s^+ \rightarrow \pi^+ K^{*0}$	$T_V, (A_V)$	3.3		1.5 ± 0.7	2.52	2.35	2.25 ± 0.39
$D_s^+ \rightarrow \pi^0 K^{*+}$	$C_V, (A_V)$	0.29		0.1 ± 0.1	0.8	1.0	
$D_s^+ \rightarrow K^+ \rho^0$	$C_P, (A_P)$	2.4		1.0 ± 0.6	1.9	2.5	2.7 ± 0.5
$D_s^+ \rightarrow K^+ \omega$	$C_P, (A_P)$	0.72		1.8 ± 0.7	0.6	0.07	< 2.4
$D_s^+ \rightarrow K^+ \phi$	$T_V, C_P, (A_V)$	0.15		0.3 ± 0.3	0.166	0.166	0.184 ± 0.045
$D_s^+ \rightarrow K^0 \rho^+$	$T_P, (A_P)$	19.5		7.5 ± 2.1	9.1	9.6	
$D_s^+ \rightarrow \eta K^{*+}$	$T_P, C_V, (A_P, A_V)$	0.24		1.0 ± 0.4	0.2	0.2	
$D_s^+ \rightarrow \eta' K^{*+}$	$T_P, C_V, (A_P, A_V)$	0.24		0.6 ± 0.2	0.2	0.2	


Observation/evidence of $D \rightarrow \omega \pi$

Double tag method

PRL116(2016)082001



Decay mode	This work	Previous measurements
$D^+ \rightarrow \omega \pi^+$	$(2.74 \pm 0.58 \pm 0.17) \times 10^{-4}$	$< 3.4 \times 10^{-4}$ at 90% C.L.
$D^0 \rightarrow \omega \pi^0$	$(1.05 \pm 0.41 \pm 0.09) \times 10^{-4}$	$< 2.6 \times 10^{-4}$ at 90% C.L.
$D^+ \rightarrow \eta \pi^+$	$(3.13 \pm 0.22 \pm 0.19) \times 10^{-3}$	$(3.53 \pm 0.21) \times 10^{-3}$
$D^0 \rightarrow \eta \pi^0$	$(0.67 \pm 0.10 \pm 0.05) \times 10^{-3}$	$(0.68 \pm 0.07) \times 10^{-3}$




Studies of singly cabibbo-suppressed decays is limited by data set and background


Benefit the understanding of SU(3) symmetry breaking and CP violation, improve theory calculation

Two-body decays of D_s^+


首次确认重子衰变 $D_s^+ \rightarrow p\bar{n}$



(a) Short-distance effects




(b) Long-distance effects



$$\mathcal{B}_{D_s^+ \rightarrow p\bar{n}} = (1.22 \pm 0.10) \times 10^{-3}$$


首次确认纯湮灭衰变 $D_s^+ \rightarrow \omega\pi^+$
并首次测定 $D_s^+ \rightarrow \omega K^+$



Under the model-independent approach:

$$A(D_s^+ \rightarrow \omega\pi^+) = \frac{1}{\sqrt{2}} (A_P + A_V)$$


($A_{P,V}$: P, V contain $\bar{q}'$ of $q\bar{q}'$ configuration)



Signal mode	Branching fraction (10^{-3})
$D_s^+ \rightarrow \omega\pi^+$	$1.85 \pm 0.30(stat.) \pm 0.19(sys.)$
$D_s^+ \rightarrow \omega K^+$	$1.13 \pm 0.24(stat.) \pm 0.14(sys.)$

Amplitude analyses of D decays


PRD89(2014)052001



Mode	Partial Branching Fraction (%)
$D^+ \rightarrow K_S^0 \pi^+ \pi^0$ Non Resonant	$0.32 \pm 0.05 \pm 0.25^{+0.28}_{-0.25}$
$D^+ \rightarrow \rho^+ K_S^0, \rho^+ \rightarrow \pi^+ \pi^0$	$5.83 \pm 0.16 \pm 0.30^{+0.45}_{-0.15}$
$D^+ \rightarrow \rho(1450)^+ K_S^0, \rho(1450)^+ \rightarrow \pi^+ \pi^0$	$0.15 \pm 0.02 \pm 0.09^{+0.07}_{-0.11}$
$D^+ \rightarrow \bar{K}^*(892)^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$	$0.250 \pm 0.012 \pm 0.015^{+0.025}_{-0.024}$
$D^+ \rightarrow \bar{K}_0^*(1430)^0 \pi^+, \bar{K}_0^*(1430)^0 \rightarrow K_S^0 \pi^0$	$0.26 \pm 0.04 \pm 0.05 \pm 0.06$
$D^+ \rightarrow \bar{K}^*(1680)^0 \pi^+, \bar{K}^*(1680)^0 \rightarrow K_S^0 \pi^0$	$0.09 \pm 0.01 \pm 0.05^{+0.04}_{-0.08}$
$D^+ \rightarrow \bar{\kappa}^0 \pi^+, \bar{\kappa}^0 \rightarrow K_S^0 \pi^0$	$0.54 \pm 0.09 \pm 0.28^{+0.36}_{-0.19}$
$NR + \bar{\kappa}^0 \pi^+$	$1.30 \pm 0.12 \pm 0.12^{+0.12}_{-0.30}$
$K_S^0 \pi^0$ S-wave	$1.21 \pm 0.10 \pm 0.16^{+0.19}_{-0.27}$

Help to determine the absolute BF, strong phase, benefit γ/ϕ_3


Previous analyses only from MarkIII and E691



Component	Branching fraction (%)	PDG value (%)
$D^0 \rightarrow \bar{K}^* \rho^0$	$0.99 \pm 0.04 \pm 0.04 \pm 0.03$	1.05 ± 0.23
$D^0 \rightarrow K^- a_1^+(1260)(\rho^0 \pi^+)$	$4.41 \pm 0.22 \pm 0.30 \pm 0.13$	3.6 ± 0.6
$D^0 \rightarrow K_1^-(1270)(\bar{K}^* \pi^-) \pi^+$	$0.07 \pm 0.01 \pm 0.02 \pm 0.00$	0.29 ± 0.03
$D^0 \rightarrow K_1^-(1270)(K^- \rho^0) \pi^+$	$0.27 \pm 0.02 \pm 0.04 \pm 0.01$	
$D^0 \rightarrow K^- \pi^+ \rho^0$	$0.68 \pm 0.09 \pm 0.20 \pm 0.02$	0.51 ± 0.23
$D^0 \rightarrow \bar{K}^{*0} \pi^+ \pi^-$	$0.57 \pm 0.03 \pm 0.04 \pm 0.02$	0.99 ± 0.23
$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$	$1.77 \pm 0.05 \pm 0.04 \pm 0.05$	1.88 ± 0.26

Λ_c^+ decays

Measurements before 2014



➤ Λ_c^+ was observed in 1979

➤ All decays of Λ_c^+ were measured with high energy data and relative to $pK^-\pi^+$, which suffers an error of 25%. No absolute measurement using threshold Λ_c^+ data

➤ Only about 60% decays are known



Λ_c^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Hadronic modes with a p: $S = -1$ final states			
$p\bar{K}^0$	(2.3 ± 0.6) %		873
$pK^-\pi^+$	[a] (5.0 ± 1.3) %		823
$p\bar{K}^*(892)^0$	[b] (1.6 ± 0.5) %		685
$\Delta(1232)^{++}K^-$	(8.6 ± 3.0) × 10 ⁻³		710
$\Lambda(1520)\pi^+$	[b] (1.8 ± 0.6) %		627
$pK^-\pi^+$ nonresonant	(2.8 ± 0.8) %		823
$p\bar{K}^0\pi^0$	(3.3 ± 1.0) %		823
$p\bar{K}^0\eta$	(1.2 ± 0.4) %		568

Systematic studies of Λ_c^+ , search for new decays, absolute BF measurements are important to explore Λ_c^+ decay mechanisms⁴³

Improved BFs of $\Lambda_c^+ \rightarrow$ hadronic decays


BELLE, PRL113(2014)042002

BESIII, PRL116(2016)052001




$$B[\Lambda_c^+ \rightarrow pK^-\pi^+] = (6.84 \pm 0.24^{+0.21}_{-0.27})\%$$

ST: ~ 15000




DT: ~ 1000



Mode	This work (%)	PDG (%)
pK_S^0	$1.52 \pm 0.08 \pm 0.03$	1.15 ± 0.30
$pK^-\pi^+$	$5.84 \pm 0.27 \pm 0.23$	5.0 ± 1.3
$pK_S^0\pi^0$	$1.87 \pm 0.13 \pm 0.05$	1.65 ± 0.50
$pK_S^0\pi^+\pi^-$	$1.53 \pm 0.11 \pm 0.09$	1.30 ± 0.25
$pK^-\pi^+\pi^0$	$4.53 \pm 0.23 \pm 0.30$	3.4 ± 1.0
$\Lambda\pi^+$	$1.24 \pm 0.07 \pm 0.03$	1.07 ± 0.28
$\Lambda\pi^+\pi^0$	$7.01 \pm 0.37 \pm 0.19$	3.6 ± 1.3
$\Lambda\pi^+\pi^-\pi^+$	$3.81 \pm 0.24 \pm 0.18$	2.6 ± 0.7
$\Sigma^0\pi^+$	$1.27 \pm 0.08 \pm 0.03$	1.05 ± 0.28
$\Sigma^+\pi^0$	$1.18 \pm 0.10 \pm 0.03$	1.00 ± 0.34
$\Sigma^+\pi^+\pi^-$	$4.25 \pm 0.24 \pm 0.20$	3.6 ± 1.0
$\Sigma^+\omega$	$1.56 \pm 0.20 \pm 0.07$	2.7 ± 1.0


Much better
precision

First absolute BFs of $\Lambda_c^+ \rightarrow \Lambda l^+ \nu$



$$B[\Lambda_c^+ \rightarrow \Lambda e^+ \nu] = (3.63 \pm 0.38 \pm 0.20)\%$$

3 fb⁻¹ data help to explore FF studies




$$B[\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu] = (3.49 \pm 0.46 \pm 0.26)\%$$

$$\Gamma[\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu] / \Gamma[\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e] = 0.96 \pm 0.16 \pm 0.04$$

Calibrate theoretical calculations:
(1.4-9.2)%



Theoretical Models	predicated branching fraction for $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$
MBM [1]	1.9%
NRQM [1]	2.6%
SU(4)-symmetry limit [2]	9.2%
RSQM [3]	4.4%
QCM [4]	5.62%
SQM [5]	1.96%
NRQM2 [6]	2.15%
NRQM3 [7]	1.42%
QCD SR1 [8]	$(3.0 \pm 0.9)\%$
QCD SR2 [9]	$(2.6 \pm 0.4)\%$
QCD SR3 [9]	$(5.8 \pm 1.5)\%$
STSR [10]	2.22% for $\Lambda_c^+ \rightarrow \Lambda l^+ \nu_l$
STNR [10]	1.58% for $\Lambda_c^+ \rightarrow \Lambda l^+ \nu_l$
HOSR [10]	4.72% for $\Lambda_c^+ \rightarrow \Lambda l^+ \nu_l$
HONR [10]	4.2% for $\Lambda_c^+ \rightarrow \Lambda l^+ \nu_l$
LCSR _s [11]	$(3.0 \pm 0.3)\%$ for $\Lambda_c^+ \rightarrow \Lambda l^+ \nu_l$ (CZ-type)
PDG 2014 [14]	$(2.1 \pm 0.6)\%$
BESIII	$(3.63 \pm 0.38 \pm 0.20)\%$

促进LQCD对形状因子的计算



Measurements of SCS decays $\Lambda_c^+ \rightarrow pK^+K^-/\pi^+\pi^-$

BESIII, PRL117(2016)232002




These help to distinguish predictions from different theoretical models and understand contributions from factorizable effects

Decay modes	$\mathcal{B}_{\text{mode}}/\mathcal{B}_{\text{ref.}}$	$\mathcal{B}_{\text{mode}}$	$\mathcal{B}(\text{PDG})$
$\Lambda_c^+ \rightarrow p\pi^+\pi^-$	$(6.70 \pm 0.48 \pm 0.25) \times 10^{-2}$	$(3.91 \pm 0.28 \pm 0.15 \pm 0.24) \times 10^{-3}$	$(3.5 \pm 2.0) \times 10^{-3}$
$\Lambda_c^+ \rightarrow p\phi$	$(1.81 \pm 0.33 \pm 0.13) \times 10^{-2}$	$(1.06 \pm 0.19 \pm 0.08 \pm 0.06) \times 10^{-3}$	$(8.2 \pm 2.7) \times 10^{-4}$
$\Lambda_c^+ \rightarrow pK^+K^-$ (non- ϕ)	$(9.36 \pm 2.22 \pm 0.71) \times 10^{-3}$	$(5.47 \pm 1.30 \pm 0.41 \pm 0.33) \times 10^{-4}$	$(3.5 \pm 1.7) \times 10^{-4}$

Observation of $\Lambda_c^+ \rightarrow nK_S\pi^+$

BESIII, PRL118(2017)112001



Help to understand SU(3) and isospin symmetry and determine strong phase

Cai-Dian Lv et al, PRD93(2016)056008

$$\cos \delta$$

$$= \frac{\mathcal{B}(n\bar{K}^0\pi^+) - \mathcal{B}(pK^-\pi^+)}{2\sqrt{\mathcal{B}(p\bar{K}^0\pi^0)(\mathcal{B}(pK^-\pi^+) + \mathcal{B}(n\bar{K}^0\pi^+) - \mathcal{B}(p\bar{K}^0\pi^0))}}$$

$$R_p = \frac{\mathcal{B}(\Lambda_c \rightarrow p\bar{K}^0\pi^0)}{\mathcal{B}(\Lambda_c \rightarrow pK^-\pi^+)}, \quad R_n = \frac{\mathcal{B}(\Lambda_c \rightarrow n\bar{K}^0\pi^+)}{\mathcal{B}(\Lambda_c \rightarrow pK^-\pi^+)}$$

$$\mathcal{B}[\Lambda_c^+ \rightarrow nK_S\pi^+] = (1.82 \pm 0.23 \pm 0.11)\%$$

$$\Gamma[\Lambda_c^+ \rightarrow n\bar{K}^0\pi^+] / \Gamma[\Lambda_c^+ \rightarrow pK^-\pi^+] = 0.62 \pm 0.09$$

$$\Gamma[\Lambda_c^+ \rightarrow n\bar{K}^0\pi^+] / \Gamma[\Lambda_c^+ \rightarrow p\bar{K}^0\pi^+] = 0.97 \pm 0.16$$

First measurement of BF of Λ_c^+ decay containing neutron

$$\cos \delta = -0.24 \pm 0.08$$



$$|I^{(1)}|/|I^{(0)}| = 1.14 \pm 0.11$$

involving a neutron. Under the isospin symmetry, its amplitude is related to those of the most favored proton modes $\Lambda_c^+ \rightarrow pK^-\pi^+$ and $\Lambda_c^+ \rightarrow p\bar{K}^0\pi^0$ as $\mathcal{A}(n\bar{K}^0\pi^+) + \mathcal{A}(pK^-\pi^+) + \sqrt{2}\mathcal{A}(p\bar{K}^0\pi^0) = 0$. Hence, precise measure-

[2,3]. In the three-body Λ_c^+ decay to $N\bar{K}\pi$, the total decay amplitudes can be decomposed into two isospin amplitudes of the $N\bar{K}$ system as isosinglet ($I^{(0)}$) and isospin-one ($I^{(1)}$). In the factorization limit, the color-allowed tree diagram, in which the π^+ is emitted and the $N\bar{K}$ is an isosinglet, dominates $I^{(0)}$, and $I^{(1)}$ is expected to be small compared to $I^{(0)}$ as it can only proceed through the color-suppressed tree diagrams. Though the factorization scheme is spoiled in

Observation of $\Lambda_c^+ \rightarrow \Sigma^- \pi^+ \pi^+ \pi^0$

BESIII, PLB772(2017)388



Preliminary results :



$$B[\Lambda_c^+ \rightarrow \Sigma^- \pi^+ \pi^+] = (1.81 \pm 0.17 \pm 0.09)\%$$

$$B[\Lambda_c^+ \rightarrow \Sigma^- \pi^+ \pi^+ \pi^0] = (2.11 \pm 0.33 \pm 0.14)\% \quad \text{[First observation]}$$

The previous one is consistent with and more precise than the PDG value of $[\Lambda_c^+ \rightarrow \Sigma^- \pi^+ \pi^+] = (2.3 \pm 0.4)\%$.

Evidence of $\Lambda_c^+ \rightarrow p\eta$ and search of $\Lambda_c^+ \rightarrow p\pi^0$

BESIII, PRD95(2017)111102(RC)



$$B[\Lambda_c^+ \rightarrow p\pi^0] < 2.7 \times 10^{-4} \quad 90\% \text{CL}$$

	$\Lambda_c^+ \rightarrow p\eta$	$\Lambda_c^+ \rightarrow p\pi^0$	$\frac{B_{\Lambda_c^+ \rightarrow p\pi^0}}{B_{\Lambda_c^+ \rightarrow p\eta}}$
BESIII	1.24 ± 0.29	< 0.27	< 0.24
Sharma <i>et al.</i> [3]	$0.2^a(1.7^b)$	0.2	$1.0^a(0.1^b)$
Uppal <i>et al.</i> [4]	0.3	0.1–0.2	0.3–0.7
S. L. Chen <i>et al.</i> [12]	...	0.11–0.36 ^c	...
Cai-Dian Lü <i>et al.</i> [13]	...	0.45	...

^aAssumed to have a positive sign for the p-wave amplitude of $\Lambda_c^+ \rightarrow \Xi^0 K^+$.

^bAssumed to have a negative sign for the p-wave amplitude of $\Lambda_c^+ \rightarrow \Xi^0 K^+$.

^cCalculated relying on different values of parameters b and α .


$$B[\Lambda_c^+ \rightarrow p\eta] = (1.24 \pm 0.28 \pm 0.10) \times 10^{-3}$$

Larger threshold Λ_c^+ data at BESIII

国际粲重子 Λ_c^+ 衰变实验研究的里程碑：**BESIII**开辟使用近阈数据绝对测量 Λ_c^+ 衰变的新领域

	Golden hadronic mode	$\delta B/B$	Golden SL mode	$\delta B/B$
D^0	$B(K\pi)=(3.88\pm 0.05)\%$	1.3%	$B(K^0\pi)=(3.55\pm 0.05)\%$	1.4%
D^+	$B(K\pi\pi)=(9.13\pm 0.19)\%$	2.1%	$B(K^0\pi)=(8.83\pm 0.22)\%$	2.5%
D_s	$B(KK\pi)=(5.39\pm 0.21)\%$	3.9%	$B(\phi\pi)=(2.49\pm 0.14)\%$	5.6%
Λ_c	$B(pK\pi)=(5.0\pm 1.3)\%$ (PDG2014)	26%	$B(\Lambda\pi)=(2.1\pm 0.6)\%$ (PDG2014)	29%
	$= (6.8\pm 0.36)\%$ (BELLE)	5.3%	$= (3.63\pm 0.43)\%$ (BESIII)	12%
	$= (5.84\pm 0.35)\%$ (BESIII)	6.0%	$= (3.18\pm 0.32)\%$ (HFAG)	10%
	$= (6.46\pm 0.24)\%$ (HFAG)	3.7%		

更高能量**4.62-4.63 GeV**，更大近阈 Λ_c^+ 样本？改进已有衰变精度，半轻形状因子，寻找**40%**未知衰变...



Senior Research Positions

Latest Issue | **Archive** | Jobs | Links | Buyer's guide | White papers | Events | Contact us

Search Go

REGISTER NOW

Register as a member of cerncourier.com and get full access to all features of the site. Registration is free.

CERN COURIER

Mar 18, 2016

BESIII makes first direct measurement of the Λ_c at threshold

The charmed baryon, Λ_c , was first observed at Fermilab in 1976. Now, 40 years later, the Beijing Spectrometer (BESIII) experiment at the Beijing Electron-Positron Collider II (BEPCII) has measured the absolute branching fraction of $\Lambda_c^+ \rightarrow pK^+\pi^+$ at threshold for the first time.

Beam-constrained mass distribution

DIGITAL EDITION

CERN Courier is now available as a regular digital edition. Click here to read the digital edition.

KEY SUPPLIERS

JANIS
Cryogenic Systems

Linde

LATEST CERN COURIER ARTICLES

- ▶ Sneeze dynamics
- ▶ The longest proof
- ▶ Electron-hole collider
- ▶ Imaging with muons
- ▶ Towards a nuclear clock

37天数据已发表7篇物理文章，其中4篇PRL

Summary

- $D^{0(+)}$ 研究取得一些重要物理成果

- D^+ 衰变常数 f_{D^+}

- 形状因子 $f^{D \rightarrow K(\pi)}_+(q^2)$

- CKM矩阵元 $|V_{cs(d)}|$

- $D^0\bar{D}^0$ 混合参数 $y_{CP}, \delta_{K\pi}$

- $D^0 \rightarrow K_S \pi^+ \pi^-$ 强相差初步结果

→精密检验格点QCD计算
和CKM矩阵幺正性、探讨
 $D^0\bar{D}^0$ 混合、约束 γ/ϕ_3 测量

- Λ_c^+ 衰变的系统研究结束了其发现近40年来无近阈数据绝对测量的历史

- 2016年, 在4.178 GeV采集了 $3.2 \text{ fb}^{-1} D_s^+$ 数据。已取得 D_s^+ 衰变常数 $f_{D_s^+}$ 、CKM矩阵元 $|V_{cs}|$ 等初步结果

- 更多物理结果将在未来1-2年完成

谢谢!

Observation of DCS decay of $\Lambda_c^+ \rightarrow pK^+\pi^-$

BELLE, PRL117(2016)011801





FIG. 1. Typical external (internal) W -emission diagrams for (a) [(c)] $\Lambda_c^+ \rightarrow pK^+\pi^-$ and (b) [(d)] $\Lambda_c^+ \rightarrow pK^-\pi^+$, and (e) a typical W -exchange diagram of $\Lambda_c^+ \rightarrow pK^-\pi^+$.



$$B[\Lambda_c^+ \rightarrow pK^+\pi^-]/B[\Lambda_c^+ \rightarrow pK^-\pi^+] \sim \tan^4\theta$$

$$\sin\theta \sim 0.225 \pm 0.001$$

$$B[\Lambda_c^+ \rightarrow pK^+\pi^-]/B[\Lambda_c^+ \rightarrow pK^-\pi^+] \sim (2.35 \pm 0.27 \pm 0.21)\%$$

Search for penta-quark in $\Lambda_c^+ \rightarrow p K^+ K^- \pi^0$

BELLE, PRD96(2016)051102(RC)







FIG. 1. Feynman diagram for the decay (a) $\Lambda_c^+ \rightarrow \phi p \pi^0$ and (b) $\Lambda_c^+ \rightarrow P_s^+ \pi^0$.

Inspired by the observation of two hidden-charm pentaquark states $P_c^+(4380)$ and $P_c^+(4450)$ in $J/\psi p$ invariant mass spectrum at LCHb

Search for hidden-strangeness pentaquark states in ϕp invariant mass spectrum



$$\mathcal{B}(\Lambda_c^+ \rightarrow \phi p \pi^0) < 15.3 \times 10^{-5}, \text{ @90\%C.L.}$$

$$\mathcal{B}(\Lambda_c^+ \rightarrow K^+ K^- p \pi^0)_{\text{NR}} < 6.3 \times 10^{-5}.$$

$$\mathcal{B}(\Lambda_c^+ \rightarrow P_s^+ \pi^0) \times \mathcal{B}(P_s^+ \rightarrow \phi p) < 8.3 \times 10^{-5}$$



Measurements of $\Lambda_c^+ \rightarrow pK^+K^-/p\pi^+\pi^-/pK^-\pi^+$

LHCb, 1711.01157[hep-ex]

1 fb⁻¹ data @ 7 TeV

$\Lambda_b^0 \rightarrow \Lambda_c^+(phh')\mu^-\bar{\nu}_\mu$ selection

Prompt $\Lambda_c^+ \rightarrow phh'$ selection



$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow p\pi^-\pi^+)}{\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)} = (7.44 \pm 0.08 \pm 0.18) \%,$$

$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow pK^-K^+)}{\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)} = (1.70 \pm 0.03 \pm 0.03) \%,$$

$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow p\pi^-K^+)}{\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)} = (0.165 \pm 0.015 \pm 0.005) \%,$$

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\pi^-\pi^+) = (4.72 \pm 0.05 \pm 0.11 \pm 0.25) \times 10^{-3},$$

$$\mathcal{B}(\Lambda_c^+ \rightarrow pK^-K^+) = (1.08 \pm 0.02 \pm 0.02 \pm 0.06) \times 10^{-3},$$

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\pi^-K^+) = (1.04 \pm 0.09 \pm 0.03 \pm 0.05) \times 10^{-4},$$

Analysis of $D^+ \rightarrow K_L e^+ \nu$


- Regardless of long flight distance, K_L interact with EMC and deposit part of energy, thus giving position information
- After reconstructing all other particles, K_L can be inferred with position information and constraint $U_{\text{miss}} \rightarrow 0$

$$\overline{B}(D^+ \rightarrow K_L e^+ \nu) = (4.482 \pm 0.027 \pm 0.103)\%$$

$$A_{CP} \equiv \frac{B(D^+ \rightarrow K_L^0 e^+ \nu_e) - B(D^- \rightarrow K_L^0 e^- \bar{\nu}_e)}{B(D^+ \rightarrow K_L^0 e^+ \nu_e) + B(D^- \rightarrow K_L^0 e^- \bar{\nu}_e)}$$

$$A_{CP}^{D^+ \rightarrow K_L e^+ \nu} = (-0.59 \pm 0.60 \pm 1.50)\%$$

Simultaneous fit to event density $I(q^2)$ with 2-par. series Form Factor



$D^+ \rightarrow K_L e^+ \nu$ is measured for the first time

PRD92(2015)112008


With 6 dominant D^- single tag

$$f_{K^+}^K(0)|V_{cs}| = 0.728 \pm 0.006 \pm 0.011$$


$$r_1 = a_1/a_0 = -1.91 \pm 0.33 \pm 0.24$$

Absolute BF for $D^+ \rightarrow \bar{K}^0 e^+ \nu$ via $\bar{K}^0 \rightarrow \pi^0 \pi^0$

With 6 dominant D^- single tag



Taking τ_{D^+} , τ_{D^0} , $B[D^0 \rightarrow K^- e^+ \nu]$ and $B[D^+ \rightarrow \bar{K}^0 e^+ \nu]$ from the PDG as input




$$\frac{\Gamma[D^0 \rightarrow K^- e^+ \nu]}{\bar{\Gamma}[D^+ \rightarrow \bar{K}^0 e^+ \nu]} = 0.969 \pm 0.025$$


Agrees with isospin conservation within 1.2σ

Improved BF for $D^+ \rightarrow \bar{K}^0 \mu^+ \nu$

With 6 dominant D^- single tag



EPJC76(2016)369



Taking $B[D^0 \rightarrow K^- \mu^+ \nu]$
and $B[D^+ \rightarrow \bar{K}^0 e^+ \nu]$
from the PDG as input

$$\frac{\Gamma[D^0 \rightarrow K^- \mu^+ \nu]}{\bar{\Gamma}[D^+ \rightarrow \bar{K}^0 \mu^+ \nu]} = 0.963 \pm 0.044$$

$$\frac{\Gamma[D^+ \rightarrow \bar{K}^0 \mu^+ \nu]}{\Gamma[D^+ \rightarrow \bar{K}^0 e^+ \nu]} = 0.988 \pm 0.033$$


Support isospin conservation in
these two decays within errors

Consistent with theory
prediction 0.97 within error

Measurements of BFs of $D_s^+ \rightarrow \eta^{(\prime)} e^+ \nu$


- Benefit the understanding of the source of difference of inclusive decay rates of $D^{0(+)}$ and D_s^+
- Complementary information to understand η - η' mixing

482 pb⁻¹ data@4.009 GeV, PRD94(2016)112003




	BESIII	CLEOII 95	CLEOc09	CLEOc15	PDG [4]
$B(D_s^+ \rightarrow \eta e^+ \nu_e)[\%]$	$2.30 \pm 0.31 \pm 0.08$	—	$2.48 \pm 0.29 \pm 0.13$	$2.28 \pm 0.14 \pm 0.20$	2.67 ± 0.29
$B(D_s^+ \rightarrow \eta' e^+ \nu_e)[\%]$	$0.93 \pm 0.30 \pm 0.05$	—	$0.91 \pm 0.33 \pm 0.05$	$0.68 \pm 0.15 \pm 0.06$	0.99 ± 0.23
$\frac{B(D_s^+ \rightarrow \eta' e^+ \nu_e)}{B(D_s^+ \rightarrow \eta e^+ \nu_e)}$	$0.40 \pm 0.14 \pm 0.02$	$0.35 \pm 0.09 \pm 0.07$	—	—	—

Studies of $D_s^+ \rightarrow K^{(*)0} e^+ \nu$ at 4.178 GeV





$$B[D_s^+ \rightarrow K^0 e^+ \nu_e] = (3.25 \pm 0.38_{\text{stat}} \pm 0.14_{\text{syst}}) \times 10^{-3}$$

$$(3.9 \pm 0.9) \times 10^{-3} \text{ [PDG17]}$$



$$B[D_s^+ \rightarrow K^{*0} e^+ \nu_e] = (2.38 \pm 0.26_{\text{stat}} \pm 0.12_{\text{syst}}) \times 10^{-3}$$


$$(1.8 \pm 0.4) \times 10^{-3} \text{ [PDG17]}$$



Model	Parameter	Value	$f_+(0)$
Simple pole	$f_+(0) V_{cd} $	$0.175 \pm 0.010 \pm 0.001$	$0.778 \pm 0.044 \pm 0.004$
Modified pole model	$f_+(0) V_{cd} $	$0.163 \pm 0.017 \pm 0.003$	$0.725 \pm 0.076 \pm 0.013$
	α	$0.45 \pm 0.44 \pm 0.02$	
Series two parameters	$f_+(0) V_{cd} $	$0.162 \pm 0.019 \pm 0.003$	$0.720 \pm 0.084 \pm 0.013$
	r_1	$-2.94 \pm 2.32 \pm 0.14$	

Taking $|V_{CKM}^{\text{fitter}}|_{cd}|$ as input



Four dimensional un-binned likelihood fit is performed. K^* parameters are fixed



$$r_V = 1.67 \pm 0.34 \pm 0.16$$

$$r_2 = 0.77 \pm 0.28 \pm 0.07$$

Inclusive decay $\Lambda_c^+ \rightarrow \Lambda X$



Help to explore the source of missing decays and search for new decay. Better input for charm baryon and B physics

$$N_{sig} = N_S - (N_A + N_B)/2 - r \cdot N_D + r \cdot (N_C + N_E)/2$$

$$\mathcal{B}(\Lambda_c^+ \rightarrow \Lambda + X) = (36.98 \pm 2.18)\% \quad \text{stat. only}$$


Agrees with PDG2015 value (35±11)%,

$$\mathcal{A}_{CP} = \frac{\mathcal{B}(\Lambda_c^+ \rightarrow \Lambda + X) - \mathcal{B}(\bar{\Lambda}_c^- \rightarrow \bar{\Lambda} + X)}{\mathcal{B}(\Lambda_c^+ \rightarrow \Lambda + X) + \mathcal{B}(\bar{\Lambda}_c^- \rightarrow \bar{\Lambda} + X)}$$

Decay mode	Branching fraction(%)	$\mathcal{A}_{CP}$
$\Lambda_c^+ \rightarrow \Lambda + X$	38.02 ± 3.24	0.02 ± 0.06
$\bar{\Lambda}_c^- \rightarrow \bar{\Lambda} + X$	36.70 ± 3.04	

Absolute BFs and y_{CP} of $D^0 \rightarrow K_{S/L} \pi^0 (\pi^0)$

- Two dimensional fits to $M_{BC}(\text{tag})$ versus $M_{BC}(\text{signal})$
- Projections of DT events on the $M_{BC}(\text{sig})$ vs. $K\pi$ (for example)



Branching fractions and asymmetries


Statistical only

$$R(D \rightarrow K_{S,L} \pi^0) = \frac{Br(D \rightarrow K_S \pi^0) - Br(D \rightarrow K_L \pi^0)}{Br(D \rightarrow K_S \pi^0) + Br(D \rightarrow K_L \pi^0)}$$

Table 10: Decay rates and the asymmetries of $D \rightarrow K_{S,L}^0 \pi^0$ and $D \rightarrow K_{S,L}^0 \pi^0 \pi^0$.


$D \rightarrow K_{S,L}^0 \pi^0$			
	$Br_{K_S \pi^0}(\%)$	$Br_{K_L \pi^0}(\%)$	$R(D \rightarrow K_{S,L} \pi^0)$
$K\pi$	1.208 ± 0.041	1.061 ± 0.038	0.0646 ± 0.0245
$K3\pi$	1.212 ± 0.037	0.985 ± 0.036	0.1035 ± 0.0237
$K\pi\pi^0$	1.251 ± 0.028	0.953 ± 0.029	0.1351 ± 0.0186
All	1.230 ± 0.020	0.991 ± 0.019	0.1077 ± 0.0125

$D \rightarrow K_{S,L}^0 \pi^0 \pi^0$			
	$Br_{K_S 2\pi^0}(\%)$	$Br_{K_L 2\pi^0}(\%)$	$R(D \rightarrow K_{S,L} 2\pi^0)$
$K\pi$	1.024 ± 0.049	1.299 ± 0.080	-0.1183 ± 0.0385
$K3\pi$	0.887 ± 0.043	1.097 ± 0.073	-0.1060 ± 0.0409
$K\pi\pi^0$	1.010 ± 0.036	1.158 ± 0.060	-0.0681 ± 0.0313
All	0.975 ± 0.024	1.175 ± 0.040	-0.0929 ± 0.0209



- $y_{CP} ((K_S \pi^0, K_L \pi^0) \text{ vs. } K\pi) = (0.98 \pm 2.43)\%$

Absolute BFs and A_{CP} of $D^+ \rightarrow K_{S/L} K^+ (\pi^0)$



The first and second uncertainties are statistical and systematic


$$A_{CP} = \frac{\mathcal{B}(D^+) - \mathcal{B}(D^-)}{\mathcal{B}(D^+) + \mathcal{B}(D^-)}$$

Mode	$\mathcal{B}(D^+) (\times 10^{-3})$	$\mathcal{B}(D^-) (\times 10^{-3})$	$\bar{\mathcal{B}} (\times 10^{-3})$	$A_{CP} (\%)$
$K_S^0 K^\pm$	$3.01 \pm 0.12 \pm 0.10$	$3.10 \pm 0.12 \pm 0.10$	$3.06 \pm 0.09 \pm 0.10$	$-1.5 \pm 2.8 \pm 1.6$
$K_S^0 K^\pm \pi^0$	$5.23 \pm 0.28 \pm 0.24$	$5.09 \pm 0.29 \pm 0.22$	$5.16 \pm 0.21 \pm 0.23$	$1.4 \pm 4.0 \pm 2.4$
$K_L^0 K^\pm$	$3.13 \pm 0.14 \pm 0.13$	$3.32 \pm 0.15 \pm 0.13$	$3.23 \pm 0.11 \pm 0.13$	$-3.0 \pm 3.2 \pm 1.2$
$K_L^0 K^\pm \pi^0$	$5.17 \pm 0.30 \pm 0.21$	$5.26 \pm 0.30 \pm 0.20$	$5.22 \pm 0.22 \pm 0.21$	$-0.9 \pm 4.1 \pm 1.6$

BFs of $D^+ \rightarrow 2K_S K(\pi)^+$ and $D^0 \rightarrow 2(3)K_S$

Comprehensive or improved measurements of 3-body decays benefit the understanding of the interplay between weak and strong interactions in multibody decays, where theory is poor than 2-body decays

BF of $D^0 \rightarrow K_S K_S$ will be helpful to explore the SU(3) symmetry breaking in D decays




Comparisons of the branching fractions (in 10^{-4}) measured in this work with the PDG values

Decay modes	This work	PDG
$D^+ \rightarrow K_S^0 K_S^0 K^+$	$25.4 \pm 0.5 \pm 1.2$	45 ± 20
$D^+ \rightarrow K_S^0 K_S^0 \pi^+$	$27.0 \pm 0.5 \pm 1.2$	–
$D^0 \rightarrow K_S^0 K_S^0$	$1.67 \pm 0.11 \pm 0.11$	1.7 ± 0.4
$D^0 \rightarrow K_S^0 K_S^0 K_S^0$	$7.21 \pm 0.33 \pm 0.44$	9.1 ± 1.3

PLB765(2017)231

$D_s^+ \rightarrow \eta' X$ and $\eta' \rho^+$



~ 15.6 K ST

$$B_{\text{CLEO}}[D_s^+ \rightarrow \eta' \rho^+] = (12.5 \pm 2.2)\%$$

PRD58(1998)052002


是理论预期 $(3.0 \pm 0.5)\%$ 的4倍

F.S.Yu PRD84(2011)074019


$$B_{\text{PDG14}}^{\text{SUM}}[D_s^+ \rightarrow \eta' X] = (18.6 \pm 2.3)\%$$

$$B_{\text{MSR}}[D_s^+ \rightarrow \eta' X] = (11.7 \pm 1.8)\%$$

PRD79(2009)112008



$$N[D_s^+ \rightarrow \eta' X] = 68 \pm 14$$



$$N[D_s^+ \rightarrow \eta' \rho^+] = 210 \pm 50$$



$$B[D_s^+ \rightarrow \eta' X] = (8.8 \pm 1.8 \pm 0.5)\%$$

$$B[D_s^+ \rightarrow \eta' \rho^+] = (5.8 \pm 1.4 \pm 0.4)\%$$

与CLEOPRD88(2013)032009一致

新实验结果解决了实验和理论不一致的矛盾

BFs of $D^{0(+)} \rightarrow PP$



BESIII preliminary

- ◆ The study of the hadronic decays of charmed D mesons is of great significance in the study of the strong and weak interactions in D decays.
- ◆ The analysis on $D \rightarrow PP$ modes will provide materials for the study of SU(3) breaking effect¹. And the observation of CP violation in D decay is commonly believed to be indications of new physics.
- ◆ $D^0 \rightarrow K^- \pi^+$ is an important normalization mode.
- ◆ Most of the D decays have been studied by CLEO in 2010², other measurements come from Belle³, BaBar⁴ and CDF⁵, etc.
- ◆ Some of the branching fractions (BFs) are not well established. With the 2.93 fb^{-1} data taken at 3.773 GeV within BESIII, the results will help to improve these measurements.

Mode	$N_{\text{signal}}^{\text{net}}$	ϵ (%)	$\mathcal{B} \pm (\text{stat}) \pm (\text{sys})$	$\mathcal{B}_{\text{PDG}}$
$\pi^+ \pi^-$	21105 ± 249	66.03 ± 0.25	$(1.505 \pm 0.018 \pm 0.031) \times 10^{-3}$	$(1.421 \pm 0.025) \times 10^{-3}$
$K^+ K^-$	5543 ± 273	62.82 ± 0.32	$(4.229 \pm 0.020 \pm 0.087) \times 10^{-3}$	$(4.01 \pm 0.07) \times 10^{-3}$
$K^- \pi^+$	537745 ± 767	64.98 ± 0.09	$(3.896 \pm 0.006 \pm 0.073) \%$	$(3.93 \pm 0.04) \%$
$K_S^0 \pi^0$	66539 ± 302	38.06 ± 0.17	$(1.236 \pm 0.006 \pm 0.032) \%$	$(1.20 \pm 0.04) \%$
$K_S^0 \eta$	9532 ± 126	31.96 ± 0.14	$(5.149 \pm 0.068 \pm 0.134) \times 10^{-3}$	$(4.85 \pm 0.30) \times 10^{-3}$
$K_S^0 \eta'$	3007 ± 61	12.66 ± 0.08	$(9.562 \pm 0.197 \pm 0.379) \times 10^{-3}$	$(9.5 \pm 0.5) \times 10^{-3}$
$\pi^0 \pi^+$	10108 ± 267	48.98 ± 0.34	$(1.259 \pm 0.033 \pm 0.025) \times 10^{-3}$	$(1.24 \pm 0.06) \times 10^{-3}$
$\pi^0 K^+$	1834 ± 168	51.52 ± 0.42	$(2.171 \pm 0.198 \pm 0.060) \times 10^{-4}$	$(1.89 \pm 0.25) \times 10^{-4}$
$\eta \pi^+$	11636 ± 215	46.96 ± 0.25	$(3.790 \pm 0.070 \pm 0.075) \times 10^{-3}$	$(3.66 \pm 0.22) \times 10^{-3}$
ηK^+	439 ± 72	48.21 ± 0.31	$(1.393 \pm 0.228 \pm 0.124) \times 10^{-4}$	$(1.12 \pm 0.18) \times 10^{-4}$
$\eta' \pi^+$	3088 ± 83	21.49 ± 0.18	$(5.122 \pm 0.140 \pm 0.210) \times 10^{-3}$	$(4.84 \pm 0.31) \times 10^{-3}$
$\eta' K^+$	87 ± 25	22.39 ± 0.22	$(1.377 \pm 0.428 \pm 0.202) \times 10^{-4}$	$(1.83 \pm 0.23) \times 10^{-4}$
$K_S^0 \pi^+$	93884 ± 352	51.38 ± 0.18	$(1.591 \pm 0.006 \pm 0.033) \times 10^{-2}$	$(1.53 \pm 0.06) \times 10^{-2}$
$K_S^0 K^+$	17704 ± 151	48.45 ± 0.14	$(3.183 \pm 0.028 \pm 0.065) \times 10^{-3}$	$(2.95 \pm 0.15) \times 10^{-3}$


$$\mathcal{B} = \frac{N_{\text{signal}}^{\text{net}}}{2 \cdot N_{D^0 \bar{D}^0} (D^+ D^-) \cdot \epsilon}, N_{D^0 \bar{D}^0} = (10,621 \pm 29_{\text{stat}}) \times 10^3, N_{D^+ D^-} = (8,296 \pm 31_{\text{stat}}) \times 10^3$$

quoted from Derrick's talk given at APS2014



The $\mathcal{B}(D^0 \rightarrow K^- \pi^+)$ has been corrected by the PDG value of $\mathcal{B}(D^0 \rightarrow K^+ \pi^-)$.

For $D^0 \rightarrow K_S^0 \eta$, $D^+ \rightarrow \pi^0 \pi^+$, $D^+ \rightarrow \eta \pi^+$, $D^+ \rightarrow \eta' \pi^+$, $D^+ \rightarrow K_S^0 \pi^+$ and $D^+ \rightarrow K_S^0 K^+$, it shows better precision than the present values.

Status of rare D^0 decays



Status of rare $D_{(s)}^+$ decays



So far, no rare D decay is found