





Holographic Dark Matter Fluid in Late Time Universe

by Yun-Long Zhang (张云龙)

Asia Pacific Center for Theoretical Physics(APCTP)
@Pohang, Korea (韩国浦项)




APCTP
Asia Pacific Center for Theoretical Physics

Emergent Dark Matter in Late Universe on Holographic Screen

Rong-Gen Cai (ITP) & Sichun Sun(NTU) & Yun-Long Zhang(APCTP)

[arXiv: [1712.09326](https://arxiv.org/abs/1712.09326)]

New Era in Black Holes & Gravitational Waves (2016-)



Med ena hälften till
With one half to:



och med den andra hälften gemensamt till
and with the other half jointly to:

Rainer Weiss
LIGO/VIRGO Collaboration

Barry C. Barish
LIGO/VIRGO Collaboration

Kip S. Thorne
LIGO/VIRGO Collaboration

"för avgörande bidrag till LIGO-detektorn och observationen av gravitationsvågor"
"for decisive contributions to the LIGO detector and the observation of gravitational waves"



Gamma Ray Burst: NS-NS
Fermi-GBM 2017



High Energy Neutrino: Blazar!
IceCube 2018

Research Background of Y.-L. Zhang

Black Holes and Quantum Matters
Diffusions & Emergent Gravity & SYK Model

2016–2018 Postdoc at APCTP (亚太理论物理中心)
Young Scientist Training(YST) Program @Pohang



Holography and Effective Theory
Phase Transition & μ_{2e} & $0\nu\beta\beta$

2014–2016 Postdoc at NTU (台湾大学)
with Prof. Jiunn-Wei Chen @Taipei






Gravity and Hydrodynamics
Einstein Eqs & Navier-Stokes Eqs

2009–2014 Ph.D at ITP/CAS (中科院理论物理所)
with Prof. Rong-Gen Cai @Beijing
[2013–2014 visiting at U. of Southampton(UK)]



2005–2009 B.S. at Lanzhou U.(兰州大学) with Prof. Yu-Xiao Liu


New Physics Between Neutron Star and Black Holes?



Universal Holographic Properties of Horizon

$$\frac{\eta}{s} \simeq \frac{1}{4\pi} \frac{\hbar}{k_B}$$


$$\tau_c^{-1} \simeq \frac{k^2}{4\pi T_c}$$



$$\Omega_D^2 \simeq \frac{1}{2} \Omega_\Lambda (\Omega_D - \Omega_B)$$

$$\frac{H^2}{H_0^2} \simeq \frac{\Omega_B}{a^3} + \sqrt{\Omega_\Lambda \left(\frac{H^2}{H_0^2} + \frac{\Omega_I}{a^4} \right)}$$

Quantum Critical Liquid
Graphene & Semi-Metal & QGP



Rindler Fluid [[1705.05078](#)]
by Lee, Park, Y.-L. Zhang

Cosmological Fluid
Dark Matter & Energy




Cosmic Fluid [[1712.09326](#)]
by Cai, Sun, Y.-L. Zhang

Holographic Screens in Flat Spacetime


Holographic Rindler Fluid

Accelerating Screen in Flat Spacetime
From Conformal Fluid to Rindler Fluid



Holographic de-Sitter Fluid

de-Sitter & FRW Screen
Relation to DGP brane world Models




Successful Holographic Models — ICTP Dirac Medal 2018




S. Sachdev (Harvard)

Holographic
Strange Metal




Dom T. Son (Chicago)

Holographic
Quark-Gluon Plasma



Xiao-Gang Wen (MIT)


Emergent Phenomena
Entanglement & Network



Holographic Duality in Classical Mechanics


I. Electromagnetism

Surface Charge
Electric Field




II. Newton Gravity

Mass Sphere Shell
Gravitational Field



III. Einstein Gravity

Surface Stress Tensor
Extrinsic Curvature





Thermodynamics (1970s): Hawking Radiation

Bekenstein & Hawking, ...

Hawking Temperature $T_H = \frac{\hbar c^3}{8\pi GM k_B} = \frac{\kappa}{2\pi}$

Bekenstein-Hawking Entropy $S_{\text{BH}} = \frac{kA}{4\ell_P^2}$



0th Law: constant surface gravity



1st Law: $dE = \frac{\kappa}{8\pi} dA + \Omega dJ + \Phi dQ,$

2nd Law: non-decreasing of entropy

3rd Law: extremal black hole is not possible

Membrane paradigm(1980s): Effective Fluid

T. Doumer & K. Thorne, ...




Effective Description

$$\mathcal{T}_{ab} = -2(K_{ab} - K\gamma_{ab})$$

Membrane on Stretched horizon

Viscosity & Conductivity



Echoes from the Abyss [1612.00266 PRD'17]

Traversable Wormholes or Black Holes?







Figure Credit: ScienceNews



Is the Gravitational-Wave Ringdown a Probe of the Event Horizon?

V. Cardoso, E. Franzin, P. Pani [PRL. 116, 171101 (2016)]

AdS/CFT Duality (2000s):

Maldacena & Gubser & Witten, et al

$$Z_{CFT} = \langle e^{S_{CFT}} \rangle \stackrel{AdS/CFT}{\simeq} e^{S_{AdS}}$$

AdS/CMT Correspondence


Black Hole in a natural Cavity

Shear Viscosity $\frac{\eta}{s} \simeq \frac{1}{4\pi} \frac{\hbar}{k_B T}$


Conductivity $\sigma \simeq \pi T$

Holographic Superconductor

Holographic Non-Fermi Liquid



Motivations for the Accelerating Screen




Extremal Charged BH




Finite Temperature

Rindler Space/
Special CMT





Membrane
Black Holes




Wilsonian Approach to Fluid/Gravity Duality [Bredberg, Keeler, Lysov, Strominger, '11]


Why in Rindler Frame



EVENT HORIZONS: From Black Holes to Acceleration




A stationary observer outside the black hole would see the thermal Hawking radiation. by Pisen Chen




An accelerating observer in vacuum would see a similar Hawking-like radiation called Unruh radiation.

$$\mathcal{T}_{ab} = -2(K_{ab} - K\gamma_{ab})$$


Credit: Physics Napkins



From AdS/CFT to Holographic Rindler Fluid — with an Accelerating Screen





Holographic Screen
The Time-like boundary



What is Rindler Fluid?

Fluid dual to Rindler spacetime




Navier–Stokes Equations:	Bredberg, Keeler, Lysov, Strominger ['10,'11]
Fluid/Gravity Expansion:	Compere, McFadden, Skenderis, Taylor ['11,'12]
Entropy Current and Constraint:	Chirco, Eling, Liberati, Meyer, Oz ['12,'13]
Comparison with AdS/Fluid:	Matsuo, Natsuume, Ohta, Okamura ['12,13]
Rindler Fluid and Recurrence Relation	Cai, Li, Yang, Zhang ['13,'14]
Rindler Fluid with Momentum Relaxation	Khimphun, Lee, Park, Zhang ['17]

Rindler Hydrodynamics

Rindler Metric $ds^2 = -r d\tau^2 + \frac{1}{r} dr^2 + dx_i dx^i$

Induced Metric $ds^2 = -r_c d\tau^2 + dx_i dx^i$

Dual Tensor $\mathcal{T}_{ab} = -2(K_{ab} - K\gamma_{ab})$



Constraint equations


$$2G_{\mu b} n^\mu|_{r_c} = 2\partial^a (K_{ab} - \gamma_{ab} K) = 0 \Rightarrow \partial^a T_{ab} = 0$$

$$2G_{\mu\nu} n^\mu n^\nu|_{r_c} = (K^2 - K_{ab} K^{ab}) = 0 \Rightarrow T^2 - p T_{ab} T^{ab} = 0$$

Bredberg, Keeler, Lysov, Strominger (JHEP 07 (2012) 146)


What is the Most Perfect Fluid in the World?

Quark Gluon Plasma
in RHIC [08'] & LHC [16']



$$\frac{\eta}{s} \simeq \frac{1}{4\pi} \frac{\hbar}{k_B}$$


Quantum Critical Liquid
Graphene [09'] & Semi-Metal [16']




Black Holes
[KSS,05']
Rindler Fluid
[BKLS,11']

$$\frac{H^2}{H_0^2} \simeq \frac{\Omega_B}{a^3} + \sqrt{\Omega_\Lambda \left(\frac{H^2}{H_0^2} + \frac{\Omega_I}{a^4} \right)}$$

$$\Omega_D^2 \simeq \frac{1}{2} \Omega_\Lambda (\Omega_D - \Omega_B)$$



Dark Fluid in the Universe?
Cosmological Fluid [CSZ,17']
[1712.09326, Cai, Sun, Zhang]




From Observation to Newton's Gravity






Tycho's Data
(1590s)



Kapler's Law
(1618)






Newton's Gravity
(1687)



NATURE and Nature's Laws lay hid in Night: God said, "Let Newton be!" and all was light.

From Einstein's Gravity to Dark Universe



Newton's Gravity

Special Relativity (1905)




Einstein's Gravity (1915)



Dark Matters



Gravitational Waves (2016)



Black Holes





Dark Energy



It did not last: the Devil howling: "Ho! Let Einstein be!" restored the status quo. — J. C. Squire

From Observation to Milgrom's MOND (Modified Newton Dynamics)




Galaxy Rotation Curve
(1970s)

Tully-Fisher Relation
(1977)

Milgrom's MOND
(1983)

$$v_f^4 \simeq a_0 G_N M_B$$

$$F_N = ma \mu\left(\frac{a}{a_0}\right)$$




Dark Matter

$$a_0 \simeq \sqrt{\Lambda}$$


20 years after MOND

$$\nabla \cdot \left[\mu \left(\frac{|\nabla\Phi|}{a_0} \right) \nabla\Phi \right] = 4\pi G\rho$$

Famaey & McGaugh,
Living Rev.Rel. 15 (2012) 10



Bullet Clusters



Acoustic Power Spectrum of CMB

Table 2: Observational tests of MOND.

Observational Test	Successful	Promising	Unclear	Problematic
Rotating Systems				
solar system			X	
galaxy rotation curve shapes	X			
surface brightness $\propto \Sigma \propto a^2$	X			
galaxy rotation curve fits	X			
fitted M./L	X			
Tully-Fisher Relation				
baryon based	X			
slope	X			
normalization	X			
no size nor Σ dependence	X			
no intrinsic scatter	X			
Galaxy Disk Stability				
maximum surface density	X			
spiral structure in LSBGs	X			
thin & bulgeless disks		X		
Interacting Galaxies				
tidal tail morphology		X		
dynamical friction			X	
tidal dwarfs	X			
Spheroidal Systems				
star clusters			X	
ultrafaint dwarfs			X	
dwarf Spheroidals	X			
ellipticals	X			
Faber Jackson relation	X			
Clusters of Galaxies				
dynamical mass				X
mass-temperature slope	X			
velocity (bulk & collisional)		X		
Gravitational Lensing				
strong lensing	X			
weak lensing (clusters & LSS)			X	
Cosmology				
expansion history			X	
geometry			X	
big bang nucleosynthesis	X			
Structure Formation				
galaxy power spectrum			X	
empty voids		X		
early structure		X		
Background Radiation				
first:second acoustic peak	X			
second:third acoustic peak				X
detailed fit				X
early re-ionization	X			

Constraints on MOND from Gravitational waves

Chesler & Loeb, arXiv:1704.05116 [PRL, '17]

1) The Speed of gravitational waves

Constraint of energy loss rate from ultra-high energy cosmic rays

2) Linear equations of motion in the weak-field limit

The observed gravitational waveforms from LIGO, which are consistent with Einstein's gravity

$$S = \frac{1}{16\pi G} \int d^4x \sqrt{g} [R + \mathcal{M}^2 \mathcal{F}(\frac{\kappa}{\mathcal{M}^2}) + \lambda(A^2 + 1)] + S_{\text{mat}}$$


Einstein-Aether theory (2004, Bekenstein)

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = \mathcal{T}_{\mu\nu} + 8\pi G T_{\mu\nu}^{\text{mat}},$$


$$\nabla_{\alpha}[\mathcal{F}' J^{\alpha}_{\beta}] - \mathcal{F}' y_{\beta} = 2\lambda A_{\beta},$$

$$\mathcal{T}_{\alpha\beta} = \frac{1}{2}\nabla_{\sigma}\{\mathcal{F}'[J_{(\alpha}{}^{\sigma} A_{\beta)} - J^{\sigma}{}_{(\alpha} A_{\beta)} - J_{(\alpha\beta)} A^{\sigma}]\}$$

$$- \mathcal{F}' Y_{\alpha\beta} + \frac{1}{2}g_{\alpha\beta}\mathcal{M}^2\mathcal{F} + \lambda A_{\alpha}A_{\beta},$$



From Verlinde's Gravity to Dark Universe




projecting data on 2 dimensional surface

Area Law & Holography



Volume Law & Entanglement




Entropy Gravity (2010)



Verlinde's Gravity (2016)

$$\int_0^r \frac{GM_D^2(r')}{r'^2} dr' = \frac{M_B(r)a_0 r}{6}$$


Tully-Fisher relation



Cluster of galaxies



Parameters in LCDM



$$g_D(r) = \sqrt{a_M g_B(r)}$$

$$a_M = \frac{a_0}{6}$$

$$\bar{\rho}_D^2(r) = \left(4 - \bar{\beta}_B(r)\right) \frac{a_0}{8\pi G} \frac{\bar{\rho}_B(r)}{r}$$

$$a_0 = cH_0$$

$$\Omega_D^2 = \frac{4}{3} \Omega_B$$

No Covariant Equations of Motion!

Compare with Verlinde's Emergent Universe

Gravitational quantity		Elastic quantity		Correspondence
Newtonian potential	Φ	displacement field	u_i	$u_i = \Phi n_i / a_0$
gravitational acceleration	g_i	strain tensor	ε_{ij}	$\varepsilon_{ij} n_j = -g_i / a_0$
surface mass density	Σ_i	stress tensor	σ_{ij}	$\sigma_{ij} n_j = \Sigma_i a_0$
mass density	ρ	body force	b_i	$b_i = -\rho a_0 n_i$
point mass	m	point force	f_i	$f_i = -m a_0 n_i$

Holographic Universe vs. Emergent Universe?

$$\frac{\mathcal{T}^2}{d-1} - \mathcal{T}_{\mu\nu} \mathcal{T}^{\mu\nu} = -\frac{\rho_\Lambda c^2}{d-1} (T + \mathcal{T}).$$

Constrain Equations



$$\Delta_V \equiv \Omega_D^2 - \frac{4}{3} \Omega_B \simeq 0.36\%,$$

$$\Delta_{CSZ} \equiv \Omega_D^2 - \frac{1}{2} \Omega_\Lambda (\Omega_D - \Omega_B) \simeq -0.34\%.$$

R.G. Cai, S. Sun, Y.L. Zhang, [1712.09326](#)

LCDM Universe?


$$H(a)^2 = H_0^2 [\Omega_\Lambda + (\Omega_D + \Omega_B) a^{-3} + \Omega_R a^{-4}]$$



Summary of the Membrane Fluid


Quark Gluon Plasma

RHIC ['08] & LHC ['16]



Quantum Critical Liquid

Graphene ['09] & Semi-Metal ['16]




Black Holes
Membrane Fluid [KSS,05']
Rindler Fluid [BKLS,11']

$$\frac{\eta}{s} \simeq \frac{1}{4\pi} \frac{\hbar}{k_B}$$

$$\tau_c^{-1} \simeq \frac{k^2}{4\pi T_c}$$

$$\frac{H^2}{H_0^2} \simeq \frac{\Omega_B}{a^3} + \sqrt{\Omega_\Lambda \left(\frac{H^2}{H_0^2} + \frac{\Omega_I}{a^4} \right)}$$


$$\Omega_D^2 \simeq \frac{1}{2} \Omega_\Lambda (\Omega_D - \Omega_B)$$



Cosmological Fluid [CSZ,'17]

Dark Matter & Dark Energy

$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} - \frac{1}{L} (\mathcal{K}_{\mu\nu} - \mathcal{K} g_{\mu\nu}) = \kappa_4 T_{\mu\nu}$$



Holographic dS Universe? — de-Sitter Screen

1) Holographic Stress Tensor — Dark Sectors


$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = \kappa_4 T_{\mu\nu} + \kappa_4 \langle \mathcal{T} \rangle_{\mu\nu}, \quad \langle \mathcal{T} \rangle_{\mu\nu} \equiv \frac{1}{\kappa_4 L} (\mathcal{K}_{\mu\nu} - \mathcal{K}g_{\mu\nu})$$

Modified Einstein equations

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} - \frac{1}{L} (\mathcal{K}_{\mu\nu} - \mathcal{K}g_{\mu\nu}) = \kappa_4 T_{\mu\nu}$$

Hamiltonian constraints

$$\mathcal{K}^2 - \mathcal{K}_{\mu\nu}\mathcal{K}^{\mu\nu} = R + 2G_{MN}^{(d+1)} \mathcal{N}^M \mathcal{N}^N,$$




2) Embedding in higher dimensions — Brane Worlds

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = \mathcal{T}_{\mu\nu}^M + T_{\mu\nu}^B,$$

$$\mathcal{T}_{\mu\nu}^M \equiv (\mathcal{K}g_{\mu\sigma} - \mathcal{K}_{\mu\sigma})\mathcal{K}^\sigma{}_\nu + \mathcal{M}_{\mu\nu} - \frac{1}{2}(\mathcal{K}^2 - \mathcal{K}_{\rho\sigma}\mathcal{K}^{\rho\sigma})g_{\mu\nu},$$

$$\mathcal{M}_{\mu\nu} \equiv g_\mu{}^M g_\nu{}^N R_{MN}^{(d+1)} - g_\mu{}^M \mathcal{N}^P g_\nu{}^N \mathcal{N}^Q R_{MPNQ}^{(d+1)}.$$

Ref: 1106.2476 [Living Rev. '10]



FRW Screen in a Flat Bulk

$$S_5 = \frac{1}{2\kappa_5} \int_{\mathcal{M}} d^5x \sqrt{-\tilde{g}} \mathcal{R} + \frac{1}{\kappa_5} \int_{\partial\mathcal{M}} d^4x \sqrt{-g} \mathcal{K},$$


$$S_4 = \frac{1}{2\kappa_4} \int_{\partial\mathcal{M}} d^4x \sqrt{-g} R + \int_{\partial\mathcal{M}} d^4x \sqrt{-g} \mathcal{L}_M.$$

FRW Screen

$$ds_4^2 = -c^2 dt^2 + a(t)^2 [dr^2 + r^2 d\Omega_2]$$


Friedmann eq.

$$\frac{H(t)^2}{H_0^2} \simeq \frac{\Omega_B}{a(t)^3} + \Omega_\Lambda^{1/2} \left[\frac{H(t)^2}{H_0^2} + \frac{\Omega_I}{a(t)^4} \right]^{1/2}$$



Ref: 1712.09326 [Cai, Sun, Zhang]


DGP BraneWorld



$$\frac{H(t)^2}{H_0^2} = \frac{\Omega_M}{a(t)^3} + \Omega_\ell^{1/2} \frac{H(t)}{H_0},$$



$$\dot{\rho}_i(t) = -3H(t) [\rho_i(t) + p_i(t)/c^2],$$

Self accelerating Branch



Ref: 1106.2476 [Living Rev. '10]

Late-Time Evolution of Holographic FRW Model



State equation of Holographic dark matter Fluid

$$\Delta_B = \frac{\Omega_M}{a(t)^3} = \frac{H(t)^2}{H_0^2} - \Delta_{\mathcal{H}}, \quad \Delta_{\mathcal{H}} \equiv \frac{\rho_{\mathcal{H}}}{\rho_c} = \Omega_{\Lambda}^{1/2} \left[\frac{H(t)^2}{H_0^2} + \frac{\Omega_I}{a(t)^4} \right]^{1/2},$$

$$\Delta_{\Lambda} = \Omega_{\Lambda}, \quad \Delta_D = \Delta_{\mathcal{H}} - \Omega_{\Lambda}, \quad w_D = -1 - \frac{1}{3H(t)} \frac{\dot{\Delta}_{\mathcal{H}}}{\Delta_{\mathcal{H}}}.$$

Embedding Function

$$ds_5^2 = \tilde{g}_{AB} dx^A dx^B = dy^2 - \mathbf{n}(y, t)^2 c^2 dt^2 + \mathbf{a}(y, t)^2 [dr^2 + r^2 d\Omega_2].$$


$$\mathbf{a}(y, t)^2 = a(t)^2 + y^2 \frac{\dot{a}(t)^2}{c^2} \pm 2y \sqrt{a(t)^2 \frac{\dot{a}(t)^2}{c^2} + I},$$

$$\mathbf{n}(y, t) = \frac{\partial_t \mathbf{a}(y, t)}{\dot{a}(t)}. \quad (\kappa_5 c^2) \rho_{\mathcal{H}} = - \frac{3 \partial_y \mathbf{a}(y, t)}{\mathbf{a}(y, t)} \Big|_{y=0} = 3 \left[\frac{H(t)^2}{c^2} + \frac{I}{a(t)^4} \right]^{1/2},$$

Summary & Outlook

$$\frac{\eta}{s} \simeq \frac{1}{4\pi} \frac{\hbar}{k_B T_c}$$

$$\tau_c^{-1} \simeq \frac{k^2}{4\pi T_c}$$





$$\Omega_D^2 \simeq \frac{1}{2} \Omega_\Lambda (\Omega_D - \Omega_B)$$

$$\frac{H^2}{H_0^2} \simeq \frac{\Omega_B}{a^3} + \sqrt{\Omega_\Lambda \left(\frac{H^2}{H_0^2} + \frac{\Omega_I}{a^4} \right)}$$


Quark Critical Liquid
QGP in RHIC ['08] & LHC ['16]

Cosmological Fluid
Dark Matter['70s] & Energy['90s]



$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} - \frac{1}{L} (\mathcal{K}_{\mu\nu} - \mathcal{K} g_{\mu\nu}) = \kappa_4 T_{\mu\nu}$$

**Thanks for All
Your Attention!**



Ref: 1712.09326 [Cai, Sun, Zhang]

REPORT ON JHEP_066P_0418

DATE: JUNE 6, 2018

AUTHOR(S): RONG-GEN CAI, SICHUN SUN, YUN-LONG ZHANG

TITLE: Emergent Dark Matter in Late Universe on Holographic Screen

RECEIVED: 2018-04-09 14:35:06.0

This took quite a long while. Given the gravity of this affair, I approached two quite distinguished referee's. You may guess who is the second referee who let me wait quite long. Even he likes it! In principle your work has survived scrutiny of the most serious kind. But both referee's come up with the same recommendation: given that it is a very serious and potentially highly consequential contribution, please put in some extra effort to improve the quality of the text and the coherence of the line of arguments. Such an extra investment is just worthwhile. Surely you should exploit the suggestion of the first referee associated with the DM equation of state.

Report of referee 1

Dear Editor,

I recommend that the article "Emergent Dark Matter in Late Universe on Holographic Screen" by R.-G. Cai, S. Sun, and Y.-L. Zhang be published in

Report of referee 2

This paper proposes an holographic approach to explain the dark contributions to the cosmological energy density: dark energy as well as dark matter. The central idea is appealing: the universe is modeled as a brane embedded in a higher dimensional spacetime, in which only the ordinary (=baryonic) matter (and radiation) are described by a stress energy tensor on the brane.

The dark components of the stress energy tensor are in this approach induced by the extrinsic curvature components associated with the brane embedding.

The paper logically consists of three parts. In the first part a relation between the energy densities associated with dark energy, dark matter and baryonic matter is derived in a simplified toy model. Despite the simplicity of the model, it is striking that this relation appears to hold in the current late universe to a very good degree. This part is well presented and logically coherent.

OutLine for Holographic Hydrodynamics

- I. Gravity, Black Holes and Holography
- II. Hydrodynamics and Membrane Fluid
- III. Rindler Horizon and Relevant topics

Rindler Fluid with Weak Momentum Relaxation

JHEP 1801 (2018) 058 [arXiv: [1705.05078](https://arxiv.org/abs/1705.05078)]

by S. Khimphun, B.-H. Lee, C. Park, **Yun-Long Zhang**



S. Khimphun(Hangyang), **B.-H. Lee** (Sogang), **C.-Y. Park** (GIST)

Emergent Dark Matter in Late Universe on Holographic Screen


by R.-G. Cai, S.-C. Sun, **Yun-Long Zhang** [arXiv: [1712.09326](https://arxiv.org/abs/1712.09326)]

R.-G. Cai
(ITP-CAS)



Holographic Bell Inequality [arXiv: [1612.09513](https://arxiv.org/abs/1612.09513)]

by J.-W. Chen, S.-C. Sun, **Yun-Long Zhang** [submitted to PRD]



J. -W. Chen(NTU)

S. -C. Sun(NTU)

Y. -L. Zhang(APCTP)