

A traditional Chinese yin-yang symbol is positioned in the top left corner. It is rendered in black and white with a textured, brush-stroke-like appearance, set against a white background.

CEPC Higgs Combination

Zhang Kaili¹, Cui Zhenwei², Wang Jin³, Liu Zhen⁴

1. Institute of High Energy Physics

2. Peking University

3. University of Sydney

4. University of Pittsburgh

Workshop on the Circular Electron Positron Collider

2018.05.25

Rome


OUTLINE



- Why and How we do combination
- $H \rightarrow \mu\mu$ study
- Results of $\sigma(ZH) * \text{Br}$
- κ Framework
- Summary

Why Combination?

- Uniformed, simultaneous statistical framework
 - Get likelihood scan result Robust & Reliable;
 - Correctly consider the correlations between individual channels
 - bb/cc/gg; ZH bkg; WW fusion; width.....
 - Extensibility
 - systematic uncertainties, theoretic assumptions.....
- Currently, with MC sample (always $\mu = 1$)
 - Build Asimov(1007.1727) data from signal and bkg spectrum
 - To fit the estimated precisions of $\sigma * Br$, and κ .
 - Calculation like Significance / Upper limit also obtained;
 - Can do more with observed data in the future.
 - Results shown in Layout=CEPC_v1, ECM=250GeV, B=3.5T.



Fit techniques

- Input: Various. binned/unbinned, 1d/2d spectrum used.
- Parameter of interest: $\sigma * Br$, Higgs coupling κ
 - $N_{total} = \mu * S + B$, $\mu = \sigma * Br = \frac{\kappa\kappa}{\Gamma}$ and share the same relative uncertainty;
- Nuisance parameter: Represents systematic uncertainties
 - $\sigma(ZH)$: 0.5%; $\sigma(Lumi)$: 0.1%; more NPs can be introduced in the future.
 - currently results are **all** determined by **statistical uncertainty**.
- PDF: To describe the shape of the spectrum.
 - signal: Double sided Crystal ball; bkg: 2rd-order poly exponential.
 - RooHistPdf/RooKeysPdf used for some channels;
- Algorithm: Likelihood Scan
 - Asymmetric result, from Minuit2; $\pm 1\sigma$ deviation from profile likelihood

Fit techniques

- For each channel

- Input observables from MC sample.

- Build combined S+B Pdf

$$\text{Tot} = N_{bb} * \text{Pdf} + N_{cc} * \text{Pdf}_{cc} + \dots + N_{bkg} * \text{Pdf}_{bkg}$$

- For event number N_{bb} :

- When measure $\sigma * Br$,

$$N_{bb} = N_{bb_SM} * \mu_{bb}$$

N_{bb_SM} directly from event yield ($5ab^{-1}$)

- When measure Br ,

$$N_{bb} = N_{bb_SM} * \frac{Br}{Br_{SM}} * \frac{\sigma(ZH)}{\sigma(ZH)_{SM}}$$

$\Delta(\sigma(ZH)) = 0.50\%$

- When measure κ ,

$$N_{bb} = N_{bb_SM} * \kappa_Z^2 (\kappa_W^2) * \kappa_b^2 / \Gamma_H$$

- Channel share the same μ .

$Z \rightarrow ee, \mu\mu, qq, \nu\nu$, share the same μ_{bb}

- Events number N_{bb} is float

and the Pdf **shape fixed** all the time.

- Use Combined pdf to make Asimov data

- Scan the likelihood and obtain the 1σ deviation

Channels Table


All channels scaled to $5ab^{-1}$



Signal		Precision	Signal		Precision	Signal		Precision
Z	H		Z	H		Z	H	
H->qq			H->WW			H->ZZ		
ee	bb	1.6%	ee	l ν l ν	9.2%	$\nu\nu$	$\mu\mu$ qq	8.2%
	cc	23.6%		evqq	4.6%	$\nu\nu$	eeqq	35.2%
	gg	13.3%		$\mu\nu$ qq	3.9%	$\mu\mu$	$\nu\nu$ qq	7.3%
$\mu\mu$	bb	1.1%	$\mu\mu$	l ν l ν	7.3%	ee	eeqq	35.1%
	cc	14.8%		evqq	4.0%	ee	$\mu\mu$ qq	23.0%
	gg	8.0%		$\mu\nu$ qq	4.0%	ZH bkg contribution		19.4%
qq	bb	0.5%	$\nu\nu$	qqqq	2.0%	v ν H(WW fusion)		
	cc	11.9%		evqq	4.7%	$\nu\nu$	bb	3.1%
	gg	3.9%		$\mu\nu$ qq	4.2%	H-> $\mu\mu$		
v ν	bb	0.4%	qq	l ν l ν	11.3%	qq	$\mu\mu$	15.9%
	cc	3.9%		l ν qq	2.2%(ILC)	ee		
	gg	1.5%		ZH bkg contribution		3.0%		
H-> $\tau\tau$			H-> $\gamma\gamma, Z\gamma$			$\nu\nu$		
ee	$\tau\tau$	2.8%	$\mu\mu+\tau\tau$	$\gamma\gamma$	41.0%	H->Invisible		Br, Upper
$\mu\mu$		2.8%	$\nu\nu$		13.7%	qq	ZZ(v $\nu\nu\nu$)	0.8%
qq		1.0%	qq		10.3%	ee		0.6%
$\nu\nu$		3.1%	$\nu\nu$		Z γ (qq γ)	21.2%		$\mu\mu$



Treatment for ZH bkg

- In individual analysis, other ZH processes are tagged as bkg;
 - Signal in one channel can be bkg for another channel.
 - Should taken into account in combination.
 - $Z \rightarrow \mu\mu, H \rightarrow \tau\tau$, the main bkg is $H \rightarrow WW$.
 - These WW events should be considered in μ_{WW} .
 - Standalone WW channel 1.2% improved to 1.0% this way;
 - Combined fit for $H \rightarrow bb/cc/gg/ww/zz$ hadronic decay, Fully correlated.



Correlation: $\nu\nu H \rightarrow bb$

- 2d fit M_{jj}^{reco} & $\text{Cos } \theta_{jj}$
- Correlated with ZH process;
 - Fix ZH process, Initial error is 2.89%.
 - But must consider the uncertainty from ZH process.
- Use the likelihood from $Z \rightarrow ee/\mu\mu/qq, H \rightarrow bb$ to constrain
 - Already have the form of μ_{ZH} , no assumption made;
 - $\nu\nu H \rightarrow bb$ and $ZH \rightarrow bb$ share the anti-correlation **-46%**. (-34% in ILC(1708.08912))
- Simultaneous Fit 3.1% ; consistent with individual study 3.1%.
 - Corresponding to this, $ZH \rightarrow bb$ precision 0.33%.
 - $\sigma(\nu\nu H)$ precision 3.16%.



Correlation: Higgs width

- In Pre_CDR, width determined by

$$\Gamma_H = \frac{\Gamma_{H \rightarrow ZZ}}{Br(H \rightarrow ZZ)} \propto \frac{\sigma(ZH)}{Br(H \rightarrow ZZ)} \quad \text{and} \quad \Gamma_H = \frac{\Gamma_{H \rightarrow bb}}{Br(H \rightarrow bb)} \propto \frac{\sigma(\nu\nu H \rightarrow \nu\nu bb)}{Br(H \rightarrow bb)Br(H \rightarrow WW)}$$


- If two independent: 2.83% (consistent with pre_CDR, which gives 2.8%)
- But width correlated with all channels
 - Like correlation like $\nu\nu H \rightarrow \nu\nu bb$ and $ZH \rightarrow bb$ -46% not included -> would worsen the result
- Combined fit in 10κ framework:

$$\Delta(\Gamma_H) = 3.2\%$$

H → μμ, 3.5T, Full simulation

- Z → ee

bkg shape all after smoothing.
(10~100x bkg events used)




Cutflow	signal	ZZ	WW	ZZorWW	SingleZ	2f
Init	4.7	18	0	9	22672	8
$120 < M_{\mu^+\mu^-} < 130$	4.3	0	0	0	747	0
$89 < M_{reco}^{\mu^+\mu^-} < 94$	3.0	0	0	0	56	0
$138 < E_{\mu^+\mu^-} < 140$	2.2	0	0	0	8	0
efficiency	46.81%					

Bkg: Sz(l)e.l0mu;

- Z → mm combination to minimize

$$\delta = \left(\frac{pair1.M}{\Delta Z}\right)^2 + \left(\frac{pair2.M}{\Delta H}\right)^2$$

- $\Delta Z = 1.5, \Delta H = 0.75$




Category	signal	ZZ	WW(SW)	ZZorWW	SingleZ	2f
Preselection	6.6	17631.0	0	0	0	0.0
$120 < E_{\mu^+\mu^-} < 130$	6.0	1685.2	0	0	0	0.0
$90.6 < M_{recoil_\mu} < 93.4$	3.9	128.8	0	0	0	0.0
$90.2 < M_{\mu^+\mu^-}(Z) < 92.8$	3.2	58.1	0	0	0	0.0
$\cos_{\mu^+\mu^-}(H) < -0.603$	3.2	50.0	0	0	0	0.0
$\cos_{\mu^+\mu^-}(Z) < -0.364$	3.2	47.0	0	0	0	0.0
$138.0 < E_{\mu^+\mu^-}(H) < 139.8$	3.0	15.5	0	0	0	0.0
$P_{T_{\mu^+\mu^-}}(H) < 62.5$	3.0	14.7	0	0	0	0.0
efficiency	45.5%					

Bkg: ZZ(l).4mu;


$ZH \rightarrow \nu\nu\mu\mu, qq\mu\mu$

- $Z \rightarrow \nu\nu$
 - 38%



Cutflow	signal	ZZ	WW	ZZorWW	SingleZ	2f
Init	41.7	34901	121952	489686	25619	1635887
$120 < M_{\mu^+\mu^-} < 130$	38.4	382	16677	56029	315	49490
$MET_{\tilde{t}} > 8.5$	37.9	291	16264	53740	305	8600
$89 < M_{reco}^{\mu^+\mu^-} < 94$	28.1	96	834	2034	79	184
$\cos\theta_{\mu^+} > 0, \cos\theta_{\mu^-} < 0$	9.1	22	11	86	17	9
efficiency	21.82%					

- $Z \rightarrow qq$
 - 17%



Cutflow	signal	ZZ	WW	ZZorWW	SingleZ	2f
Init	156.3	390775	183751	463361	101164	63217
$120 < M_{\mu^+\mu^-} < 130$	141.6	3786	181	227	244	100
$M_{j1} > 4.2, M_{j2} > 2.8$	133.0	3216	111	0	9	60
$M_{jj} > 76.0$	127.5	2917	2	0	8	59
$89 < M_{reco}^{\mu^+\mu^-} < 94$	86.1	1106	0	0	0	0
efficiency	55.08%					

- Combined: 15.9%


Main bkg: ZZ(sl)mu.down, ZZ(sl)mu.up

- Considering the scheduled time, CEPC could be the first detector to see this process.


$qq\mu\mu$, 3T & 3.5T, full simulation



H mass



Z mass



Events normalized, no significant difference in mean value and resolution.



$Z \rightarrow qq, H \rightarrow \mu\mu$

Comparison



3T: 18.6%

3.5T: 17.4%



when the magnet field reduced,
2.8% signal, 4% bkg events would be lost in reconstruction.
3.1% signal, 4% bkg events would fail in preselection. (Good muon selection)
-> Signal: 81; Bkg: 1006;
Considering these, precision has reduced from 17.4% to 18.6%.
There is a slight performance downgrade from 3.5T to 3T.

Fit result of $\sigma(ZH) * Br$

(5ab ⁻¹)	Pre_CDR	Current 2018.5	ILC 250	Fcc-ee
$\sigma(ZH)$	0.51%	0.50%	1.2%	0.40%
$\sigma(ZH) * Br(H \rightarrow bb)$	0.28%	0.28%	0.6%	0.2%
$\sigma(ZH) * Br(H \rightarrow cc)$	2.2%	3.5%	3.9%	1.2%
$\sigma(ZH) * Br(H \rightarrow gg)$	1.6%	1.4%	3.3%	1.4%
$\sigma(ZH) * Br(H \rightarrow WW)$	1.5%	1.0%	3.0%	0.9%
$\sigma(ZH) * Br(H \rightarrow ZZ)$	4.3%	5.0%	8.4%	3.1%
$\sigma(ZH) * Br(H \rightarrow \tau\tau)$	1.2%	0.8%	2.0%	0.7%
$\sigma(ZH) * Br(H \rightarrow \gamma\gamma)$	9.0%	8.1%	16%	3.0%
$\sigma(ZH) * Br(H \rightarrow \mu\mu)$	17%	16%	46.6%	13%
$\sigma(vvH) * Br(H \rightarrow bb)$	2.8%	3.1%	11%	2.4%
$Br_{upper}(H \rightarrow inv.)$	0.28%	0.42%	0.4%	0.50%
$\sigma(ZH) * Br(H \rightarrow Z\gamma)$	\	4σ(21%)		

ILC:	1310.0763
FCC-ee:	1308.6176

Difference from Pre_CDR

(5ab ⁻¹)	Pre_CDR	Combined
$\sigma(ZH)$	0.51%	0.50%
$\sigma(ZH) * \text{Br}(H \rightarrow bb)$	0.28%	0.28%
$\sigma(ZH) * \text{Br}(H \rightarrow cc)$	2.2%	3.5%
$\sigma(ZH) * \text{Br}(H \rightarrow gg)$	1.6%	1.4%
$\sigma(ZH) * \text{Br}(H \rightarrow WW)$	1.5%	1.0%
$\sigma(ZH) * \text{Br}(H \rightarrow ZZ)$	4.3%	5.0%
$\sigma(ZH) * \text{Br}(H \rightarrow \tau\tau)$	1.2%	0.8%
$\sigma(ZH) * \text{Br}(H \rightarrow \gamma\gamma)$	9.0%	8.1%
$\sigma(ZH) * \text{Br}(H \rightarrow \mu\mu)$	17%	16%
$\sigma(vvH) * \text{Br}(H \rightarrow bb)$	2.8%	3.1%
$\text{Br}_{\text{upper}}(H \rightarrow \text{inv.})$	0.28%	0.42%
$\sigma(ZH) * \text{Br}(H \rightarrow Z\gamma)$	\	4σ(21%)

- CrossSection: minor update.
- bb,cc,gg: due to flavor tagging algorithm, The template gives b/c likeness, updated algorithm has less cc candidate events left.
- WW: more subchannels studied and ZH bkg contribution.
- ZZ: the extrapolation in Pre_CDR from FCC-ee too optimistic.
- $\tau\tau$: τ finding algorithm updated.
- $\gamma\gamma$: different estimation from full/fast simulation.
- vvH : consider the correlation
- $H \rightarrow \text{invisible}$: Pre_CDR studied an exotic decay $H \rightarrow \chi_1\chi_1$ and assuming 200fb⁻¹, gives 0.28%.
- Now we study the upper limit of $H \rightarrow ZZ \rightarrow vvvv$.

κ defined as the ratio of the Higgs coupling to SM expects.

$$\kappa_f = \frac{g(hff)}{g(hff; SM)}, \quad \kappa_V = \frac{g(hVV)}{g(hVV; SM)}$$

κ Framework

- Model independent implication

- Detector's benchmark;

Constrain to new physics models;

- In CEPC

- We have $\sigma(ZH) = 0.5\%$ constrain $\sigma(\kappa_Z) < 0.25\%$.
- For Production, ZH & WW fusion process, all contribute to $\kappa_Z^2; \kappa_W^2;$
- For Partial decay, no top quark κ_t like: $\kappa_Z^2, \kappa_W^2, \kappa_b^2, \kappa_c^2, \kappa_g^2, \kappa_\tau^2, \kappa_\gamma^2, \kappa_\mu^2, \dots$
- For Total width Γ_H . $\Gamma_H = \Gamma_{SM} + \Gamma_{BSM}$.
- If we assume no exotic decay, Γ_{SM} can be resolved as: all κ correlated this way;

$$\Gamma_{SM} = 0.2137\kappa_W^2 + 0.02619\kappa_Z^2 + 0.5824\kappa_b^2 + 0.08187\kappa_g^2 + 0.002270\kappa_\gamma^2 + 0.06294\kappa_\tau^2 + 0.02891\kappa_c^2$$

- $Z \rightarrow \mu\mu, H \rightarrow \tau\tau$ channel, the signal will be $\kappa_Z^2\kappa_\tau^2/\Gamma_H$; For $\nu\nu H \rightarrow bb$, it's $\kappa_W^2\kappa_b^2/\Gamma_H$

Fit result of κ


In different interpretation,
Higgs width can be independent or resolved by branch ratio.

	10κ	Pre_CDR	7κ	Pre_CDR
κ_b	1.6%	1.3%	1.0%	1.2%
κ_c	2.3%	1.7%	2.1%	1.6%
κ_g	1.6%	1.5%	1.2%	1.5%
κ_γ	4.4%	4.7%	4.3%	4.7%
κ_τ	1.6%	1.4%	1.1%	1.3%
κ_Z	0.21%	0.26%	0.17%	0.16%
κ_W	1.4%	1.2%	1.0%	1.2%
κ_μ	8.1%	8.6%		
Br_{inv}	0.42%	0.28%	From 10 κ to 7 κ , we assume <ul style="list-style-type: none"> • No exotic decay Γ_{BSM} • Drop Br_{inv} • $\kappa_\mu = \kappa_\tau$ 	
Γ_H	3.3%	2.8%		

Integration to HL-LHC

The improvement of κ_γ from $Br_{ZZ}/Br_{\gamma\gamma} = 4\%$

	10-parameter fit		7-parameter fit	
	CEPC	+HL-LHC	CEPC	+HL-LHC
Γ_h	3.2	2.5	-	-
κ_b	1.6	1.2	1.0	0.9
κ_c	2.3	2.0	2.1	1.9
κ_g	1.6	1.2	1.2	1.0
κ_W	1.4	1.1	1.0	0.9
κ_τ	1.6	1.2	1.1	1.0
κ_Z	0.21	0.21	0.17	0.16
κ_γ	4.4	1.7	4.3	1.7
κ_μ	8.1	4.9	-	-
BR_{inv}	0.31	0.31	-	-




ATL-PHYS-PUB-2014-016

*: here Br_{inv} for BSM.

Compared to ILC_(1710.07621)

ILC used more aggressive κ_γ , by ratio $Br_{ZZ}/Br_{\gamma\gamma} = 2\%$



Correlation of κ

For each entry,
upper one is CEPC result
lower one is CEPC+HL-LHC result.



7-parameter fit Correlation

K_b	100.	-35.	-52.	-59.	-44.	66.	-9.6
	-32.	100.	2.5	17.	2.9	-38.	2.4
K_c	-32.	100.	-3.4	16.	2.8	-36.	0.92
	-52.	2.5	100.	6.7	6.8	-39.	2.2
K_g	-51.	-3.4	100.	1.6	4.3	-28.	0.55
	-59.	17.	6.7	100.	-22.	-55.	2.8
K_W	-56.	16.	1.6	100.	-21.	-56.	1.1
	-44.	2.9	6.8	-22.	100.	1.8	-0.85
K_τ	-43.	2.8	4.3	-21.	100.	1.3	-0.32
	66.	-38.	-39.	-55.	1.8	100.	-4.7
K_Z	63.	-36.	-28.	-56.	1.3	100.	-7.5
	-9.6	2.4	2.2	2.8	-0.85	-4.7	100.
K_γ	-3.7	0.92	0.55	1.1	-0.32	-7.5	100.
	K_b	K_c	K_g	K_W	K_τ	K_Z	K_γ

10-parameter fit Correlation

K_b	100.	-4.4	-7.6	-19.	4.2	78.	<0.1	<0.1	<0.1	-92.
	-4.2	100.	-19.	-9.3	2.3	77.	<0.1	<0.1	<0.1	-91.
K_c	-4.2	100.	-18.	-8.6	2.2	-6.1	<0.1	<0.1	<0.1	2.7
	-7.6	-19.	100.	-25.	9.2	3.2	<0.1	<0.1	<0.1	-5.8
K_g	-6.1	-18.	100.	-19.	7.4	8.7	<0.1	<0.1	<0.1	-13.
	-19.	-9.3	-25.	100.	-22.	-6.9	<0.1	<0.1	<0.1	5.7
K_W	-19.	-8.6	-19.	100.	-21.	-7.8	<0.1	<0.1	<0.1	5.5
	4.2	2.3	9.2	-22.	100.	25.	<0.1	<0.1	<0.1	-31.
K_τ	4.2	2.2	7.4	-21.	100.	24.	<0.1	<0.1	<0.1	-31.
	78.	-6.1	3.2	-6.9	25.	100.	2.8	1.5	<0.1	-83.
K_Z	77.	-5.7	8.7	-7.8	24.	100.	-4.7	-0.61	<0.1	-83.
	<0.1	<0.1	<0.1	<0.1	<0.1	2.8	100.	<0.1	<0.1	-3.4
K_γ	<0.1	<0.1	<0.1	<0.1	<0.1	-4.7	100.	<0.1	<0.1	-1.3
	<0.1	<0.1	<0.1	<0.1	<0.1	1.5	<0.1	100.	<0.1	-1.8
K_μ	<0.1	<0.1	<0.1	<0.1	<0.1	-0.61	<0.1	100.	<0.1	-1.1
	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	100.	<0.1
Br_{inv}	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	100.	<0.1
	-92.	2.7	-5.8	5.7	-31.	-83.	-3.4	-1.8	<0.1	100.
K_Γ	-91.	2.5	-13.	5.5	-31.	-83.	-1.3	-1.1	<0.1	100.
	K_b	K_c	K_g	K_W	K_τ	K_Z	K_γ	K_μ	Br_{inv}	K_Γ

Summary

	Current
$\sigma(ZH)$	0.50%
$\sigma(ZH) * Br(H \rightarrow bb)$	0.28%
$\sigma(ZH) * Br(H \rightarrow cc)$	3.5%
$\sigma(ZH) * Br(H \rightarrow gg)$	1.4%
$\sigma(ZH) * Br(H \rightarrow WW)$	1.0%
$\sigma(ZH) * Br(H \rightarrow ZZ)$	5.0%
$\sigma(ZH) * Br(H \rightarrow \tau\tau)$	0.8%
$\sigma(ZH) * Br(H \rightarrow \gamma\gamma)$	8.1%
$\sigma(ZH) * Br(H \rightarrow \mu\mu)$	16%
$\sigma(vvH) * Br(H \rightarrow bb)$	3.1%
$Br_{upper}(H \rightarrow inv.)$	0.42%
$\sigma(ZH) * Br(H \rightarrow Z\gamma)$	4σ(21%)

	10 κ	7 κ
κ_b	1.6%	1.0%
κ_c	2.3%	2.1%
κ_g	1.6%	1.2%
κ_γ	4.4%	4.3%
κ_τ	1.6%	1.1%
κ_Z	0.21%	0.17%
κ_W	1.4%	1.0%
κ_μ	8.1%	
Br_{inv}	0.42%	
Γ_H	3.2%	

	10-parameter fit		7-parameter fit	
	CEPC	+HL-LHC	CEPC	+HL-LHC
Γ_h	3.2	2.5	–	–
κ_b	1.6	1.2	1.0	0.9
κ_c	2.3	2.0	2.1	1.9
κ_g	1.6	1.2	1.2	1.0
κ_W	1.4	1.1	1.0	0.9
κ_τ	1.6	1.2	1.1	1.0
κ_Z	0.21	0.21	0.17	0.16
κ_γ	4.4	1.7	4.3	1.7
κ_μ	8.1	4.9	–	–
BR_{inv}	0.31	0.31	–	–

- Updated fit results of CEPC Higgs are shown.
- Correlations are taken in consideration in the simultaneous framework.
- To be used in the CDR and white paper.

backup

Individual analysis

$$B_{likeness} = \frac{b_{j_1} b_{j_2}}{b_{j_1} b_{j_2} + (1 - b_{j_1})(1 - b_{j_2})}$$


- Template fit: Flavor tagging algorithm

- $Z \rightarrow ee \mu\mu qq vv, H \rightarrow bb/cc/gg$ are studied.
- 2D fit, with dijets' b/c likeness; mass info not used;
- 7 parts, $Tot=bb+cc+gg+ww+zz+tt+bkg_{sm}$.
- Build individual pdf by MC, then fit to determine fraction.
- the shape of bkg is **fixed**.
 - Which means we have a wonderful understanding with bkg,
 - may be more suitable for CEPC.
- ToyMC test to get precision
- Now plan to use 3d fit in IH;
- Systematic uncertainties ongoing;


Scan	μ_{bb}	μ_{cc}	μ_{gg}
eeH	1.3%	15.0%	8.2%
mmH	1.0%	11.3%	5.5%
qqH	0.5%	17%	7.2%
vvH	0.4%	3.9%	1.6%
Combined	0.28%	3.48%	1.44%

	preCDR	Now
$\tau\tau$	1.2%	0.81%

- Pre_CDR concludes the precision 1.2% but no description.
- Develop LICH to identify lepton. Eff>99%
- Signal and ZH events(Main WW) share the same shape
 - use $\log_{10}(D_0^2 + Z_0^2)$ fit to separate signal
 - Impact parameter, Distance from beam spot



	BR ($H \rightarrow \tau\tau$)	$\delta(\sigma \times BR) / (\sigma \times BR)$
$\mu\mu H$	6.40	2.68%
eeH (extrapolated)	6.37	2.72%
$\nu\nu H$	6.26	4.38%
qqH	6.23	0.93%
combined	6.28	0.81%





	preCDR	Now
WW	1.5%	1.0%

Signal		Precision
Z	H	
H->WW		
ee	lvlv	9.2%
	evqq	4.6%
	$\mu\nu$ qq	3.9%
$\mu\mu$	lvlv	7.3%
	evqq	4.0%
	$\mu\nu$ qq	4.0%
$\nu\nu$	qqqq	2.0%
	evqq	4.7%
	$\mu\nu$ qq	4.2%
	lvlv	11.3%
qq	lvqq	2.2%(ILC)
ZH bkg contribution		3.0%

- Currently have 17 channels of WW

Green: studied

	Z	ee	$\mu\mu$	$\nu\nu$	qq
WW	ev+ev				
	$\mu\nu+\mu\nu$				
	ev+ $\mu\nu$				
	ev+qq				
	$\mu\nu$ +qq				
	qq+qq				



	preCDR	Now
ZZ	4.3%	5.0%

Channel	Precision	Comment
$\sigma(Z(\nu\bar{\nu})H + \nu\bar{\nu}H) \times BR(H \rightarrow ZZ)$	6.9%	CEPC Fast Simulation
$BR(H \rightarrow ZZ^*)$	4.3%	Extrapolation from FCC-ee [36]





- Pre_CDR's result from extrapolating the FCC-ee.
- Now has 5 channels clear and easy to study

Signal		Precision
Z	H	
H->ZZ		
$\nu\nu$	$\mu\mu qq$	8.2%
$\nu\nu$	$ee qq$	35.2%
$\mu\mu$	$\nu\nu qq$	7.3%
ee	$ee qq$	35.1%
ee	$\mu\mu qq$	23.0%
ZH bkg contribution		19.4%


Green: studied

	Z	ee	$\mu\mu$	$\nu\nu$	qq
ZZ	ee+qq				
	$\mu\mu+qq$				
	$\nu\nu+qq$				
	ll+ll				
(Invi)	$\nu\nu+\nu\nu$				
	qq+qq				
	ll+ $\nu\nu$				





$\gamma\gamma$

	preCDR	Now
$\gamma\gamma$	9.0%	8.1%





Signal		Precision
Z	H	
H → $\gamma\gamma$		
$\mu\mu + \tau\tau$	$\gamma\gamma$	24.8%
$\nu\nu$		11.7%
qq		12.8%



	preCDR	Now
<i>invisible</i>	0.28%	0.42%

$H \rightarrow invisible$


- Moxin studied $H \rightarrow ZZ \rightarrow vvvv$
 - Large irreducible bkg, use BDT and seek upper limit.
 - Huge fluctuation, use Asimov Data to get correct fit result.
 - precision 148%, upper limit for Br: 0.42%
 - Upper limit for BSM $H \rightarrow invisible$: 0.31%



	Precision	significance	Br Upper limit
Z $\rightarrow ee$	350%		0.84%
Z $\rightarrow mm$	242%		0.62%
Z $\rightarrow qq$	226%		0.59%
Combined	148%	0.68σ	0.42%

$Z\gamma$

- $Z \rightarrow qq, H \rightarrow Z\gamma \rightarrow qq\gamma$ studied;
 - Pre_CDR not conclude;
 - Take $m_{Z\gamma-Z}$ as observable;
 - 4σ significance; Precision about 21%.



- $e\mu, ee$ process studied.
 - Since low stats and no clear ratio, not taken into fit model.