

# Using AMPT Study: The Directed Flow in Au+Au Collisions at $\sqrt{s_{NN}} = 7.7 - 200$ GeV


*Kishora Nayak<sup>1</sup>, Shusu Shi<sup>1</sup>, Nu Xu<sup>1,2</sup>*

*<sup>1</sup>Central China Normal University, Wuhan, China*

*<sup>2</sup>Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou, China*

**The 7<sup>th</sup> Asian Triangle heavy-Ion Conference (ATHIC 2018)**

*November 3-6, 2018, University of Science and Technology of China, Hefei, AnHui, China*



# Outline

- ☑ Introduction and motivation
- ☑ AMPT analysis details
- ☑ Results
  - Directed flow ( $v_1$ ) of identified hadrons
  - $v_1$  comparison with experimental data
  - Partons'  $v_1$
- ☑ Summary and outlook

# Introduction



Invariant yield, 
$$E \frac{d^3N}{dp^3} = \frac{1}{2\pi} \frac{d^2N}{p_T dp_T dy} \left( 1 + \underset{\substack{\uparrow \\ \text{Isotropic}}}{2v_1} \cos(\phi - \psi_R) + \underset{\substack{\uparrow \\ \text{Directed}}}{2v_2} \cos 2(\phi - \psi_R) + \dots \right)$$

**Directed flow ( $v_1$ )** is quantified by the first harmonic  $v_1$ :

$$v_1 = \langle \cos(\phi - \psi_R) \rangle \quad \phi = \tan^{-1} \left( \frac{p_y}{p_x} \right)$$

**Directed flow,  $v_1$ :**

- ▶ It is generated during the nuclear passage time ( $2R/\gamma \sim 0.1$  fm/c) and hence it probes the earliest stage of collision dynamics (Ref: arXiv:1407.5003)
- ▶ The spectators deflected from dense reaction zone are sensitive to the **pressure** and also have strong sensitivity towards the **EoS**
- ▶ It describes the sideward motion of the particles within the reaction plane



# Motivation

STAR

**An experimental approach of AMPT-SM model analysis**

**with reference to STAR data paper:**

**Coalescence sum rule:** If deconfined quarks already acquired azimuthal anisotropy, then coalescence leads to  $v_n$  of the resulting hadrons being the summed  $v_n$  of their constituent quarks

$$(v_n)_{(hadron)} = \sum (v_n)_{constituent\ quark}$$

**(i) Coalescence sum rule for  $v_1$ :**


$$s = K^- (\bar{u}s) - \frac{1}{3} \bar{p} (\bar{u}\bar{u}\bar{d})$$

It assumed,  $\bar{u}$ ,  $\bar{d}$  quarks have same flow and so  $s$ ,  $\bar{s}$  quarks

**(ii) Disentangle the effect of transported and produced quarks**


$$v_{1\ p} = r(y)v_{1\ \bar{p}} + [1 - r(y)]v_{1\ net\ p} ; \quad r(y) = \bar{p}/p$$

*Essential for understanding the particle production mechanism at various energies*



Phys. Rev. Lett. 120, 062301 (2018)  
(STAR Collaboration)

# AMPT analysis details




- No. of events : ~1 million for AMPT-String Melting
- AMPT-SM. : 1.5 mb,  $c\tau = 0.6$  fm/c
- Energy (GeV) : 7.7, 11.5, 14.5, 19.6, 27, 39, 54.4, 62.4, 200
- Hadrons :  $\pi^+$ ,  $\pi^-$ ,  $K^+$ ,  $K^-$ ,  $K_S^0$ ,  $p$ ,  $\bar{p}$ ,  $\Lambda$ ,  $\bar{\Lambda}$ ,  $\Xi^-$ ,  $\Xi^+$
- Partons. :  $u$ ,  $\bar{u}$ ,  $d$ ,  $\bar{d}$ ,  $s$ ,  $\bar{s}$


*The  $p_T$  cut used here for identified hadrons are similar to experimental data analysis*

*All the particles and anti-particles are identified via their corresponding Pythia-ID (PID)*

# AMPT-SM: $v_1$ vs $y$ [10-40%]



# Compared with data: $v_1$ vs $y$ [10-40%]




Black solid circles are experimental data and color markers are AMPT-SM

Phys. Rev. Lett. 120, 062301 (2018)



# AMPT-SM: Partons' $v_1$



- ✓ Strong rapidity and energy dependence of quarks and anti-quarks
- ✓ Difference between quark and anti-quark  $v_1$  value increases with decrease in energy
- ➔ Similar  $v_1$  values for quark and anti-quark at 200 GeV unlike 7.7 GeV



$$v_1(y) = Fy + F_3y^3$$

Slope ( $dv_1/dy$ )

Fluctuation




# Produced partons slope vs $\sqrt{s_{NN}}$




- ✓ Difference between quark and anti-quark  $v_1$  value increases with decrease in energy
- ✓ Similar  $v_1$  values for **u** and **d** quarks unlike **s** quark
- ✓ Negative  $v_1$  values of quarks and anti quarks decreases with increase in energy
- ➔ At 200 GeV all quarks and anti-quarks have positive  $v_1$  values unlike 7.7 GeV

# AMPT-SM: slope vs $\sqrt{s_{NN}}$ [10-40%]

STAR




$$v_{1 \text{ net } p} = \frac{v_{1 p} - r(y)v_{1 \bar{p}}}{1 - r(y)}$$



Phys. Rev. Lett. 120, 062301 (2018)


✓ Qualitative similarity between net K and data unlike net p, net  $\Lambda$  where no sign change is observed

# AMPT-SM: slope vs $\sqrt{s_{NN}}$ [10-40%]



Produced quarks

Transported quarks



- ✓ In  $\left[ \text{net } p - \frac{1}{3} \bar{p} + s \right]$ , produced u quark is replaced with s quark and  $\bar{u}$  from  $\frac{1}{3} \bar{p}$
- ✓ Transported quarks dominant at lower energies ( $< 14.5$  GeV) and more produced quarks at high energy

# Summary

- ❖ A significant species dependence of rapidity odd component of directed flow from lower (7.7 GeV) to higher energy (200 GeV) in AMPT-SM
- ❖ Higher value of net p, net  $\Lambda$  and net  $K^+$   $v_1$  at 7.7 GeV compared to 200 GeV suggests that both the contribution of transported quarks at lower energy is more compared to high energy
- ❖ *Transported and produced quarks have different  $v_1$  values*
- ❖ *Strange and anti-strange quark don't have the same flow over all energies as assumed in experiment*
- ❖ *Particle production via quark coalescence seems better at higher energies*


## Outlook

- ❖ AMPT-Default model might give us more insight to understand  $v_1$  at lower energies

**Acknowledgement** : We would like to thank *Prof. Zi-Wei Lin* for his time and in-depth discussion which was very helpful to understand the AMPT results

**Thank you!**

# STAR: Directed flow ( $v_1$ ):



$$v_1(y) = Fy + F_3y^3$$

Slope ( $dv_1/dy$ )

Fluctuation