


Event-plane Dependence of Jet-like Correlations in Au+Au Collisions at 200 GeV in STAR

Kun Jiang

University of Science and Technology of China (USTC)
(for the STAR Collaboration)





U.S. DEPARTMENT OF
ENERGY



PURDUE
UNIVERSITY

Motivation

- Energetic partons lose energy due to interactions in the dense medium
- Flow background removal is challenging in measurements of medium modifications of jets especially on the away-side
- All orders of v_n are possible and need to be subtracted
- Jet modification depends on path length (emission-angle dependent in non-central heavy-ion collisions)



More differentially: Event Shape Engineering

- Possibility to control the initial geometry / fluctuating v_2 by selecting the magnitude of flow vector q_2
- Combination of centrality selection and event shape engineering allows control of the initial geometry while keeping the average energy density (multiplicity) fixed
- Separation of volume effect and geometry effect

$$Q_{2,x} = \sum w_i \cos(2\phi_i) / \sqrt{\sum w_i}$$



$$Q_{2,y} = \sum w_i \sin(2\phi_i) / \sqrt{\sum w_i}$$

$$q_2 = \sqrt{Q_{2,x}^2 + Q_{2,y}^2}$$

w_i : weighting factor

A.M.Poskanzer, S.A.Voloshin,
PRC 58 (1998), 1671-1678

initial eccentricity $\epsilon_2 = \frac{\langle x^2 - y^2 \rangle}{\langle x^2 + y^2 \rangle}$





Event-plane Dependent Two-particle Jet-like Correlations



Liang Zhang, HP2018

Enhanced Away-side Momentum Flow

Projection of away-side p_T onto trigger axis

$$P_x \Big|_{\eta_1}^{\eta_2} = \sum_{\eta_1 < \eta < \eta_2, |\phi - \phi_{trig}| > \pi/2} p_T \cdot \cos(\phi - \phi_{trig}) \cdot \frac{1}{\varepsilon}$$


ε : single-particle acceptance \times efficiency



- For each centrality, cut on the lowest 10% of events to enhance away-side momentum flow \rightarrow "jet" = jet + jet-like hotspots

Methodology for Two-particle Correlations


Trigger particle $|\eta| < 1$



- Away-side: large recoil momentum region opposite to trigger particle
- Analyze correlations in **close-region** and **far-region**, respectively
- Flow contributions to close-region and far-region are equal
→ cancelled in their difference


$$\begin{aligned}
 \text{close-region} &= \text{flow} + \text{near-side "jet"} + \text{away-side "jet"} * \text{fraction}_{\text{close}} \\
 \text{far-region} &= \text{flow} + \text{near-side "jet"} + \text{away-side "jet"} * \text{fraction}_{\text{far}} \\
 \text{diff} &= \text{away-side "jet"} * \text{fraction}
 \end{aligned}$$

Away-side Jet-like Correlation Widths




- Moderate to high $p_{T}^{\text{assoc.}}$ particles: broadening with increasing centrality
- Shape for all $p_{T}^{\text{assoc.}}$ more similar in central than in peripheral collisions

BBC Event-plane Ψ_2 Determination



- Large η gaps between BBCs and TPC
- Minimal non-flow between trigger particles and BBC




$$\Psi_2 = (\tan^{-1} \frac{Q_{2y}}{Q_{2x}}) / 2,$$

$$Q_{2x} = \sum_i w_i \cos(2\phi_i)$$

$$Q_{2y} = \sum_i w_i \sin(2\phi_i)$$

w_i : ADC signal weight



$$R \approx \sqrt{2\cos 2(\Psi_2^{\text{East}} - \Psi_2^{\text{West}})}$$


Raw Event-plane Dependent Correlations

20-60% Au+Au @ 200 GeV
 $3 < p_T^{\text{trig}} < 10 \text{ GeV}/c$
 $1 < p_T^{\text{assoc}} < 2 \text{ GeV}/c$

$$\phi_s = \phi^{\text{trig}} - \Psi_{2,EP}$$


$$\Delta\phi = \phi^{\text{assoc}} - \phi^{\text{trig}}$$

STAR TPC



In-plane

out-of-plane





Away-side Correlation Shape

Close - Far

20-60% Au+Au @ 200 GeV


$3 < p_T^{\text{trig}} < 10 \text{ GeV}/c$
 $1 < p_T^{\text{assoc}} < 2 \text{ GeV}/c$



- Blue curves are Gaussian fits
- Gaussian width (σ) increases with ϕ_s
- Slope = 0.08 ± 0.04
- Poor BBC Ψ_2 resolution ($R = 0.136 \pm 0.002$)
 → correct via unfolding



Resolution-corrected Correlation Functions

Resolution-corrected two-particle correlations:



20-60%
AuAu 200 GeV
 $3 < p_T^{\text{trig}} < 10$ GeV/c
 $1 < p_T^{\text{assoc}} < 2$ GeV/c

Resolution-corrected away-side (close - far) correlations:




- EP resolution is corrected via unfolding
- BBC Ψ_2 resolution = 0.136 ± 0.002
- Measured slope = 0.08 ± 0.04 (stat.)
- Corrected slope = 0.66 ± 0.27 (stat.)



Event-plane Dependent Two-particle Jet-like Correlations with Event Shape Engineering



Ryo Aoyama, APS-JPS

Two-particle Correlations with ESE



$$C(\Delta\phi) \propto \frac{N_{pair}^{real}(\Delta\phi)}{N_{pair}^{mix}(\Delta\phi)}$$

trigger angle selection w.r.t. EP





- Background subtraction: assuming ZYAM with inclusive-triggered correlations
- v_2 , v_3 and v_4 contributions are subtracted

Polar Representation of Correlation Function

Au+Au $\sqrt{s_{NN}} = 200$ GeV 30-40%


$p_T^t \otimes p_T^a = 4-10 \otimes 1-2$ (GeV/c)

STAR preliminary radius = $1/N^t dN^{ta}/d\Delta\phi$




- Two axes: **trigger axis** and **EP axis**
- $\Delta\phi$ with starts from **red line** and rotate toward counter-clockwise direction
- Radius: the amplitudes of correlated yield

Flow Subtracted Correlation Functions



- **Near-side**

- Out-of-plane trigger: No difference between large- q_2 and small- q_2
- In-plane trigger: peak height difference is enhanced



- **Away-side**

- Out-of-plane trigger: yields are almost fully suppressed both in large- q_2 and small- q_2
- Remnant yield in the EP direction has q_2 dependence
- In-plane trigger: peak height difference is enhanced

Indications of low- p_T particles preferentially escaping towards in-plane direction

Conclusions

- **Event-plane dependent two-particle jet-like correlations shape in 200 GeV Au+Au collisions are reported**
 - Data-driven method to subtract away-side flow background of all harmonics
 - The width of the away-side jet-like peak is found to increase with ϕ_s
 - Consistent with in-medium path length dependence
- **Event-plane dependent two-particle correlations with event shape engineering are measured.**
 - Separation between large- q_2 and small- q_2 events enhances difference of correlation shape while preserving average multiplicity
 - new handle to differentially study partonic energy loss mechanisms
 - Low- p_T particles preferentially escape toward in-plane direction
 - path length dependent? escape mechanism?

Outlook


- Centrality and p_T dependencies
- New data with Event Plane Detector

Thank you !

Back-up slides



v_2 with ESE



- ◆ v_2 is measured via event plane method with TPC-EP with taking 1.0 η gap
- ◆ 20% largest and smallest q_2 vectors are selected with the same region as TPC-EP
- ◆ Top 20% q_2 selection leads to ~10% larger v_2 events
- ◆ Bottom 20% q_2 selection leads to ~8% smaller v_2 events