



Quarkonium measurements in heavy-ion collisions at the STAR experiment

Qian Yang (杨钱)
(for the STAR Collaboration)
Shandong University




Outline

- **Motivation**
- **STAR experiment**
- **J/ ψ measurements in p+Au collisions**
- **Υ measurements in p+Au and Au+Au collisions**
- **Summary**




Quarkonium: sensitive probe to QGP


Heavy quarkonium: heavy mass ($m_c = \sim 1.5 \text{ GeV}/c^2$, $m_b = \sim 4.5 \text{ GeV}/c^2$)
→ early creation.
long lifetime



$T = 0$



$0 < T < T_c$



$T > T_c$



Illustration: A. Rothkopf

Quarkonium is a sensitive probe of the deconfinement in the QGP: color screening dissociation

Cold nuclear matter effects




- **nPDF effect**
 - Modification of gluon distributions in nucleus
 - Shadowing and Anti-shadowing
- **Nuclear absorption effect**
 - Break-up of quarkonium by remnant of incident nuclei.
- **Co-mover effect**
 - Break-up of quarkonium by co-moving hadrons outside of nuclear remnant.





Υ : a cleaner probe at RHIC



Υ is a cleaner probe at RHIC:

- Regeneration is expected to be small

[A. Emerick, X. Zhao and R. Rapp: EPJ A48, 72 (2012)]

[X. Du, M. He, and R. Rapp: PRC 96, 054901 (2017)]

- Co-mover absorption is expected to be small for $\Upsilon(1S)$

[Z. Lin and C. Ko: PLB 503, 104 (2001)]




The Solenoidal Tracker at RHIC

MTD

Dimuon trigger and muon identification
($|\eta| < 0.5$, 45% in $0 < \varphi < 2\pi$)

J/ψ and $\Upsilon \rightarrow \mu^+ + \mu^-$: TPC + MTD

$\Upsilon \rightarrow e^+ + e^-$: TPC + BEMC




TPC

Tracking (momentum measurement, identification) ($|\eta| < 1$, $0 < \varphi < 2\pi$)

BEMC

Trigger and identification of high- p_T electrons
($|\eta| < 1$, $0 < \varphi < 2\pi$)


Inclusive J/ ψ R_{pA} at $\sqrt{s_{NN}} = 200$ GeV



EPS09+NLO, Ma & Vogt, Private Comm.
 nCTEQ, EPS09+NLO, Lansberg Shao,
 Eur.Phys.J. C77 (2017) no.1, 1
 Comp. Phys. Comm. 198 (2016) 238-259
 Comp. Phys. Comm. 184 (2013) 2562-2570

- Model calculations with only nPDF effect can touch upper limit of data within uncertainties.
- Data favor nPDF effects with additional nuclear absorption.

Inclusive Υ R_{pA} at $\sqrt{s_{NN}} = 200$ GeV



PHENIX: PRC 87 (2013) 044909
 STAR: PLB 735 (2014) 127

- Indication of $\Upsilon(1S+2S+3S)$ suppression in p+Au collisions


→ Cold nuclear matter effects:

$$R_{pA} (|y| < 0.5): 0.82 \pm 0.10 \text{ (stat.) } {}_{+0.08}^{-0.07} \text{ (sys.)} \pm 0.10 \text{ (global)}$$

Inclusive Υ R_{AA} vs. N_{part} at $\sqrt{s_{NN}} = 200$ GeV



Combination of dielectron and dimuon channels



$Y(1S)$:

- Stronger suppression towards central collisions



$Y(2S+3S)$:

- Stronger suppression in more central collisions

- More suppressed than $Y(1S)$ in 0-10% central collisions — sequential melting



$\Upsilon(1S)$ suppression





CMS: PLB 770, 357 (2017)

$\Upsilon(1S)$ suppression is similar at RHIC and LHC and no significant p_T dependence:

- Medium temperature is higher at LHC due to higher collision energy
- Regeneration contribution is larger at LHC
- CNM
- Strong suppression of excited Υ states that feed-down to $\Upsilon(1S)$



$\Upsilon(2S+3S)$ suppression



$\Upsilon(2S+3S)$:



- Indication of less suppression at RHIC than at LHC.

STAR: $\Upsilon(2S+3S)$ R_{AA} : 0.35 ± 0.08 (stat.) ± 0.10 (sys.) ($0 < p_T < 10$ GeV/c, 0-60%)

CMS: $\Upsilon(2S)$ R_{AA} : 0.08 ± 0.05 (stat.) ± 0.03 (sys.) ($0 < p_T < 5$ GeV/c, 0-100%)



$\Upsilon(1S)$ suppression



[CMS: PLB 770, 357 (2017)]

[B. Krouppa, A. Rothkopf, M. Strickland: PRD 97, 016017 (2018)]

[X. Du, M. He, and R. Rapp: PRC 96, 054901 (2017)]

$\Upsilon(1S) R_{AA}$:

- Both Rothkopf's and Rapp's models describe data



Rothkopf's model: use a lattice-vetted heavy-quark potential

Rapp's model: use in-medium binding energies predicted by thermodynamic

T-matrix calculations using internal-energy potentials, from lattice QCD



$\Upsilon(2S+3S)$ suppression



[CMS: PLB 770, 357 (2017)]

[B. Krouppa, A. Rothkopf, M. Strickland: PRD 97, 016017 (2018)]

[X. Du, M. He, and R. Rapp: PRC 96, 054901 (2017)]

$\Upsilon(2S+3S)$ R_{AA} :

- Rapp's model describes data
- Rothkopf's model calculation is slightly lower than data in 30-60%

Summary



- **p+Au collisions at $\sqrt{s_{NN}} = 200$ GeV**

- Indication of Υ suppression
- $J/\psi R_{pA}$ favors additional nuclear absorption effect on top of nPDF effect

- **Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV**

The precision of Υ measurements is improved by combining results of dielectron and dimuon channels from dataset taken in different years (2011, 2014 and 2016)

$\Upsilon(1S)$:

- Indication of stronger suppression towards central collisions
- Similar suppression as at LHC
- Both models are consistent with data at RHIC and LHC

$\Upsilon(2S+3S)$:

- More suppressed than $\Upsilon(1S)$ in 0-10% — sequential melting
- Indication of less suppression at RHIC than at the LHC