

The spectra and flow of light nuclei in relativistic heavy ion collisions at RHIC and the LHC


Wenbin Zhao

Peking University

Collaborators: **Huichao Song**, Lilin Zhu, Hua Zheng, **Che Ming Ko**


December 20th 2018

The process of heavy-ion collisions




U. W. Heinz, hep-ph/0407360.

When light nuclei formed in heavy-ion collisions




- Hadrons emitted from the region that reaches limiting temperature and abundances fixed at chemical freeze-out.
- Light nuclei can be formed by baryons after kinetic freeze-out, which is sensitive to nucleon emission source.
- Light nuclei produced at chemical freeze-out might break up and re-form during the chemical freeze-out and the kinetic freeze-out.
- Studying the light nuclei production will help to understand the nucleon emission source and the nucleosynthesis mechanism in heavy-ion collisions.

light nuclei generations in RHIC and LHC



L. Adamczyk et al. [STAR Collaboration], Phys. Rev. C 94, no. 3, 034908 (2016).




S. Acharya et al. [ALICE Collaboration], Eur. Phys. J. C 77, no. 10, 658 (2017).

- Blast wave underpredicts $v_2(p_T)$ of d and ^3He at RHIC, but well describes spectra and $v_2(p_T)$ of d in LHC.
- AMPT + coalescence reproduces the $v_2(p_T)$ of d at Au + Au 200 GeV. Simple coalescence overestimates $v_2(p_T)$ of d at LHC Pb + Pb.

Transport and Coalescence model

- Transport model: with production and annihilation of d in AMPT.
- Coalescence model: AMPT+Coalescence.





- Transport and Coalescence models give similar d spectra.

Y. Oh, Z. W. Lin and C. M. Ko, Phys. Rev. C **80**, 064902 (2009).

Blast Wave + Coalescence

Blast Wave + Coalescence

- Transverse size R_0 is constant (independent on p_T), nucleons are uniform inside a cylinder.





- The Blast Wave + Coalescence can't describe spectra and $v_2(p_T)$ of light nuclei. The constant R_0 is unrealistic.

X. Yin, C. M. Ko, Y. Sun and L. Zhu, Phys. Rev. C **95**, no. 5, 054913 (2017).

Blast Wave + R_0 - p_T correlations + Coalescence

- With additional $R_0 = 10e^{0.23(p_T - 0.9)}$ correlations of hadrons:



- The Blast Wave + R_0 - p_T correlations + Coalescence reproduces spectra and $v_2(p_T)$ of light nuclei. The R_0 - p_T correlations are better.

X. Yin, C. M. Ko, Y. Sun and L. Zhu, Phys. Rev. C **95**, no. 5, 054913 (2017).

iEBE-VISHNU + Coalescence

Coalescence Model

The production probability of nucleus of atomic number A is:

$$\frac{d^3 N_A}{d \mathbf{P}_A^3} = g_A \int \prod_{i=1}^A p_i^\mu d^3 \sigma_{i\mu} \frac{d^3 \mathbf{p}_i}{E_i} f(\mathbf{x}_i, \mathbf{p}_i, t) \\ \times f_A(\mathbf{x}'_1, \dots, \mathbf{x}'_A; \mathbf{p}'_1, \dots, \mathbf{p}'_A; t') \delta^{(3)} \left(\mathbf{P}_A - \sum_{i=1}^A \mathbf{p}_i \right),$$

where g_A is statistical factor, \mathbf{x}_i and \mathbf{p}_i are in fireball frame. \mathbf{x}'_i and \mathbf{p}'_i are Lorentz transformed to the rest frame of produced nucleus. And

$$\int p^\mu d^3 \sigma_\mu \frac{d^3 \mathbf{p}}{E} f_p(\mathbf{x}, \mathbf{p}, t) = N_p, \quad \int p^\mu d^3 \sigma_\mu \frac{d^3 \mathbf{p}}{E} f_n(\mathbf{x}, \mathbf{p}, t) = N_n$$

Wigner function

Wigner function for two-body Coalescence:

$$f_2(\rho, \mathbf{p}_\rho) = 8g_2 \exp \left[-\frac{\rho^2}{\sigma_\rho^2} - \mathbf{p}_\rho^2 \sigma_\rho^2 \right],$$
$$\rho = \frac{1}{\sqrt{2}}(\mathbf{x}'_1 - \mathbf{x}'_2), \quad \mathbf{p}_\rho = \sqrt{2} \frac{m_2 \mathbf{p}'_1 - m_1 \mathbf{p}'_2}{m_1 + m_2}, \quad (1)$$


Wigner function for three-body Coalescence:

$$f_3(\rho, \lambda, \mathbf{p}_\rho, \mathbf{p}_\lambda) = 8^2 g_3 \exp \left[-\frac{\rho^2}{\sigma_\rho^2} - \frac{\lambda^2}{\sigma_\lambda^2} - \mathbf{p}_\rho^2 \sigma_\rho^2 - \mathbf{p}_\lambda^2 \sigma_\lambda^2 \right], \quad (2)$$


$$\begin{aligned} \lambda &= \sqrt{\frac{2}{3}} \left(\frac{m_1 \mathbf{x}'_1 + m_2 \mathbf{x}'_2}{m_1 + m_2} - \mathbf{x}'_3 \right), \\ \mathbf{p}_\lambda &= \sqrt{\frac{3}{2}} \frac{m_3 (\mathbf{p}'_1 + \mathbf{p}'_2) - (m_1 + m_2) \mathbf{p}'_3}{m_1 + m_2 + m_3}, \end{aligned} \quad (3)$$

iEBE-VISHNU

- spectra of π , K and p .




- $v_n\{p_T\}$ for π , K and p




- Hydrodynamics can well describe dynamics evolutions of heavy-ion collisions.

- Event-by-event v_n distributions.




- Event-plane correlations



W. Zhao, H. j. Xu and H. Song, Eur. Phys. J. C **77**, no. 9, 645 (2017).

• Symmetry Cumulant

Pb+Pb 2.76 A TeV




W. Zhao, H. j. Xu and H. Song, Eur. Phys. J. C 77,

no. 9, 645 (2017). Naghmeh Mohammadi, Quark

Matter 2018, (unpublished).

• Non-linear response coefficients

Pb+Pb 2.76 A TeV v.s. 5.02 A TeV



Hydrodynamic model is the standard model to describe the dynamics evolutions of heavy-ion collisions.

Set-up

iEBE-VISHNU + Coalescence

- The coalescence processes for d : $p + n \rightarrow d$.
- Two coalescence processes are included for production of ${}^3\text{He}$:
 $p + p + n \rightarrow {}^3\text{He}$ and $d + p \rightarrow {}^3\text{He}$.

Table 1: Statistical factor (g), radius (R), oscillator frequency (ω), and width parameter ($\sigma_\rho, \sigma_\lambda$) for deuteron and helium-3. Radii are taken from Ref. [?].


Nucleus	g	R (fm)	ω (sec $^{-1}$)	$\sigma_\rho, \sigma_\lambda$ (fm)
deuteron	3/4	2.1421	0.1739	2.473
$p + p + n \rightarrow {}^3\text{He}$	1/4	1.9661	0.5504	1.390
$d + p \rightarrow {}^3\text{He}$	1/3	1.9661	0.3389	1.536

RESULTS


Light nuclei production in Au + Au 200 GeV

- iEBE-VISHNU + Coalescence

Au+Au 200 GeV, 0-80%




Au+Au 200 GeV, 0-80%



- iEBE-VISHNU + Coalescence gives a good predictions for the spectra and $v_2(p_T)$ of d and ${}^3\text{He}$ in Au+Au 200 GeV.
- iEBE-VISHNU gives a good description of space momentum correlations of hadrons naturally.

W. Zhao, L. Zhu, H. Zheng, C. M. Ko and H. Song, Phys. Rev. C **98**, no. 5, 054905 (2018)


Spectra of light nuclei in Pb + Pb 2.76 and 5.02 TeV



- iEBE-VISHNU + Coalescence gives a good description for the spectra of d and underestimate spectra of ${}^3\text{He}$ in Pb + Pb 2.76 TeV.
- The spectra of d and ${}^3\text{He}$ in Pb + Pb 5.02 TeV is higher and flatter.

W. Zhao, L. Zhu, H. Zheng, C. M. Ko and H. Song, Phys. Rev. C **98**, no. 5, 054905 (2018)

$v_2(p_T)$ of light nuclei in Pb + Pb 2.76 and 5.02 TeV




- iEBE-VISHNU + Coalescence well reproduce the $v_2(p_T)$ of d in Pb + Pb 2.76 TeV.
- iEBE-VISHNU + Coalescence predicts the $v_2(p_T)$ of d and ${}^3\text{He}$ in Pb + Pb 5.02 TeV. And the $v_2(p_T)$ of d is slightly lower.

W. Zhao, L. Zhu, H. Zheng, C. M. Ko and H. Song, Phys. Rev. C **98**, no. 5, 054905 (2018)

- iEBE-VISHNU + Coalescence do a great job in describing the spectra and elliptic flow of light nuclei at Au + Au 200 GeV and Pb + Pb 2.76 TeV.
- iEBE-VISHNU generates the proper space-momeutum correlations of hadrons at kinetic freezeout.
- Coalescence together with proper space-momeutum correlations of hadrons might the mechanism of light nuclei productions in heavy-ion collisions.
- ^3He is largely underestimated in Pb + Pb 2.76 TeV.

The productions of ${}^3\text{He}$



- Including the $d + p \rightarrow {}^3\text{He}$ enhances the spectra of ${}^3\text{He}$ by two.
- $p + p + n \rightarrow {}^3\text{He} + d + p \rightarrow {}^3\text{He}$ can describe spectra in Au + Au 200 GeV, but underestimate that in Pb + Pb 2.76 TeV.
- Production of ${}^3\text{He}$ need be further studied, such as including 3-body interactions in transport model and the Pauli principle effects in Wigner function of ${}^3\text{He}$.

W. Zhao, L. Zhu, H. Zheng, C. M. Ko and H. Song, Phys. Rev. C **98**, no. 5, 054905 (2018)

Summary

- Studying light nuclei helps to understand the nucleon emission source and nucleosynthesis mechanism in heavy-ion collisions.
- iEBE-VISHNU + Coalescence can well describe the spectra and elliptic flow of light nuclei at Au + Au 200 GeV and Pb + Pb 2.76 TeV. iEBE-VISHNU generates the proper space-momentum correlations of hadrons at kinetic freezeout.
- Coalescence might be the mechanism of light nuclei production in heavy-ion collisions.
- The production of ^3He requires further study.

Thanks