

Search for heavy Higgs at the LHC

Xin Chen, **Yue Xu**, Hang Chen

Tsinghua University

23rd Mini-workshop on the frontier of LHC
Chongqing, May 18, 2019

<https://arxiv.org/abs/1905.05421>





- 1 Introduction
- 2 Interaction and Process
- 3 Heavy Higgs with Boosted Boson jet
- 4 95% Confidence Level Exclusion
- 5 Conclusion


Introduction

Many BSM predict heavy Higgs particles decaying to heavy quarks or bosons

- 2HDM, MSSM, and so on...

ATLAS and CMS have some researches on heavy Higgs



- gluon-gluon fusion (ggF)
- vector-boson fusion (VBF)
- associated production with vector boson (VH)



Introduction



• $H \rightarrow ZZ \rightarrow 4\ell$ research in ATLAS

CERN-EP-2017-251



• $X \rightarrow WW$ research in CMS

CMS-PAS-HIG-17-033



Dim-4 and Dim-6 Interaction

- In our paper: one SM-like light Higgs(h) and one generic heavy Higgs(H)
- A generic heavy Higgs: both dim-4 and effective dim-6 interactions with SM particles
- dim-4 operator Lagrangian:

$$\mathcal{L}_{hWW}^{(4)} = \rho_h g m_W h W^\mu W_\mu$$

$$\mathcal{L}_{hZZ}^{(4)} = \rho_h \frac{g m_W}{2 \cos^2 \theta_W} h Z^\mu Z_\mu$$

$$\mathcal{L}_{HWW}^{(4)} = \rho_H g m_W H W^\mu W_\mu$$

$$\mathcal{L}_{HZZ}^{(4)} = \rho_H \frac{g m_W}{2 \cos^2 \theta_W} H Z^\mu Z_\mu$$

$$\rho_h = \frac{g_h^2 \nu_h}{g^2 \nu}, \quad \rho_H = \frac{g_H^2 \nu_H}{g^2 \nu}$$

ν_h and ν_H : VEVs , g_h and g_H : gauge couplings

ν and g : SM VEV and coupling , θ_W : weak mixing angle



Dim-4 and Dim-6 Interaction

- dim-6 operator effective Lagrangian:

$$\mathcal{L}_{HVV}^{(6)} = \sum_n \frac{f_n}{\Lambda^2} \mathcal{O}_n, \quad \Lambda = 5 \text{ TeV}$$

$$\mathcal{L}_{HWW}^{(6)} = \rho_H g m_W \frac{f_W}{2\Lambda^2} (W_{\mu\nu}^+ W^{-\mu} \partial^\nu H + h.c.)$$

$$- \rho_H g m_W \frac{f_{WW}}{\Lambda^2} W_{\mu\nu}^+ W^{-\mu\nu} H$$

$$\mathcal{L}_{HZZ}^{(6)} = \rho_H g m_W \frac{c^2 f_W + s^2 f_B}{2c^2 \Lambda^2} Z_{\mu\nu} Z^\mu \partial^\nu H$$

$$- \rho_H h m_W \frac{c^4 f_{WW} + s^4 f_{BB}}{2c^2 \Lambda^2} Z_{\mu\nu} Z^{\mu\nu} H$$

$$s = \sin\theta_W, \quad c = \cos\theta_W$$



Production Channels

- Yukawa coupling between heavy Higgs and fermions is small
- The associated VH(V=W/Z) production channel is considered(VBF production channel is accompanied by large bkg)
- heavy Higgs decay modes:

$$H \rightarrow WW \rightarrow l\nu jj$$

$$H \rightarrow WW \rightarrow jjjj$$

$$H \rightarrow ZZ \rightarrow lljj$$

$$l = e/\mu$$

- final states with 2ℓ and 3ℓ from three bosons' decays are used



Background Estimation

2 ℓ channel:

- single boson($\ell\ell$) plus four QCD partons
- diboson ($\ell\ell + jj$) plus two QCD partons
- triboson process $\ell\ell + 4j$ (VBF included)
- $t\bar{t}$ with $t \rightarrow \ell\nu b$
- $t\bar{t}Z$ with $Z \rightarrow \ell\ell$


3 ℓ channel:

- diboson($3\ell + \nu$) plus two QCD partons
- triboson process $3\ell + \nu + jj$ (VBF included)
- $t\bar{t}V$ with 3ℓ in the final state



Boosted Boson jet

- $V \rightarrow jet + jet$ process is accompanied by large SM bkg
- boosted boson jet: high p_T , variable τ_2 and τ_1 describing jet substructure



by Toshi SUMIDA




600GeV Heavy Higgs Mass: 2lep final state

Region 1:

$V_0 H \rightarrow l^- l^+ + V_1(\text{subjet of leading fatjet}) V_2(\text{subjet of leading fatjet})$
cut:

- $p_{T,V_0} > 950 \text{ GeV}$
- $80 \text{ GeV} < m_{V_0} < 100 \text{ GeV}$
- $\text{fatjet_sj_n of leading fatjet} = 2$
- $70 \text{ GeV} < m_{V_1}, m_{V_2} < 150 \text{ GeV}$
- $\frac{\tau_2}{\tau_1}$ of leading fatjet < 0.45
- $p_{T,HH} > 750 \text{ GeV}$




600GeV Heavy Higgs Mass: 2lep final state

Region 2:

$$V_0 H \rightarrow l^- l^+ + V_1(jet) V_2(jet + jet)$$

cut:

- $p_{T,V_0} > 550 \text{ GeV}$,
 $p_{T,V_1} > 300 \text{ GeV}$, $p_{T,V_2} > 150 \text{ GeV}$
- $80 \text{ GeV} < m_{V_0} < 100 \text{ GeV}$,
 $70 \text{ GeV} < m_{V_1} < 150 \text{ GeV}$,
 $70 \text{ GeV} < m_{V_2} < 110 \text{ GeV}$
- $\frac{\tau_2}{\tau_1}$ of $V_1 < 0.4$
- $p_{T,HH} > 550 \text{ GeV}$




600GeV Heavy Higgs Mass: 2lep final state

Region 3:

$$V_0 H \rightarrow jet + V_1(l^+ l^-) V_2(jet + jet)$$

cut:

- $p_{T,V_0} > 700 GeV$,
 $p_{T,V_1} > 300 GeV$, $p_{T,V_2} > 50 GeV$
- $70 GeV < m_{V_0} < 150 GeV$,
 $80 GeV < m_{V_1} < 100 GeV$,
 $75 GeV < m_{V_2} < 115 GeV$
- $\frac{\tau_2}{\tau_1}$ of $V_0 < 0.6$
- $p_{T,HH} > 700 GeV$




600GeV Heavy Higgs Mass: 2lep final state

Region 4:

$$V_0 H \rightarrow jet + V_1(l^+ l^-) V_2(jet)$$

cut:

- $p_{T,V_0} > 700 GeV$,
 $p_{T,V_1} > 300 GeV$, $p_{T,V_2} > 250 GeV$
- $70 GeV < m_{V_0} < 150 GeV$,
 $80 GeV < m_{V_1} < 100 GeV$,
 $75 GeV < m_{V_2} < 150 GeV$
- $\frac{\tau_2}{\tau_1}$ of $V_0 < 0.52$, $\frac{\tau_2}{\tau_1}$ of $V_2 < 0.52$
- $p_{T,HH} > 700 GeV$




600GeV Heavy Higgs Mass: 3lep final state

Region 1:

$$V_0 H \rightarrow l^\pm \nu_{l^\pm} + V_1(l^+ l^-) V_2(jet)$$

cut:

- $p_{T,V_0} > 600 GeV, p_{T,\nu_{l^\pm}} > 50 GeV$
- $80 GeV < m_{V_1} < 100 GeV,$
 $60 GeV < m_{V_2} < 160 GeV$
- $\frac{\tau_2}{\tau_1}$ of $V_2 < 0.6$
- $p_{T,HH} > 600 GeV$




600GeV Heavy Higgs Mass: 3lep final state

Region 2:

$$V_0 H \rightarrow l^\pm \nu_{l^\pm} + V_1(l^+ l^-) V_2(\text{fatjet})$$

cut:

- $p_{T,V_0} > 600\text{GeV}$, $p_{T,\nu_{l^\pm}} > 50\text{GeV}$
- $80\text{GeV} < m_{V_1} < 100\text{GeV}$,
 $70\text{GeV} < m_{V_2} < 140\text{GeV}$
- $\frac{\tau_2}{\tau_1}$ of $V_2 < 0.5$
- $p_{T,HH} > 600\text{GeV}$




600GeV Heavy Higgs Mass: 3lep final state

Region 3:

$$V_0 H \rightarrow l^\pm \nu_{l^\pm} + V_1(l^+ l^-) V_2(jet + jet)$$

cut:

- $p_{T,V_0} > 600 GeV, p_{T,\nu_{l^\pm}} > 50 GeV$
- $80 GeV < m_{V_1} < 100 GeV,$
 $60 GeV < m_{V_2} < 120 GeV$
- $p_{T,HH} > 600 GeV$




600GeV Heavy Higgs Mass: 3lep final state

Region 4:

$$V_0 H \rightarrow l^+ l^- + V_1 (l^\pm \nu_{l^\pm}) V_2 (jet)$$

cut:

- $p_{T,V_0} > 600 GeV, p_{T,\nu_{l^\pm}} > 50 GeV$
- $80 GeV < m_{V_0} < 100 GeV,$
 $60 GeV < m_{V_2} < 160 GeV$
- $\frac{\tau_2}{\tau_1}$ of $V_2 < 0.6$
- $p_{T,HH} > 600 GeV$




600GeV Heavy Higgs Mass: 3lep final state

Region 5:

$$V_0 H \rightarrow l^+ l^- + V_1(l^\pm \nu_{l^\pm}) V_2(\text{fatjet})$$

cut:

- $p_{T,V_0} > 600\text{GeV}$, $p_{T,\nu_{l^\pm}} > 50\text{GeV}$
- $80\text{GeV} < m_{V_0} < 100\text{GeV}$,
 $70\text{GeV} < m_{V_2} < 140\text{GeV}$
- $\frac{\tau_2}{\tau_1}$ of $V_2 < 0.5$
- $p_{T,HH} > 600\text{GeV}$




600GeV Heavy Higgs Mass: 3lep final state

Region 6:

$$V_0 H \rightarrow l^+ l^- + V_1 (l^\pm \nu_{l^\pm}) V_2 (jet + jet)$$

cut:


- $p_{T,V_0} > 600 GeV, p_{T,\nu_{l^\pm}} > 50 GeV$
- $80 GeV < m_{V_0} < 100 GeV,$
 $60 GeV < m_{V_2} < 120 GeV$
- $p_{T,HH} > 600 GeV$




600GeV Heavy Higgs Mass

- $\int L dt = 300 fb^{-1}$
- $\rho_H = 0.05$, $f_W = 1000$, $f_{WW} = 1000$

2lep



3lep



300GeV Heavy Higgs Mass: 2lep final state

Region	Process
1	$V_0 H \rightarrow l^- l^+ + V_1(\text{subjet of L-fatjet}) V_2(\text{subjet of L-fatjet})$
2	$V_0 H \rightarrow l^- l^+ + V_1(\text{jet}) V_2(\text{jet} + \text{jet})$
3	$V_0 H \rightarrow \text{jet} + V_1(l^+ l^-) V_2(\text{jet} + \text{jet})$


- take out process: $V_0 H \rightarrow \text{jet} + V_1(l^+ l^-) V_2(\text{jet})$
- a bit tighter mass window cut



900GeV Heavy Higgs Mass: 2lep final state

Region	Process
1	$V_0 H \rightarrow l^- l^+ + V_1(\text{jet}) V_2(\text{jet} + \text{jet})$
2	$V_0 H \rightarrow \text{jet} + V_1(l^- l^+) V_2(\text{jet})$
3	$V_0 H \rightarrow l^- l^+ + V_1(\text{jet}) V_2(\text{jet})$

- take out process: $V_0 H \rightarrow l^- l^+ + V_1(\text{subset of leading fatjet}) V_2(\text{subset of leading fatjet})$
- take out process: $V_0 H \rightarrow \text{jet} + V_1(l^+ l^-) V_2(\text{jet} + \text{jet})$
- add process: $V_0 H \rightarrow l^- l^+ + V_1(\text{jet}) V_2(\text{jet})$



95% Confidence Level Exclusion


- the mass of gauge boson is equal to SM mass

$$\rho_h \frac{\nu_h}{\nu} + \rho_H \frac{\nu_H}{\nu} + \dots = 1$$

- take $\rho_h = 1$ and $\rho_H = 0.05$ as benchmark value
- get CL of each parameter point by "template" fits with a large quantity of toy experiments



95% Confidence Level Exclusion



Conclusion

- Two and three lepton final states are considered
- SM background can be suppressed by applying of boosted boson jets
- We focus on dim-6 operator effect and the dim-4 effect is small
- A big part of parameter space which is among unitarity bound can be excluded
- This part of phase space is waiting for discovery

