



Contribution ID: 171

Type: **Parallel**

Possible interpretation of N(1685)

Wednesday, 21 August 2019 09:55 (20 minutes)

$N(1685)$ was firstly reported in the photoproduction of η meson off the quasi-free neutron. Unlike other low-lying nucleon resonances whose Breit-Wigner (BW) widths are at least 100 MeV, the BW width of the $N(1685)$ is less than 30 MeV. The interpretation of the $N(1685)$ is still an open question.

The non-strange pentaquark mass of all quark configurations is evaluated in a constituent quark model, where the Cornell potential is employed and all model parameters are predetermined by comparing the theoretical and experimental masses of low-lying baryons which are believed to be mainly $3q$ states. The state with the $[31]_{FS}[22]_F[31]_S(q^4\bar{q})$ configuration and quantum numbers $I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$ is predicted to be the lowest pentaquark state. Its mass is derived about 1670 MeV, which encourages one to expect that the $N(1685)$ could be the lowest pentaquark state.

Primary author: XU, Kai (Suranaree University of Technology)

Co-authors: KAEWSNOD, Attaphon (S); Mr SRISUPHAPHON, Sorakrai (Department of Physics, Faculty of Science, Burapha University, Chonburi 20131, Thailand); YAN, Yupeng (Suranaree University of Technology)

Presenter: XU, Kai (Suranaree University of Technology)

Session Classification: Session 2: Baryon spectroscopy

Track Classification: Session 2: Baryon spectroscopy