



Contribution ID: 6

Type: **Parallel**

## Hadronic charm decays at BESIII

*Saturday, 17 August 2019 14:55 (20 minutes)*

BESIII has collected data samples corresponding to luminosities of  $2.93 \text{ fb}^{-1}$ ,  $3.19 \text{ fb}^{-1}$  and  $0.567 \text{ fb}^{-1}$  at center-of-mass energies of 3.773, 4.178, and 4.6 GeV, respectively. The data set collected at 3.773 GeV contains quantum-correlated  $D^0\bar{D}^0$  pairs that allow to access the phase differences between amplitudes. We report the measurements of strong phase differences in  $D^0(-\bar{D}^0)$  decays, especially for  $K_S^0/L\pi^+\pi^-$ , which can reduce the  $\gamma/\phi^3$  measurement uncertainty at LHCb and Belle II. In addition, we report the measurements of the absolute branching fraction and amplitude analysis of  $D^+$ ,  $D^0$ ,  $D_s^+$  and  $\Lambda_{c^+}$

**Primary author:** Dr LI, Pei-Rong (Lanzhou University)

**Presenter:** Dr LI, Pei-Rong (Lanzhou University)

**Session Classification:** Session 4: Hadron decays, production and interactions

**Track Classification:** Session 4: Hadron decays, production and interactions