



Contribution ID: 188

Type: Parallel

New spectrum of negative-parity doubly charmed baryons: Possibility of two quasistable states

Tuesday, August 20, 2019 2:25 PM (20 minutes)

The discovery of Ξ_{cc}^{++} by the LHCb Collaboration triggers predictions of more doubly charmed baryons. By taking into account both the P -wave excitations between the two charm quarks and the scattering of light pseudoscalar mesons off the ground state doubly charmed baryons, a set of negative-parity spin-1/2 doubly charmed baryons are predicted already from a unitarized version of leading order chiral perturbation theory. Moreover, employing heavy antiquark-diquark symmetry the relevant low-energy constants in the next-to-leading order are connected with those describing light pseudoscalar mesons scattering off charmed mesons, which have been well determined from lattice calculations and experimental data. Our calculations result in a spectrum richer than that of heavy mesons. We find two very narrow $J^P = 1/2^- \Omega_{cc}^P$, which very likely decay into $\Omega_{cc}\pi^0$ breaking isospin symmetry. In the isospin-1/2 Ξ_{cc}^P sector, three states are predicted to exist below 4.2 GeV with the lowest one being narrow and the other two rather broad. We suggest to search for the Ξ_{cc}^P states in the $\Xi_{cc}^{++}\pi^-$ mode. Searching for them and their analogues are helpful to establish the hadron spectrum.

Primary authors: Dr YAN, Maojun (ITP, CAS); Dr LIU, Xiao-Hai (Center for Joint Quantum Studies, Department of Physics, School of Science, Tianjin University)

Co-authors: Prof. ZOU BINGSONG, Bingsong (IHEP, CAS); Dr HANHART, Christoph (IAS, Forschungszentrum Juelich, Germany); Dr GUO, Feng-Kun (ITP, CAS); Prof. MEISSNER, Ulf-G. (Universitaet Bonn and Forschungszentrum Juelich)

Presenter: Dr YAN, Maojun (ITP, CAS)

Session Classification: Session 2: Baryon spectroscopy

Track Classification: Session 2: Baryon spectroscopy