XVIII International Conference on Hadron Spectroscopy and Structure (HADRON2019)



Contribution ID: 16

Type: Parallel

Tetraquark mixing framework to explain the two light-meson nonets

Wednesday, 21 August 2019 11:30 (20 minutes)

We propose a tetraquark mixing framework for the two light-meson nonets in the $J^P = 0^+$ channel, the light nonet $f_0(500)$, $f_0(980)$, $a_0(980)$, $K_0^*(800)$ and the heavy nonet $f_0(1370)$, $f_0(1500)$, $a_0(1450)$, $K_0^*(1430)$. According to this framework, one can introduce two types of tetraquark with different spin configuration, $|J, J_{12}, J_{34}| = |000| |011|$, where J is the spin of the tetraquark, J_{12} the diquark spin, J_{34} the antidiquark spin. They differ by the color configuration also but both have the same flavor structure. The two tetraquark types seem to have interesting correspondence with the two nonets in PDG. Indeed, the two tetraquarks mix strongly through the hyperfine color-spin interaction and the eigenstates that diagonalize the hyperfine masses can be identified with the two nonets in PDG. We report that their hyperfine mass splitting can generate the mass gap between the two nonets qualitatively. We also discuss interesting signatures in the decays of this tetraquark model.

References:

EPJC(2017)77:173 Hungchong Kim, Myung-Ki Cheoun, K.S.Kim

EPJC(2017)77:435 K.S.Kim, Hungchong Kim

PRD97,094005(2018) Hungchong KIm,K.S.Kim,Myung-Ki Cheoun, Makoto Oka

PRD99,014005(2019) Hungchong KIm,K.S.Kim,Myung-Ki Cheoun, Daisuke Jido, Makoto Oka

Primary author: Dr KIM, Hungchong (Korea Aerospace University)

Presenter: Dr KIM, Hungchong (Korea Aerospace University)

Session Classification: Session 1: Meson spectroscopy

Track Classification: Session 1: Meson spectroscopy