

XVIII International Conference on Hadron Spectroscopy and Structure
(HADRON2019)



Contribution ID: 71

Type: **Parallel**

On decays of $X(3872)$ to $\chi_{cJ}\pi^0$ and $J/\psi\pi^+\pi^-$

Saturday, 17 August 2019 15:55 (20 minutes)

By describing the $X(3872)$ using the extended Friedrichs scheme, in which $D\bar{D}^*$ is the dominant component, we calculate the decay rates of the $X(3872)$ to π^0 and a P -wave charmonium χ_{cJ} state with $J = 0, 1$, or 2 , and its decays to $J/\psi\pi^+\pi^-$ where $\pi^+\pi^-$ are assumed to be produced via an intermediate ρ state. The decay widths of $X(3872) \rightarrow \chi_{cJ}\pi^0$ for $J = 0, 1, 2$ are of the same order. However, this model calculation exhibits that the decay rate of $X(3872)$ to $\chi_{c1}\pi^0$ is one order of magnitude smaller than its decay rate to $J/\psi\pi^+\pi^-$.

Primary author: Dr ZHOU, Zhi-Yong (Southeast University, China)

Co-author: Dr XIAO, Zhiguang (USTC)

Presenter: Dr ZHOU, Zhi-Yong (Southeast University, China)

Session Classification: Session 3: Exotic hadrons and candidates

Track Classification: Session 3: Exotic hadrons and candidates