




*I. Vassiliev for the CBM Collaboration*



- Physics case
- MSH reconstructions
- BES-I MSH measurements, models and predictions
- Tests with experimental data
- Summary



# Physics case: Exploring the QCD phase diagram




**The equation-of-state at high  $\rho_B$**   
 collective flow of hadrons,  
 particle production at threshold energies:  
**multi-strange hyperons, hypernuclei**

**Deconfinement phase transition at high  $\rho_B$**   
 excitation function and flow of  
 strangeness ( $K, \Lambda, \Sigma, \Xi, \Omega$  and  $\varphi$ )


**Chiral symmetry restoration at high  $\rho_B$**   
 in-medium modifications of hadrons ( $\rho$ )  
 excitation function of **multi-strange (anti)hyperons**

**QCD critical endpoint**  
 excitation function of event-by-event fluctuations  
 ( $\pi, K, p, \Lambda, \Xi, \Omega \dots$ )

**Projects to explore the QCD phase diagram at large  $\mu_B$ :**  
 RHIC (STAR) beam energy-scan, HADES, NA61@SPS,  
 MPD@NICA: bulk observables  
**CBM: bulk and rare observables, high statistic!**




# Experiments exploring dense QCD matter



**CBM:**  
*unprecedented*  
**(high) rate**  
**capability**

- determination of (displaced) vertices with high resolution ( $\approx 50 \mu\text{m}$ )
- identification of leptons and hadrons
- fast and radiation hard detectors
- self-triggered readout electronics
- high speed data acquisition and
- online event selection
- powerful computing farm *and 4D tracking*
- software triggers

# Strangeness world data



No data available at FAIR energy


In the AGS (SIS100) energy range, only about 300  $\Xi^-$ -hyperons have been measured in Au+Au collisions at 6A GeV

High-precision measurements of excitation functions of multi-strange hyperons in A+A collision with different mass numbers A at SIS100 energies have a discovery potential to find a signal for the onset of deconfinement in QCD matter at high net-baryon densities.

*What about models?*



# Strangeness data BES-I




Helen Caines  
EMMI Workshop  
GSI February 2019

Non-trivial  $v_s$  dependence  
of strange baryons

Strangeness results submitted  
for publication soon

$\Omega^\pm$  (!)







# BES-I MSH measurements and models

\* STAR BES data by  
Helen Caines  
EMMI Workshop  
GSI February 2019

- Thermal model by  
A. Andronic  
32 CBM week


- Very reasonable agreement by both models
- More strangeness needed (PHSD)




# QGP and Chiral symmetry restoration

“Chiral symmetry restoration versus deconfinement in heavy-ion collisions at high baryon density”

W. Cassing, A. Palmese, P. Moreau, and E. L. Bratkovskaya Phys.Rev. C93 (2016), 014902, arXiv:1510.04120 [nucl-th]



Multi-step production of multi-strange baryons  $\Xi$  and  $\Omega$





Chiral symmetry restoration (CSR) change the flavor decomposition – more s-sbar pairs produced.

Droplets of QGP allow to interact s-sbar quarks and create more multi-strange (anti)baryons.

- Presence of QGP significantly increase yield of  $\Omega^+$  at FAIR energy
- CSR effect increase yield of  $\Omega^-$  and  $\Omega^+$  at FAIR energy



# Performance of the CBM track finder



AuAu 10 AGeV/c

165  $\pi$

170 p



26 K

15  $\Lambda$

20

$K_S^0$

0.3  $\Xi^-$




- For studies several theoretical models like UrQMD and PHSD are used.
- Track finder is based on the Cellular Automaton method.
- High efficiency for track reconstruction of more than **92%**, including fast (more than 90%) and slow (more than 65%) secondary tracks.
- Time-based track finder is developed, efficiency is stable with respect to the interaction rate.
- Low level of split and wrongly reconstructed (ghost) tracks.

**minimum bias : 6ms/core track finder, 1 ms/core particle finder**



# High rate scenario: MSH reconstruction with 4D tracking



# 4D Track Finder in CBMROOT


100 AuAu 10 AGeV mbias events

Efficiency, %	3D	0.1 MHz	1 MHz	10 MHz
All tracks	92.5 %	93.8 %	93.5 %	91.7 %
Primary high-p	98.3 %	98.1 %	97.9 %	96.2 %
Primary low-p	93.9 %	95.4 %	95.5 %	94.3 %
Secondary high-p	90.8 %	94.6 %	93.5 %	90.2 %
Secondary low-p	62.2 %	68.5 %	67.6 %	64.3 %
Clone level	0.6 %	0.6 %	0.6 %	0.6 %
Ghost level	1.8 %	0.6 %	0.6 %	0.6 %
True hits per track	92%	93 %	93 %	93%
Hits per MC track	7.0	7.0	6.97	6.70


Timeslices from CBMROOT  
Timebased digitisation, cluster and hit finder

# Particle identification with PID detectors


**ToF: hadron identification**



**RICH: electron identification**




**TRD: d-He separation**




**PID detectors:**

- ToF (Time of Flight) — hadron identification;
- RICH (Ring Imaging Cherenkov detector) — electron identification;
- TRD (Transition Radiation detector) — electron and heavy fragments identification.

PID detectors of CBM will allow a clear identification of charged tracks.











# KF Particle Finder for the CBM Experiment



- More than 150 decays. All decays are reconstructed in one go.
- Based on the Kalman filter method - mathematically correct parameters and their errors.
- KF Particle Finder is successfully tested in STAR and allows to reconstruct up to 2 times more signal.
- STAR developments are fully merged with the KF Particle Finder repository.



## 5M central AuAu collisions 10A GeV/c










- CBM will allow clean reconstruction of rare strange probes with high efficiency and high statistics.
- Tools for the multi-differential physics analysis are prepared.

# Multi strange hyperons reconstruction with missing mass method








5M central AuAu collisions 10A GeV/c




- Comparable efficiencies, better control over the systematic errors
- $\Sigma^+$  and  $\Sigma^-$  physics: completes the picture of strangeness production



# CBM KFParticle Finder @ STAR 4.4M Au+Au events sqrt(s) = 7.7




- CBM KF Particle Finder is successfully applied to the STAR data in a wide energy range.
- STAR data are excellent platform to test and improve our reconstruction software.








# Testing CBM algorithms online: STAR express analysis


Maksym Zyzak, 33<sup>rd</sup> CBM Collaboration Meeting, Darmstadt








- 600K events of the 2019 data with TFG19c xCalibration are processed.
- ToF was at the stage of calibration and is not used in the current analysis.
- Obtained mass peaks of short-lived particles have parameters similar to the offline reconstruction.
- With TFG19d xCalibration we expect to obtain more clean samples.




A. Andronic et al., Phys. Lett. B697 (2011) 203


5M mbias events Au+Au at 10AGeV/c  
50 sec at 0.1MHz IR (1.8 k/sec)



Expected collection rate:  $\sim 60$   ${}^6_{\Lambda\Lambda}$  He  
in 1 week at **10MHz IR** (not day-1)



- According to the current theoretical predictions CBM will be able to perform comprehensive study of hypernuclei, including:
  - precise measurements of lifetime;
  - excitation functions;
  - flow.
- It has a huge potential to register and investigate double  $\Lambda$  hypernuclei.







# DCM with CBM detector 5M mbias C + C collisions

## About 50 sec of data taking assuming $10^5$ IR




A.S.Botvina, K.K.Gudima, J.Pochodzalla.

Production of hypernuclei in peripheral relativistic ion collisions.

Phys. Rev. C , v. 88, p. 054605, 2013.




$\Lambda N, \Lambda NN, {}^4_\Lambda\text{H}$  and  ${}^4_\Lambda\text{He}$



# Day-1: Expected particle yields

## Au+Au @ 6, 10 AGeV



Particle (mass MeV/c <sup>2</sup> )	Multiplicity central ev. 6 AGeV	Multiplicity central ev. 10 AGeV	decay mode	BR	$\epsilon$ (%)	yield in 90 days 6AGeV	yield in 90 days 10 AGeV	IR MHz
$\bar{\Lambda}$ (1115)	$4.6 \cdot 10^{-4}$	0.034	$\bar{p}\pi^+$	0.64	19.7	$1.1 \cdot 10^7$	$8.3 \cdot 10^8$	<b>0.1</b>
$\Xi^-$ (1321)	0.054	0.222	$\Lambda\pi^-$	1	9.9	$1.0 \cdot 10^9$	$4.3 \cdot 10^9$	<b>0.1</b>
$\Xi^+$ (1321)	$3.0 \cdot 10^{-5}$	$5.4 \cdot 10^{-4}$	$\bar{\Lambda}\pi^+$	1	8.7	$5.0 \cdot 10^5$	$9.1 \cdot 10^6$	<b>0.1</b>
$\Omega^-$ (1672)	$5.8 \cdot 10^{-4}$	$5.6 \cdot 10^{-3}$	$\Lambda K^-$	0.68	4.4	$3.4 \cdot 10^6$	$3.3 \cdot 10^7$	<b>0.1</b>
$\Omega^+$ (1672)	-	$7 \cdot 10^{-5}$	$\bar{\Lambda}K^+$	0.68	3.9	0 (QGP?)	$3.8 \cdot 10^5$	<b>0.1</b>
${}^3_{\Lambda}\text{H}$ (2993)	$4.2 \cdot 10^{-2}$	$3.8 \cdot 10^{-2}$	${}^3\text{He}\pi^-$	0.25	12.7	$2.7 \cdot 10^8$	$2.5 \cdot 10^8$	<b>0.1</b>
${}^4_{\Lambda}\text{He}$ (3930)	$2.4 \cdot 10^{-3}$	$1.9 \cdot 10^{-3}$	${}^3\text{He}p\pi^-$	0.32	11.4	$1.7 \cdot 10^7$	$1.4 \cdot 10^7$	<b>0.1</b>
${}^5_{\Lambda\Lambda}\text{He}$ (5047)		$5.0 \cdot 10^{-6}$	${}^3\text{He}2p2\pi$	0.01	3	15	250	<b>0.1</b>
${}^6_{\Lambda\Lambda}\text{He}$ (5986)		$1.0 \cdot 10^{-7}$	${}^4\text{He}2p2\pi$	0.01	1.2			<b>0.1</b>




# Summary


- CBM detector is an excellent device to measure not only bulk observables, but strangeness, hypernuclei and other rare probes with high statistic.
- The CBM experiment will provide multidifferential high precision measurements of strange hadrons including multi-strange (anti)-hyperons.
- High precision measurements of excitation functions of multi-strange hyperons in A+A collision with different mass numbers A at SIS100 energies have a discovery potential to find a signal for the onset of deconfinement in QCD matter at high net-baryon densities
- The discovery of (double-)  $\Lambda$  hypernuclei and the determination of their lifetimes will provide information on the hyperon-nucleon and hyperon-hyperon interactions, which are essential ingredients for the understanding of the nuclear matter EoS at high densities, and, hence, of the structure of neutron stars.



# CBM Collaboration: 64 Institutes, ~600 members















# Strange particle reconstruction performance



5M central AuAu collisions 10A GeV/c ( $\bar{E}^-$ )



- CBM will allow clean reconstruction of rare strange probes with high efficiency and high statistics.
- Tools for the multi-differential physics analysis are prepared.