



Recent results about B physics at ATLAS

Weimin Song (宋维民)

Jilin University

weimin.song@stfc.ac.uk

Outline

Introduction of heavy flavour physics program at ATLAS


Four topics:

1. Search for the X(5568) in $B_s^0 \pi^\pm$ final states (PRL 120, 202007 (2018))
2. Observation of an Excited B_c^\pm Meson State (PRL 113, 212004 (2014))
3. ϕ s and $\Delta\Gamma$ s measurement in the $B_s^0 \rightarrow J/\Psi \phi$ channel (JHEP 08 (2016) 147, ATLAS-CONF-2019-009)
4. B_s^0 (and B^0) $\rightarrow \mu\mu$ measurement (JHEP 04 (2019) 098)


Heavy Flavour physics program at ATLAS

- ❖ Precision measurement to find hint of derivation from SM: rare decays, such $B_s^0 \rightarrow \mu^+ \mu^-$ branching fraction measurement.....
- ❖ Production and decay of heavy flavour hadrons to understand the strong interaction, such as the discovery of $B_c(2S)$
- ❖ Usually, two muons with a common vertex with invariant mass near J/Ψ are required: the inner tracker and muon detector are used

7/10/2019



Search for the $X(5568)$ in $B_s^0\pi^\pm$ final states





D0 Collaboration reported evidence of the $X(5568) \rightarrow B_s^0 \pi^\pm$, $B_s^0 \rightarrow J/\psi \phi$, and reported consistent result in the semi-leptonic decay of B_s^0 :

Mass ~ 5568 MeV; Width ~ 20 MeV
Good candidate for tetraquark state

LHCb, CMS at LHC and CDF at Tevatron revealed no signal with similar techniques.


PRL 117, 022003 (2016)

PHYSICAL




Search for the $X(5568)$ in $B_s^0 \pi^\pm$ final states

- ❖ Di-muon trigger is used
- ❖ Four final states from $B_s^0 \rightarrow J/\psi \phi \rightarrow \mu\mu KK$ are fitted to a common vertex
- ❖ Mass constrain of $J/\psi \rightarrow \mu\mu$; mass cut on $1008.5 < m(KK) < 1030.5$ MeV
- ❖ Decay time of $B_s^0 > 0.2$ ps
- ❖ Primary vertex is chosen as the one with least a_0 , calculated based on the B_s^0 vertex and momentum direction
- ❖ One track assumed to be π from the primary vertex




S: double Gaussian; B: Exponential

Search for the X(5568) in $B_s^0\pi^\pm$ final states



$p_T(B_s^0) > 10$ GeV



$p_T(B_s^0) > 15$ GeV

No obvious X(5568) is observed!


Search for the $X(5568)$ in $B_s^0\pi^\pm$ final states

Mass range: 5550–5700 MeV

Width: 21.9 MeV


$P_T(B_s^0) > 10$ GeV

Upper limit on the production rate is set



$$\rho_X \equiv \frac{\sigma(pp \rightarrow X + \text{anything}) \times \mathcal{B}(X \rightarrow B_s^0 \pi^\pm)}{\sigma(pp \rightarrow B_s^0 + \text{anything})} = \frac{N(X)}{N(B_s^0)} \times \frac{1}{\epsilon^{\text{rel}}(X)}$$


Observation of an Excited B_c^\pm Meson State



Observation of an Excited B_c^\pm Meson State


1. Select the J/Ψ into two opposite charged muons:

$p_T(\text{high}) > 6 \text{ GeV}$; $p_T(\text{low}) > 4 \text{ GeV}$; vertex fit;
mass constrain to PDG;



2. Select B_c^\pm by adding a pion:

$p_T > 4 \text{ GeV}$; common vertex to J/Ψ ;
cut on the impact parameter of the pion;
 $p_T(B_c^\pm) > 15 \text{ GeV}$ (18 GeV) for 7 TeV (8 TeV)



Observation of an Excited B_c^\pm Meson State

3. Select two pions from the primary vertex:



$$pT > 400 \text{ MeV}$$

4. Fit the mass difference distribution:

Gaussian for signal, third-order polynomial for background



Significance of the new
structure

$> 5 \sigma$




Fine structure?

With the similar technique but larger sample, both CMS and LHCb observed two structures, and updated results on ATLAS to be expected .




arXiv:1904.0008



PHYSICAL REVIEW LETTERS 122, 132001

ϕ_s and $\Delta\Gamma_s$ measurement in the $B_s^0 \rightarrow J/\Psi \phi$

$$V_{us}V_{ub}^* + V_{cs}V_{cb}^* + V_{ts}V_{tb}^* = 0$$




$$\phi_s = -2 \beta_s = -0.0363 \pm 0.016 \text{ rad}$$

Charles et al, PRD84,033005

ϕ_s , the CP violating phase is defined as the phase difference between mixing amplitude and decay amplitude; In SM, it is small, and related to CKM matrix.

World averaged results before Moriond 2019




From HFLAV

New result based on 80 fb^{-1} at 13 TeV will be discussed here

Methodology

Flavour tagger: OST (opposite-side-tagging); lepton charge in semi-leptonic decay of B meson provides strong discrimination.






CP State tagger: CP even if $L = 0$ or 2 ; CP odd if $L = 1$. L is the orbital angular momentum.

Tagger: weighted sum of charge in a cone

$$Q_x = \frac{\sum_i^{N \text{ tracks}} q_i \cdot (p_{Ti})^\kappa}{\sum_i^{N \text{ tracks}} (p_{Ti})^\kappa},$$

Tag method	Efficiency [%]	Effective Dilution [%]	Tagging Power [%]
Tight muon	4.50 ± 0.01	43.8 ± 0.2	0.862 ± 0.009
Electron	1.57 ± 0.01	41.8 ± 0.2	0.274 ± 0.004
Low- p_T muon	3.12 ± 0.01	29.9 ± 0.2	0.278 ± 0.006
Jet	5.54 ± 0.01	20.4 ± 0.1	0.231 ± 0.005
Total	14.74 ± 0.02	33.4 ± 0.1	1.65 ± 0.01



Fit results with RUN 2 data







Simultaneous un-binned maximum-likelihood fit contains nine parameters:

$$\Delta\Gamma_s, \phi_s, \Gamma_s, |A_0(0)|^2, |A_{\parallel}(0)|^2, \delta_{\parallel}, \delta_{\perp}, |A_S(0)|^2 \text{ and } \delta_S.$$


Information used in the fit: mass of B_s^0 ; proper decay time and its uncertainty; tagging probability; the transversity angles (defined in the next page)

Parameter	Value	Statistical uncertainty	Systematic uncertainty
ϕ_s [rad]	-0.068	0.038	0.018
$\Delta\Gamma_s$ [ps ⁻¹]	0.067	0.005	0.002
Γ_s [ps ⁻¹]	0.669	0.001	0.001
$ A_{\parallel}(0) ^2$	0.219	0.002	0.002
$ A_0(0) ^2$	0.517	0.001	0.004
$ A_S(0) ^2$	0.046	0.003	0.004
δ_{\perp} [rad]	2.946	0.101	0.097
δ_{\parallel} [rad]	3.267	0.082	0.201
$\delta_{\perp} - \delta_S$ [rad]	-0.220	0.037	0.010

Fit projection




Result: combined with RUN1 result




Parameter	Value	Statistical uncertainty	Systematic uncertainty
$\phi_s [\text{rad}]$	-0.076	0.034	0.019
$\Delta\Gamma_s [\text{ps}^{-1}]$	0.068	0.004	0.003
$\Gamma_s [\text{ps}^{-1}]$	0.669	0.001	0.001
$ A_{\parallel}(0) ^2$	0.220	0.002	0.002
$ A_0(0) ^2$	0.517	0.001	0.004
$ A_S ^2$	0.043	0.004	0.004
$\delta_{\perp} [\text{rad}]$	3.075	0.096	0.091
$\delta_{\parallel} [\text{rad}]$	3.295	0.079	0.202
$\delta_{\perp} - \delta_S [\text{rad}]$	-0.216	0.037	0.010

Result: combined with LHCb new result




New HFLAV average
 $\phi_s = -0.0544 \pm 0.0205$
 $\Delta\Gamma_s = 0.0762 \pm 0.0033 \text{ ps}^{-1}$

ATLAS HL-LHC projection




B_s^0 (and B^0) $\rightarrow \mu\mu$ measurement



**New result based on 26 fb^{-1} at 13 TeV
2015+2016**

Methodology: take $B^+ \rightarrow J/\psi K^+$ as reference

$$\mathcal{B}(B_{(s)}^0 \rightarrow \mu^+ \mu^-) = \frac{N_{d(s)}}{\varepsilon_{\mu^+ \mu^-}} \times [\mathcal{B}(B^+ \rightarrow J/\psi K^+) \times \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)] \frac{\varepsilon_{J/\psi K^+}}{N_{J/\psi K^+}} \times \frac{f_u}{f_{d(s)}}$$






Abundant and well measured branching fraction!

Background


(Left) Particle reconstructed decays: lower dimuon invariant mass

(Middle) Peaking background: from muon misidentification


(Right) Continuum background: dominant, flat distribution; reduced with BDT.




Fits




(a)



(b)




(c)



(d)

4 BDT intervals, and the first two contribute mostly to background modelling

Results



● SM :

$$\text{Br}(B_s \rightarrow \mu\mu) = (3.65 \pm 0.23) \times 10^{-9}$$

$$\text{Br}(B^0 \rightarrow \mu\mu) = (1.06 \pm 0.09) \times 10^{-10}$$

● Best fit of Run 2 data :

$$\text{Br}(B_s \rightarrow \mu\mu) = (3.2 \pm 0.9) \times 10^{-9}$$

$$\text{Br}(B^0 \rightarrow \mu\mu) = (-1.3 \pm 2.1) \times 10^{-10}$$


● Run 1 + Run 2 result @ 95% CL

$$\text{Br}(B_s \rightarrow \mu\mu) = (2.8 \pm 0.8) \times 10^{-9}$$

$$\text{Br}(B^0 \rightarrow \mu\mu) < 2.1 \times 10^{-10}$$

B^0 limit is most stringent at the moment

ATLAS HL-LHC projection



Summary

1. $X(5568)$ is searched with ATLAS data, but no hint;
2. Excited B_c^\pm Meson State is observed with ATLAS data;
3. ϕ s and $\Delta\Gamma$ s are measured in the $B_s^0 \rightarrow J/\Psi \phi$, and results are consistent with SM
4. B_s^0 (and B^0) $\rightarrow \mu\mu$ are measured, no surprise

Thank you very much!