

Parton energy loss in dense QCD medium

张本威

(Ben-Wei Zhang)

华中师范大学

Central China Normal University

第18届全国中高能核物理大会 长沙 2019.6.21-25日



Outline

- Introduction
- Leading hadron productions
- Full jet observables
 - 1) Gauge boson tagged jets
 - 2) heavy flavor jet
 - 3) Jet charge
- Summary

Jet quenching



Parton energy has been proposed as an excellent probe of the hot/dense matter created at HIC.

Single Hadron Tomography



Jet quenching at RHIC and LHC

$$R_{AA} = \frac{\text{Yield}_{AA} / \langle N_{\text{binary}} \rangle_{AA}}{\text{Yield}_{pp}}$$




Fingerprints of jet quenching

领头强子




整体喷注


Leading hadron production




Leading hadron production



Full jets




Full jets



Large p_T hadrons in HIC

Jet quenching with higher twist approach

The parton energy loss due to medium-induced gluon radiation has been calculated with higher twist approach.



$$\begin{aligned} \tilde{D}_{q \rightarrow h}(z_h, \mu^2) &\equiv D_{q \rightarrow h}(z_h, \mu^2) + \int_0^{\mu^2} \frac{d\ell_T^2}{\ell_T^2} \frac{\alpha_s}{2\pi} \int_{z_h}^1 \frac{dz}{z} [\Delta\gamma_{q \rightarrow qg}(z, x, x_L, \ell_T^2) D_{q \rightarrow h}(z_h/z) \\ &+ \Delta\gamma_{q \rightarrow gq}(z, x, x_L, \ell_T^2) D_{g \rightarrow h}(z_h/z)] , \end{aligned}$$

$$\begin{aligned} \frac{1}{N_{\text{bin}}^{AB}(b)} \frac{d\sigma_{AB}^h}{dy d^2p_T} &= \sum_{abcd} \int dx_a dx_b f_{a/A}(x_a, \mu^2) f_{b/B}(x_b, \mu^2) \\ &\times \frac{d\sigma}{d\hat{t}}(ab \rightarrow cd) \frac{\langle \tilde{D}_c^h(z_h, Q^2, E, b) \rangle}{\pi z_c} + \mathcal{O}(\alpha_s^3). \end{aligned}$$


X Guo, X N Wang, PRL(2001); X Guo, X N Wang, NPA (2001);
BWZ, X N Wang, NPA(2003) ; BWZ, E Wang, X N Wang, PRL (2004)

η in heavy-ion collisions at NLO

12


- Production of eta meson in HIC has been calculated;
- Flavor composition has very small effect on the ratio η/π^0 .

$$(dE/dx)_{\text{rad}} = -C_2 \hat{q} L$$



$$G^{\eta, \pi^0}(p_T) = \frac{\int f_g(\frac{p_T}{z_h}) \cdot D_{g \rightarrow \eta, \pi^0}(z_h, p_T) \frac{dz_h}{z_h^2}}{\frac{1}{p_T} \frac{d\sigma_{\pi^0, \eta}}{dp_T}}$$



$$G^{\pi^0}(p_T) \approx G^{\eta}(p_T)$$



η/π^0 in HIC at NLO

13



- η/π^0 ratio is almost same (~ 0.5) for p+p, Au+Au and Pb+Pb collision.
- Prediction on η/π^0 ratio has been confirmed by ALICE.





ALICE, PRC 98(2018)044901

ρ production in A+A at NLO


- A broken SU(3) model has been utilized to get ρ FFs in p+p.
- Dominant contributions of quark fragmentations both in p+p and in A+A.



Production of ϕ in HIC at NLO




Identified meson in HIC at NLO



iEBE-VISHNU
hydro


Global extraction of \hat{q}



Jets in HIC

What is a jet?

- A jet is a spray of final-state particles roughly moving in the same direction and defined by jet finding algorithms.
- At LO pQCD, jet \approx parton.
- In pQCD local-parton-hadron duality (LPHD) is used
- Jet: more precise and powerful





$$E_T = \sum_{i \in \text{jet}} E_{T,i}$$

$$y = \sum_{i \in \text{jet}} y_i E_{T,i} / E_T$$

$$\phi = \sum_{i \in \text{jet}} \phi_i E_{T,i} / E_T$$

$$R_{ij} = \sqrt{(y_i - y_j)^2 + (\phi_i - \phi_j)^2}$$

Tagged jet production in HIC





"golden channel"

$Z^0 + \text{jet in A+A: Iaa}$


- A sharp transition from tagged jet suppression above $\sim p_T$ of Z to tagged jet enhancement below $\sim p_T$ of Z

$$I_{AA}^{\text{jet}}(R, \omega_{\min}) = \frac{1}{\langle N_{\text{bin}} \rangle} \frac{d\sigma_{AA}}{dp_T(Z) dp_T(Q)} \bigg/ \frac{d\sigma_{pp}}{dp_T(Z) dp_T(\text{jet})}$$

NLO



Z^0 + jet in A+A: asymmetry




LO +PS
(parton
shower)

$$x_{JV} = p_T^J / p_T^V$$

	$\Delta\langle x_{JZ} \rangle$			
p_T^Z (GeV)	40 – 50	50 – 60	60 – 80	80 – 120
CMS [24]	0.061 ± 0.059	0.123 ± 0.051	0.124 ± 0.052	0.068 ± 0.042
Rad. + Coll. $g = 2.0$	0.022	0.050	0.075	0.086
Rad. + Coll. $g = 2.2$	0.024	0.058	0.093	0.119

Angular correlation in Z+jet

- NLO calculations fail at angular difference $\sim \pi$;
- LO+PS calculations fail at small angular difference.
- Z+jet in A+A: NLO+PS+Eloss




CMS, PLB

722 (2013) 238


NLO +PS
+Eloss

Sherpa: NLO+PS

- Initial state parton shower(PS)
- Final state PS
- NLO matrix elements (ME)
- Signal process
- Fragmentation
- Hadron decays
- Underlying event
- QED radiation





Angular Correlation of Z+jet (I)




Angular Correlation of Z+jet (II)

- Suppression to Z+1jet processes is rather small;
- Considerable suppression to Z+ (>2)jets is observed due to jet quenching effect; kinematical cut: $p_T > 30$ GeV




Averaged number of tagged jets

$$R_{jZ} = N_{jZ}/N_Z$$



Momentum imbalance







$$\Delta\langle x_{jZ} \rangle = \langle x_{jZ} \rangle_{p+p} - \langle x_{jZ} \rangle_{Pb+Pb}$$


$p_T^Z \text{ (GeV)}$	40-50	50-60	60-80	> 80
CMS data	0.07 ± 0.106	0.12 ± 0.148	0.13 ± 0.158	0.06 ± 0.088
$\Delta\langle x_{jZ} \rangle$	0.075	0.106	0.128	0.143

W+jet in HIC


S Zhang's talk




double b-jet production



flavor creation (FCR)





gluon splitting (GSP)




flavor excitation (FEX)

p+p baseline for double b-jet

- Simulations with NLP+PS by Sherpa should give very nice descriptions on inclusive jets, dijets as well as double b-jet.




Mean values of momentum imbalance



Angle correlation of double b-jet


W Dai's talk



D⁰-jet in p+p

- D-jet denotes a jet with D meson, and CMS has measured radial distributions of D-jet in p+p and A+A.



$$r = \sqrt{(\Delta\phi_{JD})^2 + (\Delta\eta_{JD})^2}$$



D⁰-jet in A+A



- D-jet denotes a jet with D meson, and CMS has measured radial distributions of D-jet in p+p and A+A.

S Wang's talk



Z tagged b-jet production (I)

- B-jet production in association with Z boson in HIC could be calculated in the same formalism (NLO+PS+Eloss)





$$\Delta\phi_{Zb} = |\phi_Z - \phi_{bjet}|$$

Z tagged b-jet production (II)

- B-jet production in association with Z boson in HIC could be calculated in the same formalism (NLO+PS+Eloss)

$$\Delta R_{zb} = \sqrt{(\Delta\phi_{zb})^2 + (\Delta\eta_{zb})^2}$$

$$A_{Zbb} = \frac{\Delta R_{Zb}^{max} - \Delta R_{Zb}^{min}}{\Delta R_{Zb}^{max} + \Delta R_{Zb}^{min}}$$





Jet charge in p+p

- Proposed by Feynman & Field (1978)
- Very useful to discriminate q/g

$$Q_{\kappa}^i = \sum_{h \in \text{jet}} z_h^{\kappa} Q_h$$



Krohn, Schwartz, T. Lin, Waalewijn, PRL (2013)



Jet charge in A+A

- R_{CP} of jet charge in A+A is larger than unity.
- Very useful to discriminate q/g

$$R_{CP} = \frac{\langle Q_j \rangle_{PbPb_{0-10\%}}}{\langle Q_j \rangle_{PbPb_{60-80\%}}}$$



S Chen, BWZ, E Wang (2019)

Summary

- A systematic study of identified mesons at NLO in HIC has been made.
- A framework of combining NLO+PS for initial hard production with parton energy loss in the QGP has been developed.
- Our calculations provide nice descriptions of experimental data on full jet observables.
- Calculations of pQCD with final-state jet quenching and initial-state CNM can explain a huge amount of data on hard probes with a unified picture.

Quark Matter 2019 at Wuhan

The header features a dark blue background with various colorful particle physics diagrams, including quark-antiquark pairs and meson-like structures, scattered across the top. Below these, a stylized orange and white cityscape of Wuhan is visible, with the Yangtze River and bridges. The text 'QUARK MATTER 2019' is prominently displayed in the center in a large, bold, orange and white font.

QUARK MATTER 2019


Wuhan, China 4-9 November

- 国际高能核物理领域规模最大、最重要的学术会议
- 预计~800人参加
- 华中师范大学和华南师范大学主办
- 清华大学、北京大学、复旦大学、中国科技大学、山东大学、中国科学院大学、湖州大学、中科院近代物理所等协办
- 会议报告 (abstract) 提交截止时间2019年7月1日
- 会议网站: <http://qm2019.ccnu.edu.cn/>

Backup




Jet charge in p+p



Z+jet in p+p: NLO+PS

- Results with NLO+PS by Sherpa give good descriptions on angular correlation and momentum imbalance of in p+p



Linear Boltzmann Transport Model

- Elastic scattering:

$$\begin{aligned}
 p_1 \cdot \partial f_1(p_1) &= - \int dp_2 dp_3 dp_4 (f_1 f_2 - f_3 f_4) |M_{12 \rightarrow 34}|^2 \\
 &\times (2\pi)^4 \delta^4(P_1 + P_2 - P_3 - P_4) \\
 dp_i &\equiv \frac{d^3 p_i}{2E_i (2\pi)^3}, \quad |M_{12 \rightarrow 34}|^2 = C g^2 (s^2 + u^2) / (t + \underline{\mu}^2)^2 \\
 f_i &= 1 / (e_i^{p \cdot u / T} \pm 1) \quad (i = 2, 4), \quad f_i = (2\pi)^3 \delta^3(\vec{p} - \vec{p}_i) \delta^3(\vec{x} - \vec{x}_i) \quad (i = 1, 3)
 \end{aligned}$$

X N Wang, Y Zhu, PRL(2013); He, Luo, Wang, Zhu, PRC (2015)

- Inelastic scattering by the higher twist approach:

$$\frac{dN_g}{dx dk_{\perp}^2 dt} = \frac{6\alpha_s C_A P(x) \hat{q}}{\pi k_{\perp}^4} \left(\frac{k_{\perp}^2}{k_{\perp}^2 + x^2 M^2} \right)^2 \sin^2 \left(\frac{t - t_i}{2\tau_f} \right)$$

Guo, X N Wang, PRL(2002); BWZ, X Wang, NPA(2003);

BWZ, E Wang, X N Wang, PRL (2004); Majumder, PRD(2012)

Inclusive jet and b-jet in HIC

