



BESIII 上 Lambda 超子极化的测量

平荣刚


(For BESIII Collaboration)

Base on: BESIII, arXiv: 1808.08917,
[Nature Physics \(2019\)](#)

Institute of High Energy Physics, CAS
pingrg@ihep.ac.cn

Beijing Electron Positron Collider (BEPC)

beam energy: 1.0 – 2.3 GeV



2004: started BEPCII upgrade,
BESIII construction

2008: test run

2009 - now: BESIII physics run

- 1989-2004 (BEPC):


$$L_{\text{peak}} = 1.0 \times 10^{31} / \text{cm}^2 \text{s}$$

- 2009-now (BEPCII):


$$L_{\text{peak}} = 1.0 \times 10^{33} / \text{cm}^2 \text{s}$$

BESIII Detector


MDC



R inner: 63mm ;
R outer: 810mm
Length: 2582 mm
Layers: 43



CsI(Tl) EMC




Crystals: 28 cm (15 X_0)

Barrel: $|\cos\theta| < 0.83$

Endcap:

$0.85 < |\cos\theta| < 0.93$

RPC MUC



BMUC: 9 layers – 72 modules
EMUC: 8 layers – 64 modules

TOF


BTOF: two layers

ETOF: 48 scintillators for each

MRPC --- new ETOF



BESIII J/ψ data sets



The analysis based on
2009+2012 J/ψ data:

~1.31 billion decays

Total 10.047 billion J/ψ decays we have

Role of polarization physics

- Probing of spin degree freedom

Test the dynamic of SM and low energy hadron interaction

Existant exp. : RHIC, Jlab, GRAAL, CERN and DESY

Spin observable, spin-dependent structure function and parton distribution


Spin crisis at eighties

- BEPCII/BESIII, unpolarized beam, inaccessible polarization of final state by BESIII


- Polarized beam for post-BEPCII options, CPV in tau decay, Hyperon weak decay,.....
- Useful tool: transverse polarization of hyperon, spontaneous production at e^+e^- collision
- $\Lambda \rightarrow p\pi^-$ decay plays important role in particle physics

Transverse polarization of baryons in e^+e^- collisions

FFs in continuum



J/ψ decays



$$\Gamma_\mu^e(k_1, k_2) = -ie_\psi \gamma_\mu$$

$$\Gamma_\mu^\Lambda(p_1, p_2) =$$

$$-ie_g \left[G_M^\psi \gamma_\mu - \frac{2M}{Q^2} (G_M^\psi - G_E^\psi) Q_\mu \right]$$

Time likespin $\frac{1}{2}$ baryon FFs:

Dubnickova, Dubnicka, Rekaló

Nuovo Cim. A109 (1996) 241


Gakh, Tomasi-Gustafsson Nucl.Phys. A771 (2006) 169

Czyz, Grzelinska, Kuhn PRD75 (2007) 074026


Fäldt EPJ A51 (2015) 74; EPJ A52 (2016) 141

W. Lu, et.al., Phys.Lett., B368, 261 (1996)

Transverse polarization of baryons in e^+e^- collisions



Unpolarized e^+e^- beam $\rightarrow$
transverse polarization baryon



$$\bar{P}_Y(\cos \theta_\Lambda) = \frac{\sqrt{1 - \alpha_\psi^2} \cos \theta_\Lambda \sin \theta_\Lambda}{1 + \alpha_\psi \cos^2 \theta_\Lambda} \sin(\Delta\Phi)$$

P_Y along $\mathbf{k}_{e^+} \times \mathbf{p}_\Lambda$



$\Lambda \rightarrow p\pi^-$ as polarimeter

$$\frac{dN}{d\Omega} = \frac{1}{4\pi} (1 + \alpha_{\Lambda} \vec{P} \cdot \hat{q}) = \frac{1}{4\pi} (1 + \alpha_{\Lambda} P_{\Lambda} \cos\theta_p)$$


Lee-Yang parameters:

$$\alpha = \frac{2 \operatorname{Re}(S^* P)}{|S|^2 + |P|^2}, \quad \beta = \frac{2 \operatorname{Im}(S^* P)}{|S|^2 + |P|^2}, \quad \gamma = \frac{|S|^2 - |P|^2}{|S|^2 + |P|^2}$$

Note: $\alpha^2 + \beta^2 + \gamma^2 = 1$



C- and P-transformation



$$\alpha_{\Lambda} = \frac{|\mathbf{B}_+|^2 - |\mathbf{B}_-|^2}{|\mathbf{B}_+|^2 + |\mathbf{B}_-|^2}, \alpha_{\bar{\Lambda}} = \frac{|\bar{\mathbf{B}}_+|^2 - |\bar{\mathbf{B}}_-|^2}{|\bar{\mathbf{B}}_+|^2 + |\bar{\mathbf{B}}_-|^2}$$

CP invariance:

$$\bar{\mathbf{B}}_{-\lambda_p} = \eta_{\Lambda} \eta_p \eta_{\pi} (-1)^{s_{\Lambda} - s_p - s_{\pi}} \mathbf{B}_{\lambda_p} = -\mathbf{B}_{\lambda_p}$$

If CP invariance:

$$\alpha_{\Lambda} = -\alpha_{\bar{\Lambda}}$$

CP-odd observables

$$\Delta = \frac{\Gamma - \bar{\Gamma}}{\Gamma + \bar{\Gamma}}$$

$$A = \frac{\alpha + \bar{\alpha}}{\alpha - \bar{\alpha}}$$

$$B = \frac{\beta + \bar{\beta}}{\beta - \bar{\beta}}$$


$$B' = \frac{\beta + \bar{\beta}}{\alpha - \bar{\alpha}}$$

- A_Λ at 10^{-5} level by CKM matrix, PDG: 0.006 ± 0.021
- Asymmetries B, B' require knowledge of both parent and daughter polarization

Proton polarization from $\Lambda \rightarrow p\pi^-$

$$\vec{P}_p = \frac{(\alpha + \vec{P}_\Lambda \cdot \hat{q})\hat{q} + \beta (\vec{P}_\Lambda \times \hat{q}) + \gamma (\hat{q} \times (\vec{P}_\Lambda \times \hat{q}))}{(1 + \alpha \vec{P}_\Lambda \cdot \hat{q})}$$

- If $P_\Lambda = 0$ then $P_p = \alpha p \cdot q$
- T-odd transverse polarization
 $\beta \neq 0$
- If CP is conserved :
 $\alpha = -\bar{\alpha}, \beta = -\bar{\beta},$
 $\gamma = \bar{\gamma} \text{ and } \Gamma = \bar{\Gamma}$



Previous Measurements

2018 PDG list

α_- **FOR $\Lambda \rightarrow p\pi^-$**

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.642 ± 0.013	OUR AVERAGE				
0.584 ± 0.046	8500	ASTBURY	1975	SPEC	
0.649 ± 0.023	10325	CLELAND	1972	OSPK	
0.67 ± 0.06	3520	DAUBER	1969	HBC	From Ξ decay
0.645 ± 0.017	10130	OVERSETH	1967	OSPK	Λ from $\pi^- p$
0.62 ± 0.07	1156	CRONIN	1963	CNTR	Λ from $\pi^- p$


α_+ **FOR $\bar{\Lambda} \rightarrow \bar{p}\pi^+$**

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
-0.71 ± 0.08	OUR AVERAGE				
$-0.755 \pm 0.083 \pm 0.063$	$\approx 8.7k$	ABLIKIM	2010	BES	$J/\psi \rightarrow \Lambda \bar{\Lambda}$
-0.63 ± 0.13	770	TIXIER	1988	DM2	$J/\psi \rightarrow \Lambda \bar{\Lambda}$

Most earlier measurement on α

- CNTR 实验, $\pi^- + p \rightarrow \Lambda + K^0$
- 非极化的 Λ 衰变产生的 p 的极化为 α , 末态 p 的极化可以通过火花室测量

Phys.Rev. 129 (1963) 1795-1807



$$W(\psi) = 1 + \alpha S \sin \epsilon \cos \psi$$




FIG. 1. Schematic diagram showing arrangement of apparatus. An example of an event has been sketched in.

$$\alpha = - \frac{2}{\pi} \frac{1}{\langle S \rangle \langle \sin \epsilon \rangle} \frac{N_+ - N_-}{N_+ + N_-},$$

1156 events


$$\langle S \rangle = 0.565$$

$$\langle \sin \epsilon \rangle = 0.84,$$

$$\alpha = 0.62.$$

PS185 at LEAR : $p\bar{p} \rightarrow \Lambda\bar{\Lambda}$

Assuming C invariance in the production $P_{y\Lambda} = P_{y\bar{\Lambda}}$



LEAR: Low Energy antiproton Ring

Spin observables were shown

$$I(\theta_p) = I_0(1 + \alpha P_y \cos \theta_p)$$

$$\text{Extraction of } \alpha P_y; \alpha P_y = \frac{\sum \cos \theta_y^{(i)}}{\sum \cos^2 \theta_y^{(i)}}$$

requires $\eta(\theta_y) = \eta(\pi - \theta_y)$

From a combined total sample of 95832 events:

$$A_\Lambda \approx \frac{\alpha_\Lambda P_\Lambda + \alpha_{\bar{\Lambda}} P_{\bar{\Lambda}}}{\alpha_\Lambda P_\Lambda - \alpha_{\bar{\Lambda}} P_{\bar{\Lambda}}} = 0.013 \pm 0.022 \quad (\text{PRC54,1877(1996)})$$

$\bar{p}$ momentum:
 $1.4765\text{GeV} \sim 1.5075\text{GeV}$





FIG. 1. Overview of the PS185 detector system. (1) segmented neutral trigger target, (2) multiwire proportional chambers (MWPC's), (3) multiwire drift chambers (MWDC's), (4) scintillator drift chambers, and (5) solenoid "baryon identifier" with drift chambers. The lower part of the figure shows a detail of the segmented target.


DM2 : $e^+e^- \rightarrow J/\psi \rightarrow \Lambda\bar{\Lambda}$

- 1847 $\Lambda\bar{\Lambda} \rightarrow p\pi^- \bar{p}\pi^+$ from $8.6 \times 10^6 e^+e^- \rightarrow J/\psi$
- No explicit assumption about Λ and $\bar{\Lambda}$ polarization needed
- CP violation signal appears in the product $\alpha_\Lambda \alpha_{\bar{\Lambda}} \mathbf{a} \cdot \mathbf{b}$,
where $\mathbf{a}$ and $\mathbf{b}$ are the p and $\bar{p}$ unit vector in the Λ and $\bar{\Lambda}$ rest frame
- Not consider the transverse polarization of Λ and $\bar{\Lambda}$

Fixing $\alpha_\Lambda = 0.642$ and fit yields

$$\alpha_{\bar{\Lambda}} = -0.63 \pm 0.13.$$


$$\Rightarrow A_\Lambda = 0.01 \pm 0.10$$





BESII with 58 million J/ψ decays

$$\frac{d\sigma}{d\Omega} \propto (1 - \alpha) \sin^2 \theta [1 + \alpha_{\Lambda} \alpha_{\bar{\Lambda}} (\cos \theta_1 \cos \bar{\theta}_1 + \sin \theta_1 \sin \bar{\theta}_1 \cos(\phi_1 + \bar{\phi}_1))] - (1 + \alpha)(1 + \cos^2 \theta) (\alpha_{\Lambda} \alpha_{\bar{\Lambda}} \cos \theta_1 \cos \bar{\theta}_1 - 1),$$

Spin-correlation



Same as DM2



~ 9000 events selected

$$\alpha_{\bar{\Lambda}} = -0.755 \pm 0.083 \pm 0.063,$$



$$A = -0.081 \pm 0.055 \pm 0.059,$$

PRD 81, 012003 (2010)

Hyperon decay parameters @ BESIII

$$e^+ e^- \rightarrow J/\psi \rightarrow \Lambda \bar{\Lambda} \rightarrow p \bar{p} \pi^+ \pi^-$$

Event selection:



- ✓ No PID request for charged tracks
- ✓ $\Lambda/\bar{\Lambda}$ reconstructed with second vertex fit
- ✓ Events reconstructed with kinematic fit

Hyperon decay parameters @ BESIII

$$e^+ e^- \rightarrow J/\psi \rightarrow \Lambda \bar{\Lambda} \rightarrow p \bar{p} \pi^+ \pi^-$$


G. Faldt and A. Kupsc
PLB, 772, 16 (2017)
EPJA, 52, 141 (2016)
EPJA, 51, 74 (2015).

$$\frac{d\sigma}{d\Omega} \propto T_0 + \sqrt{1 - \alpha_{J/\psi}^2} \sin(\Delta) (\alpha_\Lambda T_3 + \alpha_{\bar{\Lambda}} T_4) \\ + \alpha_\Lambda \alpha_{\bar{\Lambda}} [T_1 + \sqrt{1 - \alpha_{J/\psi}^2} \cos(\Delta) T_2 + \alpha_{J/\psi} T_5],$$



T_0 : angular distribution of Λ and $\bar{\Lambda}$

T_3, T_4 , transverse polarization

T_1, T_2, T_5 : spin correlation





$$e^+e^- \rightarrow J/\psi \rightarrow \Lambda\bar{\Lambda} \rightarrow p\bar{r}\pi^+\pi^-, p\bar{n}\pi^-\pi^0$$





$$e^+e^- \rightarrow J/\psi \rightarrow \Lambda\bar{\Lambda} \rightarrow p\bar{n}\pi^0\pi^+$$

47,009 events selected



Results of simultaneous fit to two data sets



Parameters	This measurement	Previous results
α_ψ	$0.461 \pm 0.006 \pm 0.007$	0.469 ± 0.027 [19]
α_-	$0.750 \pm 0.009 \pm 0.004$	0.642 ± 0.013 [8]
α_+	$-0.758 \pm 0.010 \pm 0.007$	-0.71 ± 0.08 [8]
$\bar{\alpha}_0$	$-0.692 \pm 0.016 \pm 0.006$	—
$\Delta\Phi$ (rad)	$0.740 \pm 0.010 \pm 0.008$	$\Delta\Phi = 42.4^\circ \pm 0.6^\circ (sta) \pm 0.5^\circ (sys.)$
A_Λ	$-0.006 \pm 0.012 \pm 0.007$	0.006 ± 0.021 [8]
$\bar{\alpha}_0/\alpha_+$	$0.913 \pm 0.028 \pm 0.012$	—

Systematic uncertainties


- Challenge to estimate detector asymmetry in tracking, and kinematic fit
- Systematic uncertainties well controlled

Relative systematic uncertainties (%)

Source	α_ψ	α_-	α_+	$\bar{\alpha}_0$	$\Delta\Phi$
Tracking, π^0 , $\bar{n}$	1.5	0.1	0.3	0.6	1.1
Kinematic fit	0.2	0.1	0.8	0.6	0.0
Fit method	0.0	0.5	0.4	0.4	0.1
Total	1.5	0.5	0.9	0.8	1.1


- Justify results with $J/\psi \rightarrow \Xi^- \bar{\Xi}^+ \rightarrow \Lambda \bar{\Lambda} \pi^+ \pi^- \rightarrow p \bar{p} 2(\pi^+ \pi^-)$

BESIII: 0.750 ± 0.010



(a) α_- for $\Lambda \rightarrow p\pi^-$

BESIII: -0.758 ± 0.012



(b) α_+ for $\bar{\Lambda} \rightarrow \bar{p}\pi^+$

BESIII, arXiv: 1808.08917,

[Nature Physics \(2019\)](#)

[高能新闻](#) (BESIII发现 Λ 超子横向极化并精确测量其衰变参数)

Quick response from PDG2019

α_- **FOR** $\Lambda \rightarrow p\pi^-$


2019 PDG list

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.750 \pm 0.009 \pm 0.004$	420k	ABLIKIM 2018AG	BES3	J/ψ to $\Lambda\bar{\Lambda}$
••• We do not use the following data for averages, fits, limits, etc. •••				
0.584 ± 0.046	8500	ASTBURY 1975	SPEC	
0.649 ± 0.023	10325	CLELAND 1972	OSPK	
0.67 ± 0.06	3520	DAUBER 1969	HBC	From Ξ decay
0.645 ± 0.017	10130	OVERSETH 1967	OSPK	Λ from $\pi^- p$
0.62 ± 0.07	1156	CRONIN 1963	CNTR	Λ from $\pi^- p$



α_+ **FOR** $\bar{\Lambda} \rightarrow \bar{p}\pi^+$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$-0.758 \pm 0.010 \pm 0.007$	420k	ABLIKIM 2018AG	BES3	J/ψ to $\Lambda\bar{\Lambda}$
••• We do not use the following data for averages, fits, limits, etc. •••				
$-0.755 \pm 0.083 \pm 0.063$	$\approx 8.7k$	ABLIKIM 2010	BES	$J/\psi \rightarrow \Lambda\bar{\Lambda}$
-0.63 ± 0.13	770	TIXIER 1988	DM2	$J/\psi \rightarrow \Lambda\bar{\Lambda}$

$$e^+e^- \rightarrow \gamma^* \rightarrow \Lambda\bar{\Lambda} \rightarrow p\bar{p}\pi^+\pi^-$$



Λ transverse polarization



- data set: 2.396 GeV, $L=66.9 \text{ pb}^{-1}$
- 555 candidate events
- $\Delta\Phi = 37^\circ \pm 12^\circ \pm 6^\circ$
- Maximum polarization degree is $\sim 30\%$

Ongoing analyses at BESIII

- $J/\psi \rightarrow \Lambda \bar{\Lambda}$, arXiv: 1808.08917, Nature physics(2019)
- $e^+e^- \rightarrow \Lambda \bar{\Lambda}$, arXiv: 1903.09421, Submit to PRL
- $\Lambda_c^+ \rightarrow \Lambda \pi^+, \Sigma^+ \pi^0, \Sigma^0 \pi^+, K_S p$, arXiv:1905.04707
- $J/\psi, \psi' \rightarrow \Sigma^- \bar{\Sigma}^+, \Xi^- \bar{\Xi}^+$
- $\psi' \rightarrow \Omega^- \bar{\Omega}^+$

Summary

- $\Lambda/\bar{\Lambda}$ transverse polarization (TP) significantly observed at BESIII in J/ψ or continuum processes
- BESIII 10 billion J/ψ data provides us chances to access hyperon physics.
- Polarized beam in the future super-tau charm facility (STCF) help to improve the precision
- [Hyperon workshop](#), 7-8, June @Fudan Univ.

Thanks for your attention!