



清華大學

Sequential Hadronization with Charm Conservation in Heavy Ion Collisions

Jiaxing Zhao (Tsinghua University)

arXiv: 1805.10858

In collaboration with:

Shuzhe Shi, Nu Xu and Pengfei Zhuang

The 2nd FCPPL quarknonium production workshop

Outline



Part1. Introduction and motivation

Part2. Charm Conservation and Hadronization Sequential

Part3. Results and Analysis

Part4. Summary and Outlook



Enhancement in Charm Sector in HIC



L. Zhou [STAR Collaboration] NPA967. 620(2017) *J. Adam et al. [ALICE Collaboration] JHEP. 1603. 082(2016)*

Significantly enhanced in Heavy ion collisions vs. pp collisions!

Enhancement in Charm Sector in HIC





L. Zhou [STAR Collaboration] NPA967. 620(2017) J. Adam et al. [ALICE Collaboration] JHEP. 1603. 082(2016)

Significantly enhanced in Heavy ion collisions vs. pp collisions!

- Induced by strangeness (D_s) enhancement?

Strange quark thermal production in QGP.
 ($M_s \sim 100$ MeV, $T \sim 300$ MeV)



Charm Quark Conservation



L. Zhou [STAR Collaboration] NPA967. 620(2017)

Charm Conservation play an important role!

Charm Quark Conservation





L. Zhou [STAR Collaboration] NPA967. 620(2017)

Charm Conservation play an important role!

- Ds enhancement \rightarrow D0 suppression \rightarrow extra Ds/D0 enhancement!
 ($M_c \sim 1500$ MeV, $T \sim 300$ MeV)

Charm Quark Conservation





L. Zhou [STAR Collaboration] NPA967. 620(2017)

Charm Conservation play an important role!

- Ds enhancement \rightarrow D0 suppression \rightarrow extra Ds/D0 enhancement!
 (Mc \sim 1500 MeV, T \sim 300 MeV)

Q: How to realize the charm conservation self-consistently?


Hadronization in HIC



Matsui & Satz 1986

A hot and dense matter(QGP) produced in HIC!
Initial Produced charm hadrons would be "melted" in it and will be regenerated at final stage!


Hadronization Sequence



H. Satz. J. Phys. G32, R25(2006)

Lattice results: open charm hadrons dissolve close but higher than light hadrons!

*A. Bazavov, HT. Ding, S. Mukherjee et al.
Phys. Lett. B737,210(2014)*



larger binding energy

survive at higher temperature

hadronize earlier.



Hadronization Sequence

Two-body Dirac equation is used for open charm bound states

H. Crater, J. Yoo and C. Wong. PRD 79. 034011(2009) S. Shi, X. Guo and Pengfei Zhuang. PRD 88. 014021(2013)

Solve 2-body Dirac equation + lattice Free energy
get the binding energy and wavefunction!

The dissociation temperature T_d : $\epsilon(T_d) = 0$



Coalescence Hadronization

Coalescence mechanism play an important role in hadrons production in heavy ion collisions.

$$\frac{dN}{d^2\mathbf{P}_T d\eta} = C \int \frac{P^\mu d\sigma_\mu(R)}{(2\pi)^3} \int \frac{d^4r d^4p}{(2\pi)^3} F(r_1, p_1, r_2, p_2) W(r, p)$$

Coalescence Hadronization

Coalescence mechanism play an important role in hadrons production in heavy ion collisions.

$$\frac{dN}{d^2\mathbf{P}_T d\eta} = C \int \frac{P^\mu d\sigma_\mu(R)}{(2\pi)^3} \int \frac{d^4r d^4p}{(2\pi)^3} F(r_1, p_1, r_2, p_2) W(r, p)$$

- The hadronization hypersurface determined by hydrodynamics and dissociation temperature.

$$\partial_\mu T^{\mu\nu} = 0 \quad \partial_\mu n^\mu = 0$$



Coalescence Hadronization

Coalescence mechanism play an important role in hadrons production in heavy ion collisions.

$$\frac{dN}{d^2\mathbf{P}_T d\eta} = C \int \frac{P^\mu d\sigma_\mu(R)}{(2\pi)^3} \int \frac{d^4r d^4p}{(2\pi)^3} F(r_1, p_1, r_2, p_2) W(r, p)$$

- The covariant wigner function can be constructed by wavefunction.

$$W(r, p) = \int d^4y e^{-ipy} \psi(r + \frac{y}{2}) \psi(r - \frac{y}{2})$$



Quark Distribution Function

Light quark reach chemical equilibrium in QGP


$$f(r, p) = \frac{N_s}{e^{u^\mu(r)p_\mu/T(r)} + 1}$$

Quark Distribution Function

Light quark reach chemical equilibrium in QGP

$$f(r, p) = \frac{N_s}{e^{u^\mu(r)p_\mu/T(r)} + 1}$$

Heavy quark evolution can be described by transport equation



$$f_c = \rho_c(r) [\alpha f_{th} + \beta f_{pp}]$$


$$\rho_c(x|\mathbf{b}) = r(\tau) T_A(\mathbf{x}_T) T_B(\mathbf{x}_T - \mathbf{b}) \frac{\cosh \eta}{\tau} \frac{d\sigma_{pp}^{c\bar{c}}}{d\eta}$$

Charm conservation:

$$r(\tau) = \begin{cases} 1 & \tau \leq \tau_{D_s^+} \\ 1 - N_{D_s^+}/N_c & \tau_{D_s^+} < \tau \leq \tau_{D^0} \\ 1 - N_D/N_c & \tau_{D^0} < \tau \end{cases}$$


Results

Transverse momentum spectrum of D_s , D_0 and charm baryon




Results

Transverse momentum spectrum of Ds, D0 and charm baryon




Results

Transverse momentum spectrum of D_s , D_0 and charm baryon




Results

Transverse momentum spectrum of D_s , D_0 and charm baryon




Results



Solid line: sequential coalescence + charm conservation

Dashed line: simultaneous coalescence at T_c

Results



Solid line: sequential coalescence + charm conservation

Dashed line: simultaneous coalescence at T_c



Summary

1. We have built a framework to realize **sequential hadronization** with **charm conservation** in HIC!
2. Hadronization sequence of open charm mesons determined by **2-body Dirac equation**.
3. Reasonable agreement between our theoretical calculation and experiment data.



Need more exp. data to constrain the hadronization mechanism !

Outlook

Ds/D0 Ratio at High Baryon Density



RHIC BESII
Fixed-target BESIII
CBM, NICA, HIAF








In progressing...

THANK YOU !

BACKUP

For charmed mesons we can solve Two Body Dirac Equation to get wave function and masses!





Model calculations are lower than data at higher p_T .

1. Fragmentation function of heavy flavor baryons is not well understood?
2. Effect of the inner structure of Λ_c ?