



Status of the Real-Time Analysis project at LHCb

**The 5th China LHC Physics Workshop
Dalian, 2019/10/24**

SAUR Miroslav

On behalf of the LHCb collaboration

University of Chinese Academy of Sciences

A solid red square is positioned in the top left corner of the slide.

Outline

I. LHCb detector

II. Run II trigger conditions

III. Turbo stream

IV. Real-time Analysis project for Run III

LHCb detector 2010-2018

- Single-arm forward spectrometer focused on heavy flavor (b , c) physics
- Run I (7/8 TeV, 3 fb^{-1}), Run II (13 TeV, $\sim 6 \text{ fb}^{-1}$) + special runs (pPb, PbPb, SMOG)


Vertex Locator(vertex reconstruction)

- Impact parameter resolution: $20 \mu\text{m}$
- Decay time resolution: 45 fs ($\tau_B \sim 1.5 \text{ ps}$)

Tracking system(particle reconstruction)

- $\epsilon(\text{Tracking}) \sim 96\%$
- $\delta p/p \sim 0.5\%-1\%$ (5-200 GeV)
- $\sigma(m_{B \rightarrow hh}) \approx 22 \text{ MeV}$

JINST 3 (2008) S08005
IJMPA 30 (2015) 1530022



RICH: particle ID

- $\epsilon(K \rightarrow K) \sim 95\%$
- Mis-ID: $\epsilon(\pi \rightarrow K) \sim 5\%$

Magnet

- Bending power: 4 Tm

Muon system

- μ ID: $\epsilon(\mu \rightarrow \mu) \sim 97\%$
- Mis-ID: $\epsilon(\pi \rightarrow \mu) \sim 1-3\%$

Trigger during Run II

→ Run II (2015-2018) trigger system consisted of 3 stages

1) L0 Hardware trigger

- fast detectors
- CALO and MUON information


2) High Level Trigger 1

- 10 PB disk buffer
- Around two weeks of data taking

3) High Level Trigger 2




→ Around 500 specific HLT lines (particular decays)

→ Introduction of TESLA framework → Turbo stream




Trigger - alignment & calib.

- HLT1 samples are used for alignment and calibration
- Alignment procedure of the full tracker system run automatically at the beginning of each fill
- Based on Kalman filter
- Update if the variations are significant
- RICH calibration and alignment
- Time calibration of OT
- Calibration of ECAL




Requirements for trigger

- Triggering is a crucial part of data taking
- Decision of what physics can be recorded
- Resources demanding operation
- Hard constrains: Bandwith [GB/S] \approx Accept Rate [kHz] \times Event size [kB]
- Limiting factors: both hardware and software
- Raw data bandwidth scales up quadratically with luminosity
- During the Run II already significant rates: 45 kHz for bb, 1 MHz for cc




Turbo stream

TURBO (since 2015)




10 kB

TURBO SP new 2017



TURBO++ (since 2016)





70 kB Event size

- Selecting what information should be preserved and which can be discarded
- Extensively used during the Run II
 - Around 30 % of the full output is Turbo - almost all Charm physics
 - Approximately 2/3 lines keep raw detector information (Turbo SP)
- Significant reduction of data size → more events at same bandwidth

Persistence method	Average event size (kB)
Turbo	7
Selective persistence	16
Complete persistence	48
Raw event	69


Accomplishments of Turbo

- Turbo lines proved to be necessary for keeping a significant charm program during Run II
- Suitable for a broad range of physics – from high to low rate
- One of the key ideas of Upgrade




arXiv:1909.12273

PRL 122 (2019) 211803




LHCb Upgrade I (Run III)

- Luminosity will increase 5x times and collision energy to 14 TeV
- Aim is to maintain the same performance as during Run II



LHCb Upgrade I (Run III)

- Luminosity will increase 5x times and collision energy to 14 TeV
- Aim is to maintain the same performance as during Run II
- A large scale Upgrade!



«This is a new detector at the LHC»

LHCb Upgrade I - Physics






Type	Observable	Current precision	LHCb 2018	Upgrade (50 fb ⁻¹)	Theory uncertainty
B_s^0 mixing	$2\beta_s (B_s^0 \rightarrow J/\psi \phi)$	0.10 [9]	0.025	0.008	~ 0.003
	$2\beta_s (B_s^0 \rightarrow J/\psi f_0(980))$	0.17 [10]	0.045	0.014	~ 0.01
	$A_{fs}(B_s^0)$	6.4×10^{-3} [18]	0.6×10^{-3}	0.2×10^{-3}	0.03×10^{-3}
Gluonic penguin	$2\beta_s^{\text{eff}}(B_s^0 \rightarrow \phi\phi)$	–	0.17	0.03	0.02
	$2\beta_s^{\text{eff}}(B_s^0 \rightarrow K^{*0}\bar{K}^{*0})$	–	0.13	0.02	< 0.02
	$2\beta_s^{\text{eff}}(B^0 \rightarrow \phi K_S^0)$	0.17 [18]	0.30	0.05	0.02
Right-handed currents	$2\beta_s^{\text{eff}}(B_s^0 \rightarrow \phi\gamma)$	–	0.09	0.02	< 0.01
	$\tau^{\text{eff}}(B_s^0 \rightarrow \phi\gamma)/\tau_{B_s^0}$	–	5 %	1 %	0.2 %
Electroweak penguin	$S_3(B^0 \rightarrow K^{*0}\mu^+\mu^-; 1 < q^2 < 6 \text{ GeV}^2/c^4)$	0.08 [14]	0.025	0.008	0.02
	$s_0 A_{\text{FB}}(B^0 \rightarrow K^{*0}\mu^+\mu^-)$	25 % [14]	6 %	2 %	7 %
	$A_{\text{I}}(K\mu^+\mu^-; 1 < q^2 < 6 \text{ GeV}^2/c^4)$	0.25 [15]	0.08	0.025	~ 0.02
	$\mathcal{B}(B^+ \rightarrow \pi^+\mu^+\mu^-)/\mathcal{B}(B^+ \rightarrow K^+\mu^+\mu^-)$	25 % [16]	8 %	2.5 %	$\sim 10\%$
Higgs penguin	$\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-)$	1.5×10^{-9} [2]	0.5×10^{-9}	0.15×10^{-9}	0.3×10^{-9}
	$\mathcal{B}(B^0 \rightarrow \mu^+\mu^-)/\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-)$	–	$\sim 100\%$	$\sim 35\%$	$\sim 5\%$
Unitarity triangle angles	$\gamma (B \rightarrow D^{(*)}K^{(*)})$	$\sim 10\text{--}12^\circ$ [19, 20]	4°	0.9°	negligible
	$\gamma (B_s^0 \rightarrow D_s K)$	–	11°	2.0°	negligible
	$\beta (B^0 \rightarrow J/\psi K_S^0)$	0.8° [18]	0.6°	0.2°	negligible
Charm CP violation	A_Γ	2.3×10^{-3} [18]	0.40×10^{-3}	0.07×10^{-3}	–
	ΔA_{CP}	2.1×10^{-3} [5]	0.65×10^{-3}	0.12×10^{-3}	–

CERN/LHCC 2012-007


LHCb Upgrade I (Run III)

- LHCb has a very broad physics program
- High quality data requires a perfectly calibrated and aligned detector
- Have to process 5x bigger events at 30 times the rate, L0 removed
- From Run 3 all alignments and calibrations will be fully automatic and incorporated to the software trigger
- Around 70 % of data will go to Turbo




Idea of Real-Time Analysis

- Real-Time Analysis – efficient decision about data in the full online mode
- Keeping only a signal and suppress any unnecessary information about event




Idea of Real-Time Analysis

- Real-Time Analysis – efficient decision about data in the full online mode
- Keeping only a signal and suppress any unnecessary information about event
- Triggerless readout, full software trigger on 30 MHz (readout 40 MHz, around 40 Tb/s)



Structure of RTA project

- Subdetector-like organization
- 36 institutes, around 50 FTE (>100 people involved)
- A long term project also in charge of maintenance after deployment




Local participation

- Contributions from various institutions and individuals with various focus are needed now and also for future
- Currently four local group are involved
- Central China Normal University (2)
 - Main focus: study of ghost probability for tracking
 - XU Menglin, today 18:10
- Peking university (2)
 - Main focus: calorimeter related studies (combined with THU)
- Tsinghua university (4)
 - Main focus: calorimeter related studies (combined with PKU)
- University of Chinese Academy of Sciences (2)
 - Main focus: data challenges, automatic testing framework, software infrastructure




Data Preparation

- Data must be prepared for algorithms
 - Very CPU intensive - up to 50 % of data processing time
- Need of fast and optimized code
 - Trade-off between efficiency and speed? How much efficiency we can lose?




Data Occupancy effect

- Events with higher occupancy take longer to process – possibly critical effect
- However, such an events typically are not representative of signal topologies
- Most of algorithms scale linearly, issue with forward tracking



Multi-thread HLT1 sequence

- Run II HLT1 framework is a single-thread algorithm
- Porting to multi-thread architecture: 20 % gain just from more threads





Forward tracking - Kalman fit

- Kalman filter is important part of tracking algorithm but using significant part of HLT1 time
- Run I - Full material map and B field propagation
- Run II - Simplified model for the material map





- Run III - parametrization of Kalman filter by analytic function
- Current implementation is already much faster with only slightly worse performance (mainly for low p regions)




Data Challenges

- The Upgrade framework must be thoroughly tested before real running
- As the most of the data will be using Turbo lines, mistake in any part of data processing can be fatal
- An iterative procedure closely following progress of the whole RTA project and Upgrade itself
- Tests scrutinizing new tracking and alignment algorithms for VELO and SciFi done and evaluated
- Current work is focusing on reconstruction algorithms
- Work also includes preparation of non-standard simulation samples with various conditions for all possible tests
- Even though not a work on some exact sub-detector, this task requires to have a overview knowledge of each sub-detector and coordination work between particular groups
- Together with Early Measurements Task Force the Data Challenges group will help to prepare evaluate conditions for a early Run III measurements and give proponents the opportunity to test their methods and code on Run III data-like simulations


Latest progress



LHCb-FIGURE-2019-005



LHCb-FIGURE-2019-004



LHCb-FIGURE-2019-002

Conclusion

- LHCb detector finished two successful periods of data taking
- During the current long shutdown LHCb detector is going through a large-scale upgrade of both hardware and software part of detector
- 5x higher luminosity during Run III
- Run I utilized a 'HEP-standard' trigger strategy
- Run II shown a need for faster trigger system and usefulness of a partial data recording
- RTA is a novel approach for hadron collider experiments enable substantially increase the amount of recorded data
- Such a new approach is crucial for a future collider experiments on HL-LHC, SppC/FCC-hh, ...



Thank you for your attention




**BACKUP
Slides**



Planned LHCb upgrades




LHCb upgrade Phase II (Run V)



Real-Time Analysis at Run V

- Real-Time Analysis – efficient decision about data in the full online mode
- Run V – HL-LHC





LHCb upgrade Phase II (Run V)

Topics and observables	Experimental reach	Remarks
EW Penguins Global tests in many $b \rightarrow s\mu^+\mu^-$ modes with full set of precision observables; lepton universality tests; $b \rightarrow dl^+l^-$ studies	<i>e.g.</i> 440k $B^0 \rightarrow K^*\mu^+\mu^-$ & 70k $\Lambda_b^0 \rightarrow \Lambda\mu^+\mu^-$; Phase-II $b \rightarrow d\mu^+\mu^- \approx$ Run-1 $b \rightarrow s\mu^+\mu^-$ sensitivity.	Phase-II ECAL required for lepton universality tests.
Photon polarisation \mathcal{A}^Δ in $B_s^0 \rightarrow \phi\gamma$; $B^0 \rightarrow K^*e^+e^-$; baryonic modes	Uncertainty on $\mathcal{A}^\Delta \approx 0.02$; $\sim 10k \Lambda_b^0 \rightarrow \Lambda\gamma$, $\Xi_b \rightarrow \Xi\gamma$, $\Omega_b^- \rightarrow \Omega\gamma$	Strongly dependent on performance of ECAL.
$b \rightarrow cl^-\bar{\nu}_l$ lepton-universality tests Polarisation studies with $B \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$; τ^-/μ^- ratios with B_s^0 , Λ_b^0 and B_c^+ modes	<i>e.g.</i> 8M $B \rightarrow D^*\tau^-\bar{\nu}_\tau$, $\tau^- \rightarrow \mu^-\bar{\nu}_\mu\nu_\tau$ & $\sim 100k \tau^- \rightarrow \pi^-\pi^+\pi^-(\pi^0)\nu_\tau$	Additional sensitivity expected from low- p tracking.
$B_s^0, B^0 \rightarrow \mu^+\mu^-$ $R \equiv \mathcal{B}(B^0 \rightarrow \mu^+\mu^-)/\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-)$; $\tau_{B_s^0 \rightarrow \mu^+\mu^-}$; CP asymmetry	Uncertainty on $R \approx 20\%$ Uncertainty on $\tau_{B_s^0 \rightarrow \mu^+\mu^-} \approx 0.03$ ps	
LFV τ decays $\tau^- \rightarrow \mu^+\mu^-\mu^-$, $\tau^- \rightarrow h^+\mu^-\mu^-$, $\tau^- \rightarrow \phi\mu^-$	Sensitive to $\tau^- \rightarrow \mu^+\mu^-\mu^-$ at 10^{-9}	Phase-II ECAL valuable for background suppression.
CKM tests γ with $B^- \rightarrow DK^-, B_s^0 \rightarrow D_s^+K^-$ etc. ϕ_s with $B_s^0 \rightarrow J/\psi K^+K^-, J/\psi\pi^+\pi^-$ $\phi_s^{s\bar{s}s}$ with $B_s^0 \rightarrow \phi\phi$ $\Delta\Gamma_d/\Gamma_d$ Semileptonic asymmetries $a_{sl}^{d,s}$ $ V_{ub} / V_{cb} $ with Λ_b^0, B_s^0 and B_c^+ modes	Uncertainty on $\gamma \approx 0.4^\circ$ Uncertainty on $\phi_s \approx 3$ mrad Uncertainty on $\phi_s^{s\bar{s}s} \approx 8$ mrad Uncertainty on $\Delta\Gamma_d/\Gamma_d \sim 10^{-3}$ Uncertainties on $a_{sl}^{d,s} \sim 10^{-4}$ <i>e.g.</i> 120k $B_c^+ \rightarrow D^0\mu^-\bar{\nu}_\mu$	Additional sensitivity expected in CP observables from Phase-II ECAL and low- p tracking. Approach SM value. Approach SM value for a_{sl}^d . Significant gains achievable from thinning or removing RF-foil.
Charm CP -violation studies with $D^0 \rightarrow h^+h^-$, $D^0 \rightarrow K_s^0\pi^+\pi^-$ and $D^0 \rightarrow K^\mp\pi^\pm\pi^+\pi^-$	<i>e.g.</i> $4 \times 10^9 D^0 \rightarrow K^+K^-$; Uncertainty on $A_\Gamma \sim 10^{-5}$	Access CP violation at SM values.
Strange Rare decay searches	Sensitive to $K_s^0 \rightarrow \mu^+\mu^-$ at 10^{-12}	Additional sensitivity possible with downstream trigger enhancements.

LHC timeline

LHC roadmap: according to MTP 2016-2020 V1

LS2 starting in 2019 => 24 months + 3 months BC
 LS3 LHC: starting in 2024 => 30 months + 3 months BC
 Injectors: in 2025 => 13 months + 3 months BC



Amount of data in HEP

