


Anomalous magneto-hydrodynamics and toward a full solution of complete relativistic Boltzmann equation on GPUs

反常磁流体力学和 利用GPU数值求解完整版Boltzmann方程

浦 实
中国科学技术大学


第十七届全国核物理大会暨第十三届会员代表大会
2019年10月8-12日 湖北武汉 华中师范大学

Outlines




**Heavy ion
collisions**

macroscopic



**1. Anomalous
magneto-
hydrodynamics**

microscopic



**2. Solving
Boltzmann
equations on
GPUs**



1. Anomalous MagnetoHydroDynamics

Ref: Irfan Siddique, Ren-jie Wang, Shi Pu, and Qun Wang,
arXiv: 1904.01807, accepted by PRD

Strong Electromagnetic fields


- Theoretical estimation:

Lienard-Wiechert potential + Event-by-event





A. Bzdak, V. Skokov PRC 2012 ;W.T. Deng, X.G. Huang PRC 2012;V.Roy, SP, PRC 2015;
H. Li, X.I. Sheng, Q.Wang, 2016; etc. / review: K. Tuchin 2013

Chiral Magnetic Effect



- Magnetic fields
- Nonzero axial chemical potential
- Number of Left handed fermions \neq Number of Right handed fermions





- Charge current: charge separation

$$\mathbf{j} = \frac{e^2}{2\pi^2} \mu_5 \mathbf{B},$$

Kharzeev, Fukushima, Warrigna, (08,09), etc. ...

Relativistic Heavy ion collisions

Experiments: signal VS background



STAR PRL 103, 251601(2009);
PRC 81, 054908

CMS PRL 118, 122301 (2016);
PRC 97, 044912

Anomalous MagentoHydroDynamics

- Conservation equations :

- Energy-momentum conservation

$$\partial_\mu T^{\mu\nu} = 0, \quad T^{\mu\nu} = T_F^{\mu\nu} + T_{EM}^{\mu\nu}.$$

Fluid part **Electromagnetic part**

- (anomalous) currents conservation

$$\partial_\mu j_e^\mu = 0, \quad \partial_\mu j_5^\mu = -e^2 C E \cdot B,$$

Electric Charge current

Chiral current

- Maxwell' s equation :

$$\partial_\mu F^{\mu\nu} = j_e^\nu, \quad \partial_\mu (\epsilon^{\mu\nu\alpha\beta} F_{\alpha\beta}) = 0.$$

Previous Studies: ideal MHD without CME

Preview studies:

➤ 1+1 D ideal MHD Bjorken flow

V.Roy, SP, L. Rezzolla, D. Rischke, Phys.Lett. B750, 45

– With magnetization effects

SP, V. Roy, L. Rezzolla, D. Rischke, Phys.Rev. D93, 074022

➤ 2+1 D ideal MHD Bjorken flow (perturbative)

SP, Di-Lun Yang, Phys.Rev. D93, 054042

➤ Background Magnetic field: contribution to v2

V.Roy, SP, L. Rezzolla, D. Rischke, Phys.Rev. C96, 054909

Problem:

How to add the CME to Magentohydrodynamics?

Before solving Anomalous MHD

- Our previous studies on analytic MHD provides a way to test MHD codes.
- The Italy-Frankfurt group, who has written a MHD code, tested their codes by our analytic solutions.
- Could we find analytic solution for anomalous MHD?
- Or, could we get some hints from analytic solutions?

Ideal limit of MHD

- Ideal limit:

- Electric conductivity is infinite

$$\sigma \rightarrow \infty$$

$$\mathbf{j} = \sigma(\mathbf{E} + \mathbf{v} \times \mathbf{B}) \xrightarrow{\sigma \rightarrow \infty} \mathbf{E} + \mathbf{v} \times \mathbf{B} = 0$$

Maxwell's
equation

$$\partial_t \mathbf{B} = -\nabla \times \mathbf{E} = -\nabla \times (\mathbf{v} \times \mathbf{B})$$

- No space for the CME

$$\nabla \times \mathbf{B} = \mathbf{j} + \partial_t \mathbf{E}$$

- Anomalous MHD needs finite conductivity

Constitution Eqs. for Anomalous MHD

$$\partial_\mu T^{\mu\nu} = 0,$$

$$T^{\mu\nu} = (\varepsilon + p + E^2 + B^2)u^\mu u^\nu - (p + \frac{1}{2}E^2 + \frac{1}{2}B^2)g^{\mu\nu} - E^\mu E^\nu - B^\mu B^\nu - u^\mu \epsilon^{\nu\lambda\alpha\beta} E_\lambda B_\alpha u_\beta - u^\nu \epsilon^{\mu\lambda\alpha\beta} E_\lambda B_\alpha u_\beta,$$

$$\partial_\mu j_e^\mu = 0,$$

$$\partial_\mu j_5^\mu = -e^2 C E \cdot B,$$

Electric
Conducting
flow CME

$$j_e^\mu = n_e u^\mu + \sigma E^\mu + \xi B^\mu,$$

$$j_5^\mu = n_5 u^\mu + \sigma_5 E^\mu + \xi_5 B^\mu,$$

$$\partial_\mu F^{\mu\nu} = j_e^\nu,$$

$$\partial_\mu (\epsilon^{\mu\nu\alpha\beta} F_{\alpha\beta}) = 0.$$

CSE

$$F^{\mu\nu} = E^\mu u^\nu - E^\nu u^\mu + \epsilon^{\mu\nu\alpha\beta} u_\alpha B_\beta,$$

Equation of States (EoS)

- High (chiral) chemical potential:

Energy density $\varepsilon = c_s^{-2} p$, pressure

$$n_e = a\mu_e(\mu_e^2 + 3\mu_5^2),$$
$$n_5 = a\mu_5(\mu_5^2 + 3\mu_e^2),$$
Chemical potential
Chiral chemical potential

- High temperature:

$$\varepsilon = c_s^{-2} p,$$
$$n_e = a\mu_e T^2,$$
temperature
$$n_5 = a\mu_5 T^2,$$

Bjorken boost invariance

- Profound Bjorken velocity

$$u^\mu = \gamma(1, 0, 0, z/t),$$

- Bjorken invariance: all observed quantity are independent on rapidity.
- Could the Bjorken velocity hold in electromagnetic fields?

Simplification

- Neutral fluid:
 - Electric field will not accelerate the fluid
- Force-free-like magnetic field:
- Configuration of Electromagnetic fields:

$$E^\mu = (0, 0, \chi E(\tau), 0), \quad B^\mu = (0, 0, B(\tau), 0),$$

$$\chi = \pm 1 \quad \text{τ: proper time}$$

$$E^\mu = F^{\mu\nu} u_\nu, \quad B^\mu = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} u_\nu F_{\alpha\beta},$$

**Not EM fields
In lab frame!**

Results: High temperature EoS

- Analytic solutions: (at the order of \hbar):

➤ chiral density

$$n_5(\tau) = n_{5,0} \left(\frac{\tau_0}{\tau} \right) \{ 1 + a_2 e^{\sigma \tau_0} [E_1(\sigma \tau_0) - E_1(\sigma \tau)] \},$$

➤ Energy density



τ : proper time
 σ : electric conductivity

$$\begin{aligned} \varepsilon(\tau) = \epsilon_0 \left(\frac{\tau_0}{\tau} \right)^{1+c_s^2} & \left\{ 1 + \sigma \frac{E_0^2}{\varepsilon_0} e^{2\sigma \tau_0} [\tau_0 E_{1-c_s^2}(2\sigma \tau_0) - \tau \left(\frac{\tau}{\tau_0} \right)^{c_s^2-1} E_{1-c_s^2}(2\sigma \tau')] \right. \\ & \left. + \frac{a_3}{\tau_0} e^{\sigma \tau_0} [\tau_0 E_{2-3c_s^2}(\sigma \tau_0) - \tau \left(\frac{\tau_0}{\tau} \right)^{2-3c_s^2} E_{2-3c_s^2}(\sigma \tau)] \right\}. \end{aligned}$$

a1,a2,a3 are related to the initial EM fields and chirality density

$E_n(x)$: the generalized exponential integral. $E_n(z) \equiv \int_1^\infty dt t^{-n} e^{-zt}$

Analytic solution VS numerical results



Solid line: numerical results / Dashed line: analytic

$$a_1 = eC\chi \frac{B_0 n_{5,0}}{a T_0^2 E_0} \tau_0,$$


$$a_2 = \frac{e^2 C \chi E_0 B_0}{n_{5,0}} \tau_0,$$

C: chiral anomaly

Coefficients = $\hbar/(2\pi^2)$

$$a_3 = \frac{eC\chi}{a} \frac{n_{5,0} E_0 B_0}{\epsilon_0 T_0^2} \tau_0.$$

Electromagnetic fields in the Lab frame



$$\mathbf{E}_L = (\gamma v^z B(\tau), \chi \gamma E(\tau), 0),$$

$$\mathbf{B}_L = (-\gamma v^z \chi E(\tau), \gamma B(\tau), 0),$$

τ: proper time
σ: electric conductivity

$$E(\tau) = E_0 \left(\frac{\tau_0}{\tau} \right) \left\{ e^{-\sigma(\tau-\tau_0)} - a_1 e^{-\sigma\tau} [E_{1-2c_s^2}(-\sigma\tau_0) - \left(\frac{\tau}{\tau_0} \right)^{2c_s^2} E_{1-2c_s^2}(-\sigma\tau)] \right\},$$

$$B(\tau) = B_0 \frac{\tau_0}{\tau},$$

$$a_1 = eC\chi \frac{B_0 n_{5,0}}{a T_0^2 E_0} \tau_0,$$

Decay behavior in the Lab frame:

- By decays $\sim 1/\tau$
- Bx decays $\sim \exp(-\sigma\tau)/\tau$

Much slower than decaying in vacuum

Why Bz and Ez vanish?

- We have checked the Maxwell's eq. in Lab frame.
Space and time derivatives of Bz, Ez are zero.
- Key point: the currents is different with static case!

$$\nabla \times \mathbf{B}_L = \mathbf{j}_e + \partial_t \mathbf{E}_L.$$
$$\mathbf{j}_{e,\parallel} = \sigma \mathbf{E}_{L,\parallel} + \xi \mathbf{B}_{L,\parallel}, \quad v: \text{three vector of fluid velocity}$$

$$\mathbf{j}_{e,\perp} = \sigma \gamma (\mathbf{E}_L + \mathbf{v} \times \mathbf{B}_L)_{\perp} + \xi \gamma (\mathbf{B}_L - \mathbf{v} \times \mathbf{E}_L)_{\perp},$$

- Similar to the force-free EM fields in classical electrodynamics (e.g. Woltjer states)

Qin, Liu, Li, Squire, PRL 109, 235001 (2012);
Xia, Qin, Q. Wang, PRD(2016)

Summary of Anomalous MHD

- Anomalous MHD:
Hydrodynamic eq. + Maxwell' s eq. + Chiral currents
- Analytic solutions of anomalous MHD in Bjorken flow with transverse EM fields
- Decay behavior of EM fields:
 - In lab frame, B field decays much **slower** than in the vacuum
 - By decays like $\sim 1/\tau$,
 - B_x decays like $\sim \exp(-\sigma \tau)/\tau$

2. Toward a full solution of complete relativistic Boltzmann equation on GPUs


Ref: Jun-jie Zhang, Hong-zhong Wu, SP, Guang-you Qin,
Qun Wang, in preparation

Relativistic Boltzmann Equations (BE)

- Boltzmann equations: kinetic theory
 - microscopic theory, many body-systems
- Many many applications in non-equilibrium systems:
 - Turbulence, self-similar behavior
 - Simulations for heavy ion collisions
 - Gluon condensation

Relativistic BE

$$\frac{\partial}{\partial t} f_p + \frac{\partial \mathbf{x}}{\partial t} \cdot \nabla_x f_p + \frac{\partial \mathbf{p}}{\partial t} \cdot \nabla_p f_p = C[f_p]$$




f: distribution function = function of (t,x,p)

- It is a semi-classical description of many body theory.
One can derive it symmetrically from quantum field theory with closed-time-path formulism.
(Blaizot, Iancu, Phys. Rep. 359, 355 (2002).)

Collisional term

$$C_{ab \rightarrow cd} \equiv \int \prod_{i=1}^3 \frac{d^3 k_i}{(2\pi)^3 2E_{k_i}} \frac{|M_{ab \leftrightarrow cd}|^2}{2E_p} (2\pi)^4 \delta^{(4)}(k_1 + k_2 - k_3 - p) [f_{k_1}^a f_{k_2}^b F_{k_3}^c F_p^d - F_{k_1}^a F_{k_2}^b f_{k_3}^c f_p^d],$$

For 2->2 scatterings including quarks and gluons: e.g.



Main difficulties for relativistic BE

$$\frac{\partial}{\partial t} f_p + \frac{\partial \mathbf{x}}{\partial t} \cdot \nabla_x f_p + \frac{\partial \mathbf{p}}{\partial t} \cdot \nabla_p f_p = C[f_p]$$

- 6+1 dimensional phase space
 - (3 coordinates, 3 momentum, 1 time)
- Extremely complicated collisional term
 - 5 dimensional integrals including many terms

Assuming we only have 10 grids in all space and momentum directions, so totally, we have 10^6 grids. We choose the time step could be 1000. Totally, we need to compute 10^9 times high dimensional integrals. Assuming that it costs 1 sec to compute one 5 dim integral, then totally it will cost 30 years!

Other issues in relativistic BE

- Particle number non-conservation:
comes from errors of collisional integrals.
- A usual way: Test particle method;
requires many parameters.

Could we solve a complete BE directly?
Or, with minimal parameters?

What have we done?

- A new numerical framework on GPUs :
 - A full solution of complete relativistic BE
 - High performance
 - Particle number is strictly conserved


Parameters

- Physical parameters:
 - coupling constant; initial conditions.
- Parameters for simulations:
 - Size, number of grids.
- Minimal parameters !

Collisional term via ZMCintegral


- 5 dimensional integral on each phase space grid:

Wu, Zhang, Pang, Q. Wang, 1902.07916. accepted in Computer Physics Communications




Symmetrical sampling method on GPUs

We introduce a new method to **ensure particle number conservation**.



Time evolution


Pure quark case



- space: 1 grid; momentum: $30 \times 30 \times 30 = 27,000$
- Phase space box is of size $[-3\text{fm}, 3\text{fm}]^3 \times [-2\text{GeV}, 2\text{GeV}]^3$.
- Time step $dt=0.001\text{fm}$; 20,000 steps
- on one Nvidia Tesla V100 card: costs around 2 hours

Time evolution

Gluons + quarks



- space: 1 grid; momentum: $30 \times 30 \times 30 = 27,000$
- Phase space box is of size $[-3\text{fm}, 3\text{fm}]^3 \times [-2\text{GeV}, 2\text{GeV}]^3$.
- Time step $dt=0.0005\text{fm}$; 100,000 steps
- on one Nvidia Tesla V100 card: costs around 50 hours


Rich phenomena

- Gluon condensation
- Effects of inelastic scatterings
- Turbulence, self-similar behavior,
non-thermal fixed points

Summary for BE


- We introduce a new numerical framework to derive full solutions of a complete relativistic BE on GPUs.
- Full collisional term: high dimensional integrals.
- High performance:
space 10x10x10, momentum 30x30x30, Time steps: 10^4 - 10^6 ,
on one Nvidia Tesla V100 card costs a few days!
- Particle number is strictly conserved.

Summary




**Heavy ion
collisions**

macroscopic



**1. Anomalous
magneto-
hydrodynamics**


microscopic



**2. Solving
Boltzmann
equations on
GPUs**


Thank you for your time!

Time evolution




Energy conservation

- Not conserved in HTL (physically)
- Not conserved (from errors): because of discrete grid



- But, if we increases the number of grid, the variation of energy is tiny.

Symmetric Sampling



$$d\Gamma_{ab \rightarrow cd} = \frac{1}{2E_p} |M_{ab \rightarrow cd}|^2 \prod_{i=1}^3 \frac{d^3 k_i}{(2\pi)^3 2E_{k_i}} \\ \times (2\pi)^4 \delta^{(4)}(k_1 + k_2 - k_3 - p)$$

$$\frac{d\tilde{f}_p^a(x)}{dt} = \mathcal{C}_a$$

5维积分: $k_{1y}, k_{1z}, k_{3x}, k_{3y}, k_{3z}$
 6维扫描: p_x, p_y, p_z, x, y, z


一维时间尺度迭代

即使利用ZMCintegral, 将5维积分压缩到5s, 6维 ($50 \times 25 \times 25 \times 5 \times 5 \times 5$) 扫描需要 $\sim 10^6$ s, 加上时间维度共需要 $\sim 10^9$ s ~ 30 年! !


结合ZMCintegral编程经验, 修改GPU内核程序。将5维积分压缩到 ~ 0.01 s, 6维扫描压缩至 $\sim 10^2$ s, 有希望将时间控制在 $\sim 10^5$ s ~ 1 天。Challenging! !

Chirality and massless fermions

Left handed



Right handed



Polarization by magnetic fields

